

	1
Breast Cancer Molecular Subtypes in Relation to Age, Stage and Grade among Sudanese Women Patients in Khartoum specialized oncology center (2013 – 2017)	3
	4
Abstract	5
Background: -	6
There is a lack of data regarding molecular subtypes of breast cancer among Sudanese female patients. Moreover to the best of our knowledge no study had correlated the molecular subtypes with age, stage and grade till date. This study aimed to determine breast cancer molecular subtypes among Sudanese women in relation to age, clinical stage and grade and to compare the findings to other related researches.	11
Methods: Retrospective Study of (255) Sudanese female patients diagnosed with breast cancer from September 2013 to August 2017 in Khartoum Specialized Oncology Center, an information sheet was used to collect data from patients medical records.	14
Results: The results demonstrated that, the most commonly detected molecular subtype was luminal B (34.9%), followed by triple negative and HER-2 enriched, (31.4%) and (19.2%) respectively. The least common subtype was luminal A, (14. 5%). Additionally, over half of the patients (54.4%) were classified as grade 3 and (22.4%) were diagnosed as stage IIIb.	18
Conclusion: The main conclusion that can be drawn is that, luminal B was the most commonly diagnosed subtype and it was more common in younger age group in comparison to luminal A. The majority of the patients in the study were grade 3 and stage IIIb and were diagnosed between the age of 41 and 50. Moreover, triple negative subtype was high among Sudanese women and was associated with advance grade breast cancer.	23
Key words: Subtypes, Breast Cancer, Sudanese, Women	24
	25
	26
	27
	28
	29
	30

2 1. Introduction:

3 Breast cancer is the top cancer in females leading to 327,000 deaths annually [1, 2]. In 2018,
 4 WHO reported nearly 2.1 million newly diagnosed female with breast cancer [3]. The global
 5 cancer burden of breast cancer is 11.6% [4]. According to the latest data of Globocan 2018,
 6 breast cancer ranked the most frequent cancer among women in Sudan and a total of 5,677
 7 (36.69%) Sudanese women were newly diagnosed with breast cancer [5].
 8 Several clinical and pathological parameters are used to classify the breast cancer subtypes,
 9 namely; lymph node (LN) status, tumor size, tumor grade, age, menopausal status, clinical and
 10 pathological stage, histological type as well as estrogen (ER), progesterone (PR) and HER-2
 11 receptors [6]. Detecting the molecular subtype of breast cancer is a recent and advance technique
 12 for early detection, evaluating the prognosis and management of breast cancer [7]. Thus,
 13 management options in Sudan are adapted according to breast cancer molecular subtype. The
 14 followed management treatment guidelines in Sudan are based on the National Comprehensive
 15 Cancer Network (NCCN) and European Society for Medical Oncology(ESMO) guidelines which
 16 includes; screening, diagnosing, staging, management of local/loco- regional disease, endocrine
 17 treatment in pre- and post-menopausal patients, chemotherapy and HER2-directed therapy [8, 9].
 18 Undoubtedly molecular subtypes could provide promising prognostic and predictive information
 19 and may help identify new therapeutic targets. However, it is important to understand their
 20 limitations and to evaluate their role in improving breast cancer prognosis beyond the traditional
 21 expected outcomes in a practical and cost-effective manner [10].

Breast cancer is divided into four major molecular subtypes; the most frequent type is luminal A
 which is present in 50-60% of breast cancer patients and has the best prognosis rates. The second
 subtype, luminal B is characterized by higher histological grade and is found in 15-20% of 24
 patients [11]. The third subtype triple negative breast cancer (TNBC) which varies according to
 patient race and ethnicity. Furthermore, it impacts patient's survival rates and may affects 26
 management options [12]. The fourth molecular subtype, HER2-enriched (previously the 27
 HER2+/ER- subtype) is characterized by aggressive behavior [13]. There is a variation in the
 prevalence of molecular subtypes worldwide. Luminal A tumors were the most common tumors
 among some Middle East countries, namely; Saudi Arabia [14], Jordan [15] and Egypt [16] and
 in a number of western countries; Italy [17], Germany [18] and Atlanta (America) [19]. 31

Regarding Asian countries, studies revealed that luminal A was high among Iranian patients [20], while Luminal B was more commonly found in Japanese [21] and Pakistani patients [22]. Studies in African countries unveiled that in Morocco (North Africa), Luminal B was the most common subtype [23], while triple negative had high prevalence rates in Nigeria (West Africa) [24], Uganda (sub-Saharan Africa) [25], Sudan and Eretria (North East and East Africa) [26]. Regarding race, TNBC was found to be the most prevalent subtype in African-American women [27]. Another study conducted to compare molecular subtypes between Sudanese and German women showed that triple-negative subtype was more frequent in Sudanese than German women [28].

Few studies have investigated breast cancer molecular subtypes in Sudan. Furthermore, there is no research relating molecular subtypes to age, stage and grade of breast cancer to date. Thus, in our research we tried to classify patients according to breast cancer molecular subtypes.

The aim of this study is to determine breast cancer molecular subtype among Sudanese women in relation to age, clinical stage and grade and to compare the results to other related researches.

2. Materials and Methods

2.1 Study design and setting

This is a retrospective study of histologically confirmed Sudanese women with breast at Khartoum Specialized Oncology center, in the period from September 2013 to August 2017. Khartoum Specialized Oncology center is a specialized tertiary hospital that offers chemotherapy and radiotherapy for cancer patients, located in Khartoum State, Capital of Sudan. Entitled patients are referred from all over Sudan and hence, this study's sample is therefore representative of the Sudanese population.

2.2 Sampling

255 medical records of histologically confirmed breast cancer patients were included in the study.

2.3 Data collection tools

An information sheet has been used for data collection from patient's medical records.

The data retrieved included the following:-

-Patients age when diagnosed was distributed into two groups; younger age group (50 years or less) and older age group (above 50 years).

1 -Molecular subtypes which were identified by Immunohistochemical (IHC) markers
2 (ER/PR/HER2). Four subtypes were defined, namely; luminal A, luminal b, triple negative and
3 HER-2 enriched.

4 -Breast cancer grades (I, II and III) were detected using Nottingham Bloom-Richardson grading
5 system [29].

6 -The Clinical stage of the disease was estimated from the clinical examination and was classified
7 according to American Joint Committee (AJC) and TNM classification [30].

8 **2.4 Statistical analysis**

9 Data was entered and analyzed using the Statistical Package for the Social Sciences (version
10 21.0). Chi-square Test was used to evaluate the correlation between molecular subtypes and age,
11 stage and grade were used. The results were considered significant when p (degree of
12 significance) was less than 0.05.

13 **2.5 Ethical consideration**

14 Ethical approval was obtained from Institutional review board of Omdurman Islamic university-
15 Faculty of Medicine. Data were collected after taking the necessary agreement from Khartoum State
16 Ministry of Health and Khartoum Specialized Oncology Center.

17 **3. Results:**

18 A total of 255 records of female patients diagnosed with breast cancer were enrolled in the study.
19 The mean patient's age at diagnosis was 48.8 ± 11.3 years. The majority of the patients (78.2%)
20 were diagnosed with breast cancer before the age of 60. However, only (20.8%) of the cases
21 were diagnosed above the age of 60 (Table 1).

22 The most commonly detected molecular subtype was luminal B (34.9%), followed by triple
23 negative and HER-2 enriched, (31.4%) and (19.2%), respectively. The least common subtype
24 was luminal A (14.5%) (Fig.2). The vast majority of patients (22.4%) were stage IIIb, followed
25 by stage IIa and IIb (21.1%). Furthermore, (15.7%), (8.6%) and (6.7%) of the cases were
26 diagnosed as stage IIIa, IV and I, respectively. A small minority, (4.3%) were diagnosed as stage
27 IIIc (Table 2). (54.4%) of the breast cancer patients were diagnosed as grade 3, while (39.2%)
28 were classified grade 2 and only (9.4%) were diagnosed as grade 1 (Fig. 1).

29 Out of the (255) cases, 135) cases (52.9%) were in the younger age group (≤ 50) and 120 cases
30 (47.1%) were in the older age group (>50 years). Most cases (34.9%) (n=89) were classified as
31 luminal B subtype, (51.7%) of which were in the younger age group, while (48.3%) (n=43) were

1 in the older age group. Moreover, (31.4%)(n=80) of the cases were classified as triple negative
2 subtype, (56.3%) (n=45) were in the younger age group and (47.7%) were in the older age
3 group. (14.5%) (n=37)of the cases were classified as luminal A subclass, (37.8%) (n=14) were in
4 the younger age group and (62.7%)(n=23) were in the older age group. (19.2%)(n=49) were
5 classified as HER-2 enriched subclass, (61.2%)(n=30) were in the younger age group and
6 (38.8%) (n=19) were in the older age group. However, the relationship between molecular
7 subtype of breast cancer and patients age at diagnosis was not statistically significant (p=0.162)
8 (Table 3).

9 Concerning the relationship between molecular subtypes and breast cancer stages, there was no
10 significant association (p=0.257). For patients with HER-2 enriched molecular subtypes, the
11 frequency of stage I, IIa, IIb IIIa, IIIb, IIIc and IV breast cancer, were
12 (2%),(16%),(24.4%),(18.3%),(26.5),(2%),and (10.2%), respectively. Regarding patients with
13 Luminal A molecular subtypes, stage I, IIa, IIb IIIa, IIIb, IIIc and IV breast cancer, were
14 (16%),(32%),(18.9%),(10.8%),(16.2%) and (2.7%), respectively. For patients with Luminal B
15 molecular subtypes, stage I, IIa, IIb IIIa, IIIb, IIIc and IV breast cancer, were
16 (5.6%),(22.4%),(19%),(13.4%),(22.4%),(3.4%) and (13.4%), respectively. For patients with
17 Triple negative molecular subtypes, stage I, IIa, IIb IIIa, IIIb, IIIc and IV breast cancer, were
18 (6.3%),(17.5%)(22.5%),(18.8%),(22.4%),(7.5%)and (5%), respectively (Table 4).

19 The molecular subtypes were found to be significantly associated with breast cancer grade
20 (p=0.012). Luminal B frequency of grade 1, 2 and 3 was (3.5%), 16.1%) and (15.3%),
21 respectively, while the distribution of Luminal A was (2.4%) for grade 1, (7.8%) for grade 2 and
22 (4.3%) for grade 3. (19.2%) of triple negative were grade 3, (9.8%) were grade 2, while only
23 (2.4%) were grade 1. (12.5%) of Her-2 enriched were grade 3, (5.5%) were grade 2 and (1.2%)
24 were grade 1.

25 **4: Discussion**

26 In the current study most of the patients were above or fifty years and above (52.9%), while
27 patients under fifty were (47. 1%). Furthermore, the vast majority of patients were diagnosed
28 with breast cancer between the age of 41 and 50. A similar conclusion was suggested by a study
29 done in Nigeria [24]. According to the present study most of the cases were grade 3(54.4%) and
30 (39.9%) were grade 2. However, a different finding was reported by another study, which

1 revealed that grade 2 breast cancer was more frequent among Sudanese, German and Nigerian
2 female patients, (54.6%), (60%) and (48.57%) respectively, while the percentage of stage 2
3 cancer in Sudanese women was (41.8%), German women (22%) and (43.57%) for Nigerian
4 women. The latter finding could be partially explained by the late detection of cancer cases in
5 Sudan.

6 From the results, (22.4%) of female patients were diagnosed as stage IIIb cancer, that may be
7 attributed to lack of awareness, difficult accessibility services and absence of cancer screening
8 programs.

9 With regard to molecular subtypes, luminal A is found in (50%-60%) of the patients and luminal
10 B in (15%-20%), however, in our group the majority of the cases were classified as luminal B
11 (34.9%) and only (14.5%) were luminal A [11]. A different conclusion was obtained from a
12 study done in middle east countries; where percentage of patients with luminal A subtype in
13 Saudi Arabia, Jordan and Egypt was (58.5%), (60%) and (45%,) respectively [14,15 and 16].
14 Another similar study conducted in some western countries showed the following; Italy (34%),
15 Germany (44.7%) and Atlanta (51.1%), [17, 18 and 19]. Furthermore, the prevalence of Luminal
16 A subtype was (63.8%) according to an Iranian study [20]. In Japan, Pakistan and Morocco, the
17 percentage of luminal B was (71%), (69%) and (41.8%), respectively [21, 22 and 23] which was
18 higher than our finding. The variation in the results could be linked to the distribution of the
19 different age groups in the studies.

20 The prevalence of Triple negative breast cancer subtype (TNBC) in Nigerian women was
21 (26.53%) and (21.2%) among African-American women [24, 27], nonetheless, a lower result
22 was attained by our study (31.4%). A slightly comparable value to our finding was found in
23 Ugandan women (34%) and Sudanese- Eritrean women (34.5%), [25, 26 and 28].

24 In our examination, HER-2 enriched was found in (19.2%) of the breast cancer patients.
25 Nevertheless, a lower finding was cited by another study; the HER-2 enriched frequency among
26 Jordanian women was (12%) [15], Sudanese women (15.7%), German women (6.8%) [28],
27 Sudanese Eritrean women (16%) and (9.2%) among Moroccan women [23, 26].

28 Luminal B subtype was present in (51.7%) of patients fifty years or less. whereas, the percentage
29 in older patients (i.e. above fifty) was (48.3%). On the other hand, Luminal A subtype prevailed
30 in patients over fifty years old (62.2%), while (37.8. %) was detected in younger ages (fifty years

1 or less). A similar pattern of result was obtained in Jordan, where (72%) of luminal A subtype
2 were above fifty years old [15].

3 Triple negative and HER-2 enriched subtypes were more prominent in the younger ages (fifty
4 years or less), (56.3%) and (61.2%), respectively. In patients over fifty years old, Triple negative
5 subtype was detected in (43.7%) of the patients and Her-2 enriched in (38.8%). This was
6 consistent with a study done among Sudanese and German women [28].

7 (7.8%) of Luminal B breast cancer subtype was stage IIIb, while only (2.4%) of luminal A was
8 stage IIIb, which indicates that, Luminal B is more aggressive than luminal. With regard to
9 cancer grades, the majority of luminal A breast cancer subtype (7.8%) was Grade 2. On the other
10 hand, in Grade 3, luminal B prevailed over luminal A, (15.3%) and (4.3%) respectively.
11 Therefore, it's suggested that, luminal B is associated with higher histological grades [11].

12 Another finding was that, high frequencies for grade 3 cancer was registered by Triple negative
13 and Her-2 enriched Subtypes (19.2%) and (12.5%), respectively. This implies that Triple
14 negative and Her-2 enriched are associated with aggressive and advanced stages of breast cancer
15 [13].

16 **4: Conclusion and recommendation: -**

17 Most of the Sudanese women were diagnosed with breast cancer between (41-50) years old.

18 Moreover, most of them were presented with grade 3 and stage IIIb breast cancer.

19 Luminal B was the most prevailed molecular subtypes, followed by Triple negative.

20 Luminal A was more common among old age groups (over fifty). However, Her-2 enriched and

21 TNBC subtypes were mostly Grade 3 and prevailed among younger Sudanese women.

22 Detecting the subtype of breast cancer is not only essential for following disease prognosis but
23 also for the management of the breast cancer.

24 Breast cancer screening programs and self-examination are highly recommended for the early
25 detection of the disease.

26 The effect of determining molecular subtype on survival rates is an issue for future research to
27 explore.

1 **References: please the formatting of the references with the journal specifications.**

2
3 1-ZiQi Tao • Aimin Shi • Cuntao Lu • Tao Song • Zhengguo Zhang • Jing Zhao. *Cell Biochem*
4 *Biophys, Breast Cancer: Epidemiology and Etiology, cell Biochem Biophys* (2015) 72:333–338
5 (DOI 10.1007/s12013-014-0459-6).

6 The correct format is: ZiQi Tao, Aimin Shi, Cuntao Lu, Tao Song, Zhengguo Zhang, Jing Zhao.
7 *Breast Cancer: Epidemiology and Etiology. Cell Biochem Biophys* 2015); 72:333–38.

8 2-Hadi MA, Al Madani R, Abu Arida L, Al Ghareeb B. Breast Cancer Age in Developing
9 Countries: The Narrowing Gap. *Clin Surg.* 2018; 3: 2074

10 3- Freddie Bray et al: Global cancer statistics 2018: GLOBOCAN estimates of incidence and
11 mortality worldwide for 36 cancers in 185 countries, *CA CANCER J CLIN* 2018; 68:394–424

12 4-WHO, Latest global cancer data: available at:

13 <https://www.who.int/cancer/PRGlobocanFinal.pdf>, [September, 2018].

14 5- WHO, the Global Cancer Observatory, Available at:

15 <http://gco.iarc.fr/today/data/factsheets/populations/729-sudan-fact-sheets.pdf> [March, 2019].

16
17 6 I Gannon LM, Cotter MB, Quinn CM. The classification of invasive carcinoma of the breast.
18 *Expert Rev Anticancer Ther* 2013; 13: 941-954. (PMID: 23984896).

19

20

21 7-Jane Date C Hon, Baljit Singh, Aysegul Sahin, Gang Du, Jinhua Wang, Vincent Y Wang,
22 Fang-Ming Deng, David Y Zhang, Marie E Monaco, Peng Lee *Am J Cancer Res.* 2016; 6(9):
23 1864–1872. Published online 2016 Sep 1. PMID: PMC5043099.

24
25 8-National Comprehensive Cancer Network® (NCCN®), Guidelines for Breast Cancer Patients
26 Version 4.2018, available at www.nccn.org/patients[February, [February, 8, 2019]

27
28 9- E. Senkus, S. Kyriakides, S. Ohno, et al. Primary breast Cancer, ESMO Clinical Practice
29 Guidelines for diagnosis, treatment and follow-up, *Annals of Oncology*, Volume 26, Issue
30 suppl_5, September 2015, Pages v8–v30, <https://doi.org/10.1093/annonc/mdv298>

31
32 10-Emad A Rakha, Jorge S Reis-Filho, *et al.* Breast cancer prognostic classification in the
33 molecular era: the role of histological grade *Breast Cancer Res.* 2010; 12(4): 207. Published
34 online 2010 Jul 30. doi: 10.1186/bcr2607 PMID: PMC2949637.

35
36 11- Ozlem Yersal, Sabri Barutca *World Biological subtypes of breast cancer: Prognostic and*
37 *therapeutic implications, J Clin Oncol.* 2014 Aug 10; 5(3): 412–424. Published online 2014 Aug
38 10. doi: 10.5306/wjco.v5.i3.412.

- 1 12- Ding YC, Steele L et al, Molecular subtypes of triple-negative breast cancer in women of
2 different race and ethnicity, *Oncotarget*. 2019 Jan 4; 10(2):198-2
3 doi:10.18632/oncotarget.26559.
- 4 13- O'Sullivan C.C., Loprinzi C.L., Haddad T.C; Updates in the Evaluation and Management of
5 Breast Cancer, *Mayo Clinic Proceedings*, 2018, 93 (6), pp. 794-807
6 DOI: <https://doi.org/10.1016/j.mayocp.2018.03.025>.
- 7
- 8 14. Al Tamimi DM, Shawarby MA, Ahmed A, Hassan AK, AlOdaini AA. Protein expression
9 profile and prevalence pattern of the molecular classes of breast cancer—a Saudi population
10 based study. *BMC Cancer*. 2010; 10(1):223. <https://doi.org/10.1186/1471-2407-10-223> PubMed:
11 20492711 Pied: PMC2880995.
- 12
- 13 15-Shomaf, M., Masad, J., Najjar, S., & Faydi, D. (2013). Distribution of breast cancer subtypes
14 among Jordanian women and correlation with histopathological grade: molecular sub
15 classification study. *JRSM Short Reports*, 4(10). <https://doi.org/10.1177/2042533313490516>.
- 16 16-Bodour Sahlia , Coya Tapia et al. Molecular subtype analysis determines the association of
17 advanced breast cancer in Egypt with favorable biology, *BMC Women's Health*, 2011, 11:44,
18 <https://doi.org/10.1186/1472-6874-11-44>.
- 19
- 20 17-Caldarella A, Buzzoni C, Crocetti E, Bianchi S, Vezzosi V, Apicella P, Biancalani M,
21 Giannini A, Urso C, Zolfanelli F, Paci E, Invasive breast cancer: a significant correlation
22 between histological types and molecular subgroups, *J Cancer Res Clin Oncol*. 2013 Apr;
23 139(4):617-23.
- 24
- 25 18- Fabian Riedel, Peter Sinn, Prognosis of breast cancer molecular subtypes in routine clinical
26 care: A large prospective cohort study, *BMC Cancer* 2016; 16:734, [https://doi.org/10.1186/s12885-](https://doi.org/10.1186/s12885-016-2766-3)
27 016-2766-3.
- 28 19-Gaudet MM, Press MF, et al Risk factors by molecular subtypes of breast cancer across a
29 population-based study of women 56 years or younger, *Breast Cancer Res Treat*. 2011 Nov;
30 130(2):587-97. doi: 10.1007/s10549-011-1616-x. Epub 2011 Jun 11.
- 31 20- Kadivar M, Mafi N, Joulaee A, Shamshiri A, Hosseini N. Breast cancer molecular subtypes
32 and associations with clinicopathological characteristics in Iranian women 2002-2011. *Asian Pac*
33 *J Cancer Prev*. 2012; 13:1881–1886. [PubMed] [Google Scholar].
- 34 21-Shibuta K, Ueo H, Furusawa H, Komaki K, Rai Y, Sagara Y, Kamada Y, Tamaki N, The
35 relevance of intrinsic subtype to clinicopathological features and prognosis in 4,266 Japanese
36 Women with Breast Cancer. *Breast Cancer*. 2011; 18(4):292-98. [https://doi.org/10.1007/s12282-](https://doi.org/10.1007/s12282-010-0209-6)
37 010-0209-6 PMid: 20571962. [PubMed] [Google Scholar] .
- 38
- 39 22- Hashmi AA, Aijaz S, Khan SM, Mahboob R, Irfan M, Zafar NI, Nisar M, Siddiqui M, Edhi
40 MM, Faridi N, Khan A. Prognostic parameters of luminal A and luminal B intrinsic breast cancer
41 subtypes of Pakistani patients. *World J.Sur Oncol*. 2018; 16(1):1. <https://doi.org/10.1186/s12957->

- 1 23-El Fatemi H, Chahbouni S, Jayi S, Moumna K, Melhouf MA, Bannani A, Mesbahi
2 O, Amarti A., Luminal B tumors are the most frequent molecular subtype in breast cancer of
3 North African women: an immunohistochemical profile study from Morocco, *Diagn Pathol*; 2012
4 Dec 7; 7:170. doi: 10.1186/1746-1596-7-170.
- 5 24-Kenneth A.Omoruyi1, Martins A. Nnoli1, Godwin A. Ebughe , Godstime . Irabor , Edoise M.
6 Isiwele, Prevalence of Molecular Subtypes of Breast Carcinoma in University of Calabar
7 Teaching Hospital using Immunohistochemistry as surrogates for Intrinsic DNA gene
8 characteristics, *IOSR Journal of Dental and Medical Sciences (IOSR-JDMS)* e-ISSN: 2279-
9 0853, p-ISSN: 2279-0861. Volume 17, Issue 2 Ver. 12 February. (2018), PP 64-73
- 10 25- Moses Galukande, Henry Wabinga, Florence Mirembe, Charles Karamagi, Alexzander Asea,
11 Molecular breast cancer subtypes prevalence in an indigenous Sub Saharan African population
12 *Pan Afr Med J.* 2014; 17: 249. Published online 2014 April 5. doi: 10.11604/pamj.2014.17.249.330
13 PMID: PMC4189896.
- 14
- 15 26- Sengal AT, Haj-Mukhtar NS, Elhaj AM, Bedri S, Kantelhardt EJ, Mohamedani AA.
16 Immunohistochemistry defined subtypes of breast cancer in 678 Sudanese and Eritrean women;
17 hospitals based case series, *BMC Cancer.* 2017 Dec 1; 17(1):804. doi: 10.1186/s12885-017-
18 38054. PMID: 29191181.
- 19
- 20 27- Chukwuemeka U. Ihemelandu et al: Molecular breast cancer subtypes in premenopausal and
21 postmenopausal African-American women: age-specific prevalence and survival. *J Surg*
22 *Res.* 2007 Nov; 143(1): 109–118. doi: 10.1016/j.jss.2007.03.085.
- 23
- 24 28-Sengal AT, Haj Mukhtar NS, Vetter M, Elhaj AM, Bedri S, Hauptmann S, Thomssen C,
25 Mohamedani AA, Wickenhauser C, Kantelhardt EJ. Comparison of Receptor-Defined Breast
26 Cancer Subtypes between German and Sudanese Women: A Facility-Based Cohort Study.
27 *J Glob Oncol.* 2018 Sep; 4:1-12. doi: 10.1200/JGO.2017.010082. Epub 2017 Aug 4.
28 PMID: 30241184.
- 29 29- Elston C, Ellis IO. Pathological prognostic factors in breast cancer: the value of histological
30 grade in breast cancer: experience from a large study with long-term follow-up. *Histopathology.*
31 2002; 41:154–61.
- 32 30-The American Joint Committee on Cancer: the 7th edition of the AJCC cancer staging
33 manual and the future of TNM. Edge SB, Compton CC *Ann Surg Oncol.* 2010 Jun; 17(6):1471-4.
34

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15

Table1- Distribution of patients according to the age of breast cancer at diagnosis.

Age group	Frequency	Percent
<30	7	2.7
30-40	54	21.2
41-50	74	29.0
51- 60	67	26.3
>60	53	20.8
Total	255	100.0

Table 2- Distribution of patients according to the stages of breast cancer at diagnosis.

Stage	N	%
I	17	6.7
IIa	54	21.2
IIb	54	21.2
IIIa	40	15.7
IIIb	57	22.4
IIIc	11	4.3
IV	22	8.6
Total	255	100.0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32




Fig.1- Distribution of patients according to the grade of breast cancer at diagnosis.




Fig.2- Distribution of patients according to the molecular subtypes of breast cancer at diagnosis.

1
2
3
4

Table 3- The relationship between molecular subtype of breast cancer and age at diagnosis.

Molecular subtype	Cases ≤50 years N(%)	Cases >50 years N(%)	Total N(%)	P- Value
HER-2 enriched	30(61.2)	19(38.8)	49(19.2)	0.162
Luminal A	14(37.8)	23(62.7)	37(14.5)	
Luminal B	46(51.7)	43(48.3)	89(34.9)	
Triple negative	45(56.3)	35(47.7)	80(31.4)	
Total	135(52.9)	120(47.1)	255(100)	

5
6
7
8
9
10

Table 4- The relationship between Molecular subtype of breast cancer and stage at diagnosis.

			Stage							Total	P value
			I	II a	II b	III a	III b	III c	IV		
Molecular subtype	HER-2 enriched	N (%)	1 (2)	8 (16)	12 (24.4)	9 (18.3)	13 (26.5)	1 (2)	5 (10.2)	49 (19.2)	0.257
	Luminal A	N (%)	6 (16)	12 (32)	7 (18.9)	4 (10.8)	6 (16.2)	1 (2.7)	1 (2.7)	37 (14.5)	
	Luminal B	N (%)	5 (5.6)	20 (22.4)	17 (19)	12 (13.4)	20 (22.4)	3 (3.4)	12 (13.4)	89 (34.9)	
	Triple negative	N (%)	5 (6.3)	14 (17.5)	18 (22.5)	15 (18.8)	18 (22.5)	6 (7.5)	4 (5)	80 (31.4)	
Total			17 (6.7)	54 (21.2)	54 (21.2)	57 (22.4)	40 (15.7)	11 (4.3)	22 (8.6)	255 (100)	

11

1
2
3

Table 5: The relationship between Molecular subtypes of breast cancer and histological grades.

			Grade			Total	P-Value
			1	2	3		
Molecular subtype	HER-2enriched	N (%)	3 (6.1)	14 (28.6)	32 (65.3)	49 (19.2)	0.012
	Luminal A	N (%)	6 (16.2)	20 (54.1)	11 (29.7)	37 (14.5)	
	Luminal B	N (%)	9 (10.1)	41 (46.1)	39 (43.8)	89 (34.9)	
	Triple negative	N (%)	6 (7.5)	25 (31.2)	49 (61.3)	80 (31.4)	
Total	N (%)	24 (9.4)	100 (39.2)	131 (51.4)	255 (100)		

4

UNDER PELL