

34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67




Fig 1.1




Fig 1.2

Figure 1.1, three major nutrients (Nitrogen, Potash, and Phosphate) and total fertilizer demand globally, predictions, 2015-2019. Figure. 1.2, Regional and worldwide Nitrogen demand forecast, 2015- 2019. (Adopted from FAO 2016 and ²)

Commonly, fertilizers are considered as a vital part to increase the crop's production, but the excessive use of nitrogen can lead to reducing the full potential of crop's output. Moreover, an adequate supply of nitrogen does not become part of the plant's system. This excessive part is leached and cause environmental problems and human health hazards². Only 30 to 50% NUE is recorded in plants, the remaining is used by soil microbes, leached down in soil or volatilized³ Healthy plants retain 2-4% Nitrogen³. Nitrogen plays an important role in the preparation of proteins. In the case of Nitrogen deficiency, plant's growth is stunted². From the previously performed experiments, it has been noted that nitrogen is lost from the soil. In high rain-fed areas and light texture soils (sandy soil), leaching is a common problem.

Nitrate form of N does not strongly absorb on soil surface because nitrate is the mobile in nature, and easily move beyond the soil profile by process of leaching⁴. Through this mechanism, as much as 25-50% of the applied N can be lost⁴. From the soil, N can be lost through the water as well as wind erosion. Loss of N through water erosion is a major problem for humid and sub-humid climatic areas while wind erosion is a more commonly reported mechanism of N loss in the arid and semiarid climatic region⁵. Soil Physical, chemical properties and genotypes cause 18%, 5% and 12% losses, respectively. Due to less fertility, 50% of agricultural lands are not producing the crop with full yield potential. Except this, Low NUE also due to the excessive use of Nitrogen in the fields.

68

69 Excess use of nitrogen declines the crop yield⁶. Different kind of practices is being used to
70 improve the nitrogen use efficiency (NUE). The main objective of NUE is to enhance the
71 performance of the overall cropping system. Nitrogen use efficiency (NUE) also addresses
72 the sustainability of agriculture system with respect to soil fertility and some other soil
73 quality components⁶. In this manner nutrient, expert diagnosis different N management
74 strategies such as nutrients are applied at the right time, at the right place as per requirement
75 of the crop⁷.

76 In this review, we have discussed the optimum quantity of nitrogen that is required by plants;
77 excessive use of nitrogen, problems due to excessive use and the strategies to improve the
78 nitrogen use efficiency, and its impacts on plant's output.

79

80

81 **1. Nitrogen is a key and basic element for crops:**

82 It's the foremost objective of agricultural scientists to increase the food production to meet
83 the requirements, but the environmental protection is also an important factor to save the
84 world regarding climate change⁸. Globally, nitrogen use has been increased effectively. In the
85 past 4 decades, its use has been increased to 100-fold⁹. Now scientists are recognizing the
86 needs of crops that utilize nitrogen efficiently and in a quick way³. For the proper growth of
87 plants, some nutrients are basic and very effective. Nitrogen is one of them that is responsible
88 for the full-fledged growth of crops. In the last years, agricultural scientists around the globe,
89 are taking passionately interest on the optimum use of nitrogen for the lavish growth of
90 agricultural crops^{2,10}.

91 Healthy plants contain 2-4 % nitrogen. Deficiency of nitrogen results in the appearance of
92 chlorosis in plants. By facing the deficit problem, protein quantity is decreased, while sugar
93 content is increased. Protein is made by carbon compounds, and without the availability of
94 nitrogen, these are not built¹¹. Not only for crops but nitrogen is also is an important element
95 in the lives of the living organism, but the most limiting factor only for plants. By applying
96 the adequate supply of nitrogen according to the requirements of plants, enough food can be
97 produced. Optimum use of nitrogen can lead to maximum productivity of plants¹². Deficiency
98 of macronutrients results in the stunted growth as nitrogen deficiency can limit the growth of

99 plants. To get the adequate production from plants, generally fertilizers are applied, but
100 excessive use of nitrogen is no more part of the plants. It is leached down, this can be part of
101 the environment in the form of environmental pollution.

102 Therefore, it is recommended to use the fertilizers according to the needs of plants, in this
103 way productivity and profitability are increased as well maintained. Past research explains
104 well about the positive correlation between nitrogen concentration and chlorophyll content in
105 the leaves of plants. In maize, by measuring the chlorophyll content, nitrogen requirement is
106 estimated¹³. Photosynthetic activity of plants is increased as the nitrogen increases. Majority
107 of leaf nitrogen is represented by the proteins of the Calvin cycle and thylakoids¹⁴. In one of
108 the published papers, nitrogen and chlorophyll content were measured at the flowering stage
109 and found the close relationship between nitrogen content and chlorophyll content. Moreover,
110 chlorophyll structure is composed of nitrogen. As the nitrogen application is increased, the
111 nitrogen that is derived from the soil decreases; only that part of nitrogen is used which is
112 applied through basal dressing and topdressing. Maximum nitrogen is lost by using a basal
113 dressing method than topdressing¹⁴.

114 **2. Excess of anything in life is poison:**

115 Soil fertility is declining continuously. It is considered the main problem of the green
116 revolution era. Intensive cropping is responsible for the removal of fertile nutrients. Use of
117 Inorganic fertilizers is being increased to replenish the soil. Farmers are not well aware of the
118 proper use of nitrogen fertilizers and apply without quantification. More than adequate
119 quantity is applied to agricultural soils, and many other macro and micronutrients are
120 ignored, including Potash, Phosphorus, Zinc e.t.c. Upon application of fertilizers,
121 mineralization is started and it depends upon different factors, including soil microbes,
122 irrigated water, and type of fertilizer¹⁵. One of an excess of nitrogen is lost to running waters
123 and enters in the freshwater lakes, and algal growth appears on the water surface. Due to the
124 algal growth, the creature under the water surface dies¹⁶.

125 Excessive use of nitrogen fertilizer creates many reduced yield problems. Continuous use of
126 huge quantity leads to elevation of NO₃-N concentration in groundwater, causing human
127 health disorders, moreover, day by day, its efficiency is declining. With the groundwater, it is
128 also affecting the surface freshwater resources and becoming the major factor of water
129 pollution¹⁷.

130 It is quite odd that to increase food production, more and more fertilizers are being applied in
131 agricultural land, but nobody cares about environmental pollution. Global nitrogen cycle has

132 been changed effectively. For getting maximum output, the use of nitrogen has been
133 increased¹⁸. Only 30 to 50% NUE is recorded in plants, the remaining is used by soil
134 microbes, leached down in soil or volatilized¹⁹. Plants can use nitrogen in the form of NH₃
135 (ammonium). 82% nitrogen is present in ammonia. When plants are unable to use that
136 ammonia, it is converted in nitrates and enter into the plant roots and causes soil pollution.
137 Like plants, humans are also the victim of nitrates and becomes the part of vegetables that
138 causes severe human health problems²⁰. In the agricultural system, these effluents
139 disseminate in the environmental air in the form of ammonia (NH₃), nitrate (NO₃), and
140 nitrogen oxides (NO₂). These are highly toxic to human and animals health. Therefore, it is a
141 serious concern of present era and a difficult challenge for policymakers. In one of the
142 performed research, it was revealed that if nitrogen content increases, the nitrate
143 concentration in lettuce is also increased²¹. to get sustainable production and clean
144 environment, NUE should be increased. NUE is dependent upon the performance of different
145 steps, comprising of using up, translocation, assimilation, and remobilization. These steps are
146 linked with the environmental and genetic interaction. In this hour of need, by doing well
147 management, NUE can be increased⁶.

148

149 **3. Losses of Nitrogen:**

150 **(a) Nitrogen Loses in Field**

151 Urea is a major source of nitrogen, as allied to crops, some of its quantity is taken up by the
152 crops and utilized for their growth and development, but in the soil-plant production system,
153 most of the quantity of the applied fertilizers are lost by the processes of de-nitrification, soil
154 erosion, surface runoff, leaching, volatilization of ammonia and phosphorus fixation in the
155 soil due to the lower concentration of calcium in the soil²².

156

157 **(b) Soil Erosion and Surface Runoff:**

158 From the soil, N can be lost through the water as well as wind erosion. Loss of N through
159 water erosion is a major problem for humid and sub-humid climatic areas while wind erosion
160 is a more commonly reported mechanism of N loss in the arid and semiarid climatic region⁵.

161

162 **(c) Loss through Leaching and Microbes:**

163 In high rain fed areas and light texture soils (sandy soil), leaching is a common problem.
164 Nitrate form of N does not strongly absorb on soil surface because nitrate is the mobile in

165 nature, and easily move beyond the soil profile by process of leaching. Through this
166 mechanism, as much as 25–50% of the applied N can be lost⁴. This loss can be highly
167 dependent upon the quantity of N applied, climatic conditions and crop production system
168 practices²².

169 In arid to semi-arid areas, leaching problem is documented very less. Soil microorganisms are
170 used the much quantity of applied nitrogen. If microorganism has a ready food supply in the
171 form of organic matter, they readily assimilate nitrates-nitrogen. This is one of the major
172 reasons; microbes can get about more than half the applied nitrogen from the soils.

173

174 **(d) Ammonia Volatilization and Denitrification.**

175 When ammonium and/or urea are applied on the surface of the soil, nitrogen is lost in the
176 gaseous form through the reduction process (volatilization) in which NH_4 convert into NH_3
177 gas. The discussed the phenomenon of N loss is more severe when chemical nitrogenous
178 fertilizers and organic manures is applied on soil surface through broadcasting method⁵.
179 Losses of N in the form of ammonia is a major problem for alkaline soils. Higher
180 concentration of ammonia is not recommended for the nitrification process, as it resulted in
181 un-budgeting of nitrites in the soil. This mechanism is most common in alkaline soil and
182 warm climatic conditions, and more than 20% of N may volatilize by this process and lost to
183 the atmosphere within a short period²³. Under this condition, as much as 10-15% of applied
184 nitrogen has been lost. Denitrification is a more common problem for heavy texture soil with
185 poor natural drainage²⁴.

186

187 **4. Responsible Factors for Low NUE:**

188 To get healthy and high food production, 40% population rely on nitrogen to get healthy and
189 extreme yield. Maize is using 56% of the total nitrogen production. From the total applied
190 fertilizer, only 50% is utilized by plants. While the remaining one is wasted in the form of
191 environmental pollution. The efficiency of applied nitrogen fertilizers depends upon its
192 demand and losses. (crop environmental and management factors affecting nitrogen use
193 efficiency). Agronomic management can increase or decrease of NUE, it depends upon the
194 efficient strategies to use fertilizer according to the need of maize crop. In one of the
195 conducted research, 3 kinds of strategies were applied for managing the fertilizers
196 accordingly, including OPT-1 (Optimized management strategy), OPT-2, and OPT-3. In
197 comparison to OPT-1, OPT-2 showed yield increment. Additionally, farmers fields did not

198 show a significant increase of NUE, but it depends upon the agronomical management
199 strategies (Concurrent Improvement in Maize Yield and Nitrogen Use Efficiency with
200 Integrated Agronomic Management Strategies).

201 Different factors affect the NUE like soil condition, water, and weather. on daily basis, many
202 studies are published. Agronomical management practices, Soil physical and chemical
203 properties, and genotypes cause 18%, 5%, and 12% losses, respectively. Due to less fertility,
204 50% of agricultural lands are not producing the crop with full yield potential.

205 Except this, Low NUE also due to the excessive use of Nitrogen in the fields. Excess use of
206 nitrogen declines the crop yield⁶. Except for the decrease in yield, upon increasing the
207 nitrogen rate, Photosynthetic activity is also reduced²⁵.

208

209 **5. Influences of Different Agricultural Practices:**

210 Already, worldwide, agriculture land is finite for the production of food. To meet the demand
211 for food, production per unit land area should be increased. To utilize the agricultural land
212 efficiently, proper planning and management strategies should be applied²⁶. According to the
213 estimated statistical report, by 2050, the population will be increased to 9 billion, and to feed
214 the whole world, we will be needed to increase food up to 70-100%²⁶. For increasing the
215 production, farmers have to rely on the more use of nitrogen, with the increase of its use,
216 efficient utilization is also required to get the maximum yield. Upon the unreasonable use of
217 the nitrogen fertilizers, its yield is decreased.

218 Appropriate methods, time and application rate always matters, otherwise increased nitrogen
219 rate is no more useful for plants and lost²⁷. Different methods of fertilizers are being applied.

220 Again, the point matters; which method is suitable to increase the nitrogen use efficiency?
221 Before the cultivator use, fertilizers are applied across the whole field; its called as a
222 broadcast method. This method results in non-uniform fertilizer rate across the filed. some
223 places receive more fertilizers. Banding fertilizer method is used to place the fertilizers near
224 the roots, and it is helpful in decreasing the costs and kills weeds maximum. Chih-Li Yu and
225 his team carried out a 3 years study experiment to check the soil respiration, physiological
226 parameters, and yield. Maize behaves Differently in different agricultural practices. Yet, it's
227 the reasoning of differential behave is unclear.

228 Application methods showed different behavior accordingly. Different parameters including,
229 transpiration, photosynthesis, plant height, soil respiration, and yield were measured to asses
230 the differences by adopting six different agricultural practices. Different results showed that

231 application methods do not give significantly different results but the agronomical
232 management practices increase the production of maize²⁸. Likewise the fertilizer application
233 method, application rate also matters a lot for increasing the maize production and nitrogen
234 use efficiency. Luiz Fernando Pricinotto and his team in 2014 published a study on the effects
235 of application time in maize production. Five different nitrogen rates (0, 45, 90, 135, 180)
236 kg/ha were applied. Among these, all application rates, average estimated rate, 130.1 and
237 131.5 kg/ ha proved to produce higher grain yield²⁹. Nowadays, a new kind and effective use
238 are being applied to increase the production and nitrogen use efficiency of plants because
239 CRU is coated with less soluble compounds that make it efficient to use gradually³⁰. Xiang
240 Gao et.al 2007, carried out a study to check the CRU effects on potato and environment.
241 Results clearly depicted that CRU decreases the NH₄⁺ and Nitrates NO₃⁻, thus it does not
242 permit to emit different gasses and increases the NUE³¹. Different studies proved that split
243 nitrogen fertilizer application time is a determinant of higher yield and increase the nitrogen
244 use efficiency. Pre-planting application and side dressing, both are highly effective
245 techniques to increase the yield and nitrogen use efficiency²⁸. The timing of fertilizer can
246 synchronize the demand and uptake of nitrogen fertilizers.

247

248 If management strategies are ignored, the full potential of maize yield and nitrogen recovery
249 efficiency cannot be achieved. Silas et.al 2018, carried out an experiment by using the labeled
250 Nitrogen. Nitrogen was applied at five different stages, including Oat tillering, before 15 days
251 of maize planting time, at the time of corn planting, at three-leaf growth stage V3, and split
252 application at V3 and six-leaf growth. Early nitrogen application is not suited for the
253 availability of nutrients to plants. Soil microbes use the early applied fertilizers and they
254 make it unavailable for plants. Suitable timing of fertilizers increases the nitrogen recovery
255 efficiency and nitrogen content³².

256

257 **6. Need to Increase Nitrogen Use Efficiency:**

258 **(a) The Concept and Importance of NUE**

259 Meeting this requirement in a sustainable manner, is a big challenge today, especially when
260 parallel to historical cereal yield trends which have been linear for nearly half a century.
261 Improving nitrogen use efficiency (NUE) is environmentally and economically desirable
262 traits for crops. NUE is an emerging concept for assessing crop production systems and
263 highly be influenced by fertilizer management. It indicates the potential for nutrient losses to

264 the environment from cropping systems as managers strive to meet the increasing global food
265 and fibre demand.

266

267

268 **(b) Nutrient Use Efficiency**

269 The main objective of nutrient use is to enhance the overall productivity of cropping systems
270 in a sustainable manner while minimizing losses of nutrient from the field. Nitrogen use
271 efficiency (NUE) also addresses the sustainability of agriculture system with respect to soil
272 fertility and some other soil quality components³³. Therefore, the main objective of NUE is to
273 enhance the performance of the overall cropping system³⁴. 78% nitrogen is present in the air
274 but it can not be utilized directly by plants. One acre has 34, 000 tons nitrogen but its direct
275 use is impossible for plants. Nowadays, for the increment of food and make efficient use of
276 nitrogen, highly effective management strategies are needed. Synthetic nitrogen fertilizers are
277 soluble in water and can be readily available to plants. In the 20th century, nitrogen fertilizers
278 were prepared by the Harbor-Bosch process and that was considered as the most important
279 invention³³. Now, nitrogen use is being increased.

280 In the world, China is the leading importer of chemical fertilizers. According to one of the
281 published study, for agricultural outputs, China is consuming 30% of the world's total
282 nitrogen production¹⁸. Although Harbor-Bosch process works for making synthetic fertilizer
283 is the great invention of the 20th century but its minimum and maximum use disturb the
284 plant's output and creates health hazardous for humans. Alone nitrogen is not highly useful
285 for plants to boost the production of crops. In the 19th century, two scientists put forth the law
286 of the minimum, this law clearly states that in the absence of phosphorus or potassium,
287 nitrogen can not give fruitful results. It shows nil behavior. No more yield is increased. Use
288 of nitrogen is directly involved with the plant health and environment. By using different
289 extra activities, nitrogen use efficiency can be increased³⁵. In this scientific arena, by
290 employing different biological approaches, nitrogen use efficiency can be measured quickly
291 and precisely. By improving assimilation and management, nitrogen use efficiency is
292 increased.

293 By employing different breeding schemes and biotechnological tools, new lines with the
294 higher nitrogen use efficiency can be developed. It's a very difficult task to manage the
295 fertilizers according to the requirement of plants. Its tried to manage the nitrogen fertilizer
296 alone or with the combination of other fertilizers. The nitrogen use efficiency is actually the
297 optimum assimilation of nitrogen. Leguminous crops perform better due to their higher

298 nitrogen use efficiency because it is stored in the root system, does not lose in soil or in the
299 air. Nitrogen use efficiency is a very complex trait that is associated with genetic and
300 environmental interaction. Around the world, nitrogen use efficiency is considered lower than
301 standards. Nitrogen use has been increased drastically from 79 million pounds in 2009 to 99
302 million pounds in 2012.

303 However, the use of nitrogen can be improved by designing proper plants and management
304 strategies. According to one of the study, the main problem in the decrease of nitrogen use
305 efficiency is that farmers apply more nitrogen before planting. By doing proper management
306 and previously performed experiments, farmers should use the knowledge and wait for the
307 time of active nitrogen absorption¹. Different kind of agricultural practices is being used to
308 increase nitrogen use efficiency. For managing the nitrogen use, the first step is to do the
309 analysis of plant and soil. Soil analysis components are used to manage the nitrogenous
310 fertilizers, including a quantity of soil organic matter, nitrogen-nitrate credit from the
311 previous crop data, yield targets, and nitrogen credit from irrigation water and manures.
312 Variable nitrogen management zones (MZ) should be identified to apply the fertilizers
313 accordingly, in this way nitrogen use efficiency can be improved.

314 By applying the nitrogen fertilizers according to the demand of specific soil parts, plants
315 perform uniformly and give maximum and uniform yield³⁶. Sometimes, by comparing with
316 C4 plants, nitrogen is recommended to use. For example, by making a comparison to wheat,
317 corn needs less nitrogen for a given biomass³⁷. Another technique to determine the nitrogen
318 requirement is to predict yield target by having knowledge about the previous 5 years
319 performance. Some researcher finds it useful if growing conditions are favorable but
320 sometimes, if the climate is not good, then this suggestion leads to a decrease in nitrogen use
321 efficiency. Because weather conditions are not suitable all time^{38, 39}. Worldwide, agriculturist
322 goes beyond the thinking and solve the problems by utilizing the research skills. Nowadays,
323 there are many sensitive plants are present and these are used for as responsive indicators to
324 fertilizers, weather and soil. For example, chlorophyll in increases, if more nitrogen is
325 applied. So, these plants show the concentration in the form of their phenotypic appearance.
326 And as chlorophyll content in increases, the photosynthetic activity also increases. Previous
327 studies showed that photosynthetic activity has a positive correlation with the nitrogen
328 concentration⁴⁰.

329 Nitrogen concentrations are used as an indicator of maximum crop growth. Critical nitrogen
330 requirement is the optimum amount of nitrogen that can produce maximum yield. Initially, in
331 the plants, nitrogen concentration is higher than the maturity level. As plants grow, nitrogen

332 concentration is decreased⁴¹. The ratio of actually available nitrogen in plants to the critical
333 nitrogen is called as nitrogen nutrition index (NNI). Now, agricultural scientists are using the
334 NNI (nitrogen nutrition index). This approach is being used in wheat, rice, sorghum, and
335 grasses⁴². In maize, this approach can not be used with much efficiency. At early growth
336 stages, critical nitrogen cannot provide a reliable nitrogen status. Usually, nitrogen
337 concentration is decreased as maize shifts toward maturity, and it is called nitrogen dilution⁴³.
338 Up to silage maturity, critical nitrogen dilution curve gives effective results. In corn, this
339 system could be used only at small scale⁴⁴.

340

341 **7. Strategies to Enhance NUE:**

342 **(a) Agronomic Practices**

343 Nutrient use efficiency can be enhanced by adopting local as well as scientifically available
344 means of nutrient management to ensure more efficient use of various agricultural inputs
345 such as fertilizers, irrigation water, and land that will minimize its losses while enhancing
346 beneficial use of these inputs.

347 Strategies used for enhancing the nutrients use efficiency of crops should be focused on two
348 major bases (1) either it enhances the efficacy of externally applied nutrient (2) either it
349 enhances the budget of nitrogen in the soil by reducing N losses through different
350 mechanisms and ensure more uptake of conserved N by crops⁴⁵. Application of the nutrients
351 at a suitable rate, right time, and in the right place is the major and basic principle for
352 attaining the higher nutrients use efficiency⁴⁶. Different practices based on the above-
353 discussed principle for enhancing the nitrogen use efficiency are discussed below: Best
354 nutrient management in wheat-maize cropping systems should aim to apply fertilizers based
355 on the requirement of crops and select a suitable method for maximizing the nutrients use
356 efficiency and reduce its losses⁴⁷. In this manner nutrient, expert diagnosis different N
357 management strategies such as nutrients are applied at the right time, at the right place as per
358 requirement of the crop⁷.

359

360 **(b) Right Rate:**

361 Several crops are highly dependent on location, climate and season so it is essential that
362 accurate yield goals are established and that fertilizers are applied to meet the target yield
363 (Fertilizers Europe, 2011). Excess or low supply of the nutrients will result in reduced NUE
364 and significant losses in yield and grain quality. Soil testing analysis also one of the most

365 powerful and easily conductible tools for determining the capacity of the soil for providing
366 the nutrient to crops. Soil testing approaches also be useful for formulating appropriate
367 fertilizer recommendations, good calibration data in the proper way⁷.

368

369 **(c) Right Time (site-specific nitrogen management):**

370 Great relation between crop requirement and nutrient supply is necessary to enhance the
371 NUE, especially for nitrogen. During the growing season, application of nitrogen in split
372 doses, rather than a single dose at once time are known to be effective in increasing nitrogen
373 use efficiency⁴⁸. For assessing the nitrogen status of growing crops, tissue testing is a cheapest
374 and famous method, but other diagnostic techniques are also commonly available. The use of
375 chlorophyll meters also found as an easy diagnostic tool for enhancing the nitrogen use
376 efficiency in crops⁴⁹. Use of leaf colour charts also recommended for maize crop when
377 nitrogen is applied in split doses⁵⁰.

378

379

380 **(d) Right Place:**

381 Selection of suitable application method has always been crucial in ensuring the nutrients use
382 efficacy. Selection the right placement is an important factor for determining the right
383 application rate. Currently, different placements are available, but surface or subsurface
384 application before or after planting are more common. Prior to planting, nitrogen can be
385 broadcast, or applied as a band on the soil surface, or applied as a subsurface band (15-20 cm
386 deep). Commonly, with banded application method, nutrient recovery efficiency tends to be
387 higher as compared to another method because under band application less nutrient contact
388 with the soil lessens, which reduce the chances for nutrient loss by the leaching process.
389 Selection of the Placement highly dependent on the crop and edaphic factors, which interact
390 to influence the availability and uptake of nutrients. Adequate and balanced application of
391 nutrients is one of the most common practices for enhancing the efficacy of nitrogenous
392 fertilizer both in developed and developing countries³³.

393

394 **(e) Chlorophyll Meter and Leaf Colour Chart:**

395 Chlorophyll meter (CM) can be successfully used to estimate the crop nitrogen content
396 because most of the nitrogen is found in the chloroplast of the plant⁵¹. CM helps in measuring
397 the chlorophyll content and can calibrate it for different climatic, soils and crop cultivars. It is
398 also being recommended to check the effectiveness of late applied nitrogen in standing crops

399 to enhance the protein content and crop productivity. Leaf color chart also used as an
400 indicator of leaf color, color intensity, leaf nitrogen status and helps in selecting the right time
401 of nitrogen application. As a diagnostic tool, it also provides the guideline to the farmers for
402 making appropriate decisions regarding appropriate time, appropriate dose and right method
403 of nitrogen application in standing crops. As concluded, it works on the base of relative
404 greenness of leaves which directly co-related with chlorophyll content of leaves.

405

406

407

408 **(f) Integrated Nutrient Management:**

409 Nitrogen is a basic component of leaf chlorophyll so its measurement over different
410 phenological stages serves as the indirect basis for nitrogen management in different crops⁵¹.
411 Integrated nutrient management involves balanced use of indigenous nutrient components
412 such as crop residues, organic manures, biological nitrogen fixation as well as chemically
413 available nutrients and their complementary interactions to increase the recovery of N
414 recovery⁵¹. Positive effects of the integrated use of organic as well as inorganic nitrogen are
415 either due to optimum Physio-chemical conditions of the soil or due to the better architecture
416 of root and more supply of micronutrients to the plants⁵¹. The exploitation of these positive
417 effects among the plant nutrient is the major detriment for increasing the productivity of
418 cropping system as well as the efficiency of applied nitrogen. The paired interaction of N
419 with other secondary and micronutrients could result in improvement in crops yield and
420 nitrogen use efficiency. Therefore, balanced and judicious use of nitrogenous fertilizers will
421 lead to achieving higher productivity.

422

423 **(g) Increase the Use of Modified Fertilizers and Slow Released Fertilizers:**

424 These are various fertilizer products which are used for enhancing the fertilizer use efficiency
425 of crops by reducing losses of nutrients associated with the production system. These
426 products are based on two basic concepts either they can release in slow or either interfere
427 with nutrient transformation processes and thus reduce their losses. Slow release nitrogenous
428 fertilizers and inhibitors are two important classes of fertilizers. The selection of the suitable
429 type of applied nitrogenous fertilizers has a pivot role in reducing the various nitrogen losses
430 hence, affecting the availability and recovery of nitrogen. As compared with ammonium and
431 amide containing nitrogen fertilizers, nitrate fertilizers are more susceptible to leaching. But

432 in contrast, ammonium and amide containing fertilizers are more susceptible to volatilization
433 process than nitrate fertilizers.

434 A variety of slow-release fertilizers is now easily available in the market which has the
435 potential to increase the nitrogen use efficiency and reduces the nitrogen losses⁴⁸. Polymer-
436 coated products are commonly used in agriculture, which can be designed to supply the
437 nutrients to crops in a controlled manner. Nutrient release rates are highly dependent on
438 properties of the polymer coating, soil temperature, and moisture conditions. In developing
439 countries, non-availability and high manufacturing cost are two major reasons for the limited
440 use of these compounds. In additions, some others approach to enhance the nitrogen use
441 efficiency is the use of N stabilizers which increase the nitrogen use efficiency not only by
442 minimizing leaching losses but also by reducing the de-nitrification losses⁵².

443

444

445

446

447

448 **Conclusion:**

449 To increase the crop production fertilizers play a vital role. Among fertilizers nitrogen is
450 more important as it helps the plants in the preparation of protein. Its deficiency effects the
451 growth of the plant and its excess reduces the crop yield. Plant uses an optimum level of
452 nitrogen and the remaining is leached down into the soil. The excess of nitrogen in the field
453 cause environmental problems and health hazards. Plants have low NUE. It is the need of the
454 time to increase the nitrogen efficiency of the plants. Different experiments are going on to
455 increase the NUE of plants. Agronomic practices can also help in this regard. Nitrogen given
456 at the right time and right place can increase the plant efficiency to use to effectively. Now a
457 days slow release fertilizers are also in use to control the loss of nitrogen by the plants.
458 Moreover, for the better development of plant more practices and improvement in plant is
459 needed to use nitrogen more effectively.

460

461

462

463

464

465 **References:**

466

- 467 1. Sharma, L., Sustainability, S. B.- & 2017, undefined. A review of methods to improve
468 nitrogen use efficiency in agriculture. *mdpi.com*
- 469 2. Journal, E. Analysis of improvements in nitrogen use efficiency associated with 75
470 years of spring barley breeding. **42**, (2012).
- 471 3. Ag, G., Ak, S. & Dg, M. Can less yield more ? Is reducing nutrient input into the
472 environment compatible with maintaining crop production ? **9**, (2004).
- 473 4. Helmers, M. J., Christianson, R. D. & Sawyer, J. Nitrate loss in subsurface drainage as
474 affected by nitrogen application rate and timing under a corn- soybean rotation system.
475 (2016). doi:10.31274/icm-180809-76
- 476 5. Fageria, N. K. Soil quality vs. environmentally-based agricultural management
477 practices. *Commun. Soil Sci. Plant Anal.* **33**, 2301–2329 (2002).
- 478 6. Xu, G., Fan, X. & Miller, A. J. Plant Nitrogen Assimilation and Use Efficiency.
479 doi:10.1146/annurev-arplant-042811-105532
- 480 7. Pampolino, M. F., Witt, C., Mae, J., Johnston, A. & Fisher, M. J. Development
481 approach and evaluation of the Nutrient Expert software for nutrient management in
482 cereal crops. *Comput. Electron. Agric.* **88**, 103–110 (2012).
- 483 8. Muriuki, A. W. Effect of organic and conventional farming systems on nitrogen use
484 efficiency of potato , maize and vegetables in the Central highlands of Kenya. (2017).
- 485 9. Glibert, P. M., Harrison, J., Heil, C. & Seitzinger, S. Escalating worldwide use of urea
486 – a global change contributing to coastal eutrophication. 441–463 (2006).
487 doi:10.1007/s10533-005-3070-5
- 488 10. Hirel, B. *et al.* Towards a Better Understanding of the Genetic and Physiological Basis
489 for Nitrogen Use Efficiency in Maize. (2019).
- 490 11. Weil, R. R. Nitrogen and Sulfur Economy of Soils. (2017).
491 doi:10.13140/RG.2.1.1435.0482
- 492 12. Indian, T. & Assessment, N. 12 - Nitrogen Nutrition in Crops and Its Importance in
493 Crop Quality. (2017).
- 494 13. Schepers, J. S., Francis, D. D., Vigil, M. & Below, F. E. Communications in Soil
495 Science and Plant Analysis Comparison of corn leaf nitrogen concentration and
496 chlorophyll meter readings. **3624**, (2017).
- 497 14. Journal, E. Relationship between nitrogen and chlorophyll content and spectral
498 properties in maize leaves. **2**, (1993).

- 499 15. The cost of uncertainty for nitrogen fertilizer management : A sensitivity analysis. **100**,
500 (2007).
- 501 16. Liu, C., Sung, Y., Chen, B. & Lai, H. Effects of Nitrogen Fertilizers on the Growth and
502 Nitrate Content of Lettuce (*Lactuca sativa* L .). 4427–4440 (2014).
503 doi:10.3390/ijerph110404427
- 504 17. Wang, D. & Xu, Z. *Acta Agriculturae Scandinavica* , Section B - Soil & Plant Science
505 Excessive nitrogen application decreases grain yield and increases nitrogen loss in a
506 wheat – soil system. (2015). doi:10.1080/09064710.2010.534108
- 507 18. Ali, S. B. *et al.* IMPACT OF EXCESSIVE NITROGEN FERTILIZERS ON THE
508 ENVIRONMENT AND. 1–23
- 509 19. Ng, J. M., Han, M., Beatty, P. H. & Good, A. “ Genes , Meet Gases ”: The Role of
510 Plant Nutrition and Genomics in Addressing “ Genes , Meet Gases ”: The Role of
511 Plant Nutrition and Genomics in Addressing. (2016). doi:10.1007/978-1-4939-3536-9
- 512 20. Long-term effects of semisolid beef manure application to forage grass on soil
513 mineralizable nitrogen . 2011 (2011).
- 514 21. Lemaire, G. *improve nitrogen-use efficiency*. (2015). doi:10.1016/B978-0-12-417104-
515 6.00008-X
- 516 22. Wheat, W. Nitrogen Balance in the Magruder Plots Following 109 Years in
517 Continuous Nitrogen Balance in the Magruder Plots Following 109 Years in
518 Continuous. (2003). doi:10.1081/PLN-120022364
- 519 23. Fageria, N. K. *Communications in Soil Science and Plant Analysis* Soil quality vs .
520 environmentally-based agricultural management practices. **3624**, (2007).
- 521 24. Hutchinson, C. Testing of Controlled Release Fertilizer Programs for Seep Irrigated
522 Irish Potato Production. **26**, (2007).
- 523 25. Li, Y. *et al.* Does Chloroplast Size Influence Photosynthetic Nitrogen Use Efficiency ?
524 Does Chloroplast Size Influence Photosynthetic Nitrogen Use Efficiency ? (2013).
525 doi:10.1371/journal.pone.0062036
- 526 26. Optimising yield and resource utilisation of summer maize under the conditions of
527 increasing density and reducing nitrogen fertilization. 2017 (2017).
- 528 27. Research advances on regulating soil nitrogen loss by the type of nitrogen fertilizer
529 and its application strategy . 29732872 doi:10.13287/j.1001-9332.201609.022
- 530 28. Yu, C. *et al.* Responses of corn physiology and yield to six agricultural practices over
531 three years in middle Tennessee. *Nat. Publ. Gr.* 1–9 (2016). doi:10.1038/srep27504
- 532 29. Pricinotto, L. F., Soares, P., Filho, V., Scapim, C. A. & José, O. Effects of nitrogen

- 533 rates and application time on popcorn. (2014). doi:10.5897/AJAR2013.7471
- 534 30. Tong, X., He, X., Duan, H., Han, L. & Huang, G. Evaluation of Controlled Release
535 Urea on the Dynamics of Nitrate , Ammonium , and Its Nitrogen Release in Black
536 Soils of Northeast China. (2018). doi:10.3390/ijerph15010119
- 537 31. Access, G. Controlled release urea improved the nitrogen use efficiency , yield and
538 quality of potato (*Solanum tuberosu m L .*) on silt loamy soil. 2–4 (2019).
- 539 32. Oliveira, S. M. De *et al.* Understanding N timing in corn yield and fertilizer N
540 recovery : An insight from an isotopic labeled-N determination. 1–14 (2018).
- 541 33. Hirel, B., Tétu, T., Lea, P. J. & Dubois, F. Improving Nitrogen Use Efficiency in
542 Crops for Sustainable Agriculture. (2011). doi:10.3390/su3091452
- 543 34. Barbieri, P. & Andrade, F. H. Row Spacing Effects at Different Levels of Nitrogen
544 Availability in Maize. (2000). doi:10.1007/s100870050034
- 545 35. Carvalho, E. V. De, Afféri, F. S. & Peluzio, J. M. Nitrogen use efficiency in corn (*Zea mays L*)
546 genotypes under different conditions of nitrogen and seeding date
547 Original Paper Open Access Nitrogen use efficiency in corn (*Zea mays L*) genotypes
548 under different conditions of nitrogen and seeding date. (2012).
- 549 36. Kitchen, N. R., Sudduth, K. A., Myers, D. B., Drummond, S. T. & Hong, S. Y.
550 Delineating productivity zones on claypan soil fields using apparent soil electrical
551 conductivity. **46**, 285–308 (2005).
- 552 37. Gastal, F. & Lemaire, G. N uptake and distribution in crops : an agronomical and
553 ecophysiological perspective. **53**, 789–799 (2002).
- 554 38. Puntel, L. A. *et al.* Modeling Long-Term Corn Yield Response to Nitrogen Rate and
555 Crop Rotation. **7**, 1–18 (2016).
- 556 39. Yield, D., Fertilizer, M. & Requirement, N. Delta Yield: Mapping Fertilizer Nitrogen
557 Requirement for Crops. **80**, 20–23 (1996).
- 558 40. Science, P. Leaf Nitrogen Content , Photosynthesis and Radiation Use Efficiency in
559 Peanut . 40–43 (1993).
- 560 41. Greenwood, A. D. J., Lemaire, G., Gosse, G., Cruz, P. & Draycott, A. Decline in
561 Percentage N of C3 and C4 Crops with Increasing Plant Mass Published by : Oxford
562 University Press Stable URL : <https://www.jstor.org/stable/42758332> Decline in
563 Percentage N of C3 and C4 Crops with Increasing Plant Mass. **66**, 425–436 (2019).
- 564 42. Justes, E., Mary, B. & Jean-marc, M. Determination of a Critical Nitrogen Dilution
565 Curve for Winter Wheat Crops. (1994). doi:10.1006/anbo.1994.1133
- 566 43. Pl, D. Relationships between dynamics of nitrogen uptake and dry matter

567 accumulation in maize crops . Determination of critical N concentration. 1983–1984
568 (2000).

569 44. Division, I. S. & Library, N. A. range of the critical nitrogen dilution curve for maize
570 (*Zea mays* L.) can be extended until silage maturity [2004]. 2–4 (2019).

571 45. Humphreys, E., Timsina, J., Meisner, C. A. & Masih, I. Permanent beds and rice-
572 residue management for rice-wheat systems in the Indo-Gangetic Plain: overview.

573 46. Majumdar, K., Jat, M. L., Pampolino, M. & Satyanarayana, T. Nutrient management in
574 wheat : current scenario , improved strategies and future research needs in India
575 Nutrient management in wheat : current scenario , improved strategies and future
576 research needs in India. (2013).

577 47. Basso, B., Ritchie, J. T., Cammarano, D. & Sartori, L. A strategic and tactical
578 management approach to select optimal N fertilizer rates for wheat in a spatially
579 variable field. *Eur. J. Agron.* **35**, 215–222 (2011).

580 48. Cassman, K. G. & Walters, D. T. Agroecosystems , Nitrogen-use Efficiency , and
581 Nitrogen Management Agroecosystems , Nitrogen-use Efficiency ,. (2002).

582 49. Francis, D. D. & Piekielek, W. P. Assessing Crop Nitrogen Needs with Chlorophyll
583 Meters.

584 50. Witt, B. C. & Dobermann, A. A Site-Specific Nutrient Management Approach for
585 Irrigated , Lowland Rice in Asia. **16**, 20–24 (2002).

586 51. Singh, U. Integrated Nitrogen Fertilization for Intensive and Sustainable Agriculture
587 Integrated Nitrogen Fertilization for Intensive and Sustainable Agriculture. **7528**,
588 (2008).

589 52. Shivay, Y. S. Coating of prilled urea with ecofriendly neem (*Azadirachta indica* A .
590 Juss .) formulations for efficient nitrogen use in hybrid rice COATING OF PRILLED
591 UREA WITH ECOFRIENDLY NEEM (*Azadirachta indica* A . Juss .)
592 FORMULATIONS FOR EFFICIENT NITROGEN USE IN HYBRID RICE. (2014).
593 doi:10.1556/AAgr.51.2003.1.7
594
595
596
597
598
599