

## Original Research Article

### **Preliminary Phytochemical Screening and Thin Layer Chromatography Analysis of Stem Bark Extracts of African Mistletoe Parasitic on *Vitellaria paradoxa*, *Pilostigma thonningii* and *Combretum fragrans***

#### **ABSTRACT**

**Aim:** Mistletoes are highly utilized in traditional medicine to treat different kinds of diseases such as heart diseases, diabetes and malaria, among others. The chemistry of African mistletoe is not sufficiently documented. This paper is therefore, aimed at determining the phytochemicals present in the crude extracts of mistletoe parasitic on plants that are commonly seen as hosts.

**Study Design:** This study was designed to compare the phytochemical profiles of mistletoe stem barks obtained from different plant hosts.

**Place and Duration of Study:** Department of Chemistry, University of Agriculture, Makurdi, Benue State Nigeria, between August and September, 2018.

**Methodology:** Powdered stem bark of mistletoe was extracted successively with hexane, ethyl acetate and methanol. Preliminary phytochemical screening was carried out on the extracts. Thin layer chromatography (TLC) was carried out on silica gel precoated plates in 9:1 (hexane/ethyl acetate), 5:5 (hexane/ethyl acetate), and 7:3 (ethyl acetate/methanol) mobile phases for hexane, ethyl acetate and methanol extracts respectively.

**Results:** The study revealed the presence of secondary metabolites such as alkaloids, flavonoids, tannins/phenols, cardiac glycosides, steroids and triterpenoids. It was evident from TLC analysis that mistletoes from various plant hosts contain similar chemical profile.

**Conclusion:** We therefore debunk the claim by some herbalists that medicinal values of mistletoes vary due to host plant. This is the first time a study of this kind is reported on mistletoe parasitic on *Vitellaria paradoxa*, *Pilostigma thonningii*, *Combretum fragrans*.

**Key words:** Mistletoe, Thin layer chromatography, Phytochemicals, Silica gel, Mobile phase

#### **1. INTRODUCTION**

African mistletoes are hemiparasitic plants that grow on other plants. The fact that they photosynthesize and derive some salts from host plants; they are regarded as hemiparasitic [1]. Mistletoes grow on many plants in Nigeria including *Vitellaria paradoxa*, *Pilostigma thonningii*, *Combretum fragrans*, *Parkia biglobosa* and many others. They are also rarely seen to grow on mango, guava, cocoa and kola nut trees etc [2]. It is believed that birds excrete the seeds of mistletoe through faeces on trees upon which they sit. The sticky faeces facilitate the attachment of seeds on tree branches [3]. Almost all trees could

have opportunity to host mistletoe but just a few have been seen to do so. In Igboughul District of Bali Local Government Area, Taraba State, almost 9 in 10 *Vitellaria paradoxa* (shea butter tree) and 8 in 10 *Pilostigma thonningii* host mistletoe. This could be due to their thick, fresh and easily penetrated stem barks.

Medicinal plants produce various classes of secondary metabolites such as alkaloids, tannins, steroids, phenols, saponins, flavonoids glycosides, terpenoids and others that are responsible for therapeutic and defense properties [4]. Preliminary phytochemical screening is a useful method of detecting these bioactive principles that are present in medicinal plants and may subsequently in drug discovery and development processes [5].

Mistletoes are highly utilized in traditional medicine to treat different kinds of diseases such as heart diseases, diabetes, malaria, cancer and diarrhea [6],[3] etc. However, some traditional practitioners claim that the use of mistletoe depends on the type of host plant- mistletoes from specific plants are used to treat specific diseases [6]. In Nigeria for example, mistletoes found on bamboo trees and gamba grasses are used to perform rituals especially money making. However, mistletoe on bamboo and gamba grasses is not common.

The chemistry of African mistletoe is not sufficiently documented. However, polysaccharides, lecithin, viscotoxin and peptides, were isolated and identified structurally [1],[7] The report published by Ezema *et al.* [8] revealed the presence of cardiac glycosides, steroids, saponins, carbohydrates, and terpenoids in the leaves of mistletoe parasitic on *Parkia biglobosa*. The methanolic extract of mistletoe leaves were shown to contain saponins, alkaloids, phenols, flavonoids and tannins [2].

From the survey of literature, little or no information is reported of the phytochemistry of mistletoe parasitic on *Vitellaria paradoxa*, *Pilostigma thonningii* and *Combretum fragrans*. This paper is therefore, aimed at determining the phytochemicals present in the crude extracts of mistletoe parasitic on these plants that are commonly seen as hosts in Igboughul District of Bali L.G.A of Taraba State using preliminary phytochemical screening and thin layer chromatography analyses of crude extracts.

## 2. MATERIALS AND METHODS

### 2.1 Sample Collection

Mistletoe stem were harvested from *Vitellaria paradoxa*, *Pilostigma thonningii* and *Combretum fragrans* in Igboughul District of Bali L.G.A, Taraba State in August, 2018. The barks were peeled and allowed to dry under shed. The barks were pulverized using pestle and mortar.

### 2.2 Extraction

Using cold macerations, a powdered sample (10 g) was extracted with 50 mL of hexane for 48 hour with intermittent shaking. The macerated sample was filtered using Whatman filter paper No. 5 and the filtrate allowed to evaporate to obtain crude hexane extract. The residue was allowed to dry for further extraction with ethyl acetate followed by methanol [9].

### 2.3 Preliminary Phytochemical Screening

Preliminary phytochemical screening of the hexane, ethyl acetate and methanol crude extracts each of the three plants were carried out based on routine practices described by Adawia *et al.* [10]; Sabri *et al.* [11]; Satheesh *et al.* [12].

### **2.3.1 Test for steroids and triterpenoids (Liebermann-Burchard test)**

Approximately 3 mg of an extract was mixed with 3 drops of acetic anhydride, boiled and cooled. Concentrated sulphuric acid was then added from the sides of the test tube and observed for the formation of a brown ring at the junction of two layers. Green coloration of the upper layer and the formation of deep red colour in the lower layer would indicate a positive test for steroids and triterpenoids, respectively.

### **2.3.2 Test for cardiac glycosides (Keller-Killiani Test)**

About 3 mg an extract was mixed with 3 drops of conc. glacial acetic acid and diluted ferric chloride solution and mixed. Concentrated sulphuric acid was added, and observed for the formation of two layers. Lower reddish brown layer and upper acetic acid layer which turn bluish green would indicate a positive test for glycosides.

### **2.3.3 Test for phenolics and tannins (Ferric chloride test)**

About 2 mg each of a crude extract was dissolved in 2 mL of solvent of extraction and treated with 4 drops of ferric chloride solution. Formation of bluish black colour would indicate the presence of phenols. Generally, the formation of bluish-black colour would indicate the presence of gallic tannins and bluish-green would indicate the presence of catechic tannins.

### **2.3.5 Test for flavonoids (alkaline test)**

About 5 mg of an extract was added 5 mL of diluted sodium hydroxide solution. The appearance of yellow colour which would become colourless on addition of few drops of dilute hydrochloric acid would indicate the presence of flavonoids.

### **2.3.6 Test for saponins**

The ability of saponins to produce frothing in aqueous solution and to haemolyse red blood cells was used as screening test for these compounds. Distilled water (5 mL) was added to an extract (5 mg) and strongly shaken in a test tube. Formation of a large amount of froths that would last for about 30 minutes indicated the presence of saponins.

### **2.3.7 Test for alkaloids**

About 3 mL of an extract was mixed with 1 mL of 10% HCl in a test tube and heated (**over boiling water**) for 20 minutes. This was allowed to cool and filtered; 1 mL of the filtrate was treated with few drops of Mayer's reagent. Appearance of creamy precipitate would indicate the presence of alkaloids.

## **2.4 Thin Layer Chromatography**

Approximately 2 mg of an extract was reconstituted with solvent of extraction and spotted on silica gel precoated plates. The extracts were drawn with capillary tubes and applied as spots on a stationary phase (silica-gel coated plate) about 1 cm from the base. These plates were developed in suitable solvent system of 9:1 (Hex/E.A), 5:5 (Hex/E.A), and 7:3 (E.A/MeOH) for hexane, ethyl acetate and methanol extracts respectively in a TLC tank. The developed plates were then heated at about 120 °C for charring to have a better vision of possible spot [13].



Steroids/triterpenoids	+	+	+	+	+	+	-	-	-
Cardiac glycosides	-	-	-	+	+	+	+	+	+
Phenols/Tannins	-	-	-	+	+	+	+	+	+
Flavonoids	-	-	-	+	+	+	+	+	+
Saponins	-	-	-	-	-	-	-	-	-
Alkaloids	-	-	-	-	-	-	+	+	+

Key: - = Absence, + = presence

*V.P= Vitellaria paradoxa, C.F= Combretum fragrans, P.T= Pilocostigma thonningii*

The phytochemical screening result (Table 2) showed that only steroids and triterpenoids were present in the hexane extracts of all the samples. It was shown from the result that ethyl acetate extracts contained steroids and triterpenoids, cardiac glycosides, phenols/tannins and flavonoids. In addition to alkaloids, the methanol extracts contained similar classes of compounds to ethyl acetate. Saponins were however, not detected in any of the samples. Petroleum ether (non-polar solvent) extract of leaves of mistletoe were found to contain tannin, saponins, steroids and terpenes; the ethanol (polar solvent) extract contained tannins, alkaloids, glycosides, steroids, saponins and terpenes [15]. This result is comparable to that published using leaves of mistletoe by Tabe *et al.* [2].

#### 4. CONCLUSION

The stem bark extracts of mistletoe contained alkaloids, flavonoids, steroids, triterpenoids, cardiac glycosides and tannins. Based on similar phytochemical profile possessed by mistletoes obtained from different plants, it can be concluded that the plant may have similar medicinal or biological effects irrespective of their hosts. Furthermore it can be inferred from this research that, mistletoe do not obtain their phytochemicals from host plants but rather produce by them since they undergo photosynthesis on their own. Thus, the claim by some herbalists that medicinal values of mistletoes vary due to host plant is debunked. This is the first time a study of this kind is reported on mistletoe parasitic on *Vitellaria paradoxa*, *Pilocostigma thonningii*, *Combretum fragrans*.

#### REFERENCES

1. Adesina SK, Illoh HC, Imoh IJ, Imo EJ. African Mistletoes (Loranthaceae); Ethnopharmacology, Chemistry and Medicinal Values: An update. African Journal of Traditional, Complementary and Alternative Medicine. 2013; 10(3): 161-170
2. Tabe NN, Ushie OA, Jones BB, Kendenson AC, Muktar M, Ojeka CU. Phytochemical Analysis of Methanolic Extract of Mistletoe Leaf. International Journal of Advanced Research in Chemical Science. 2018; 5(7): 7-11
3. Saleh I, Maigandi SA, Hudu MI, Abubakar MI, Shehu AU. Uses and Chemical Composition of Mistletoe Obtained From Different Species of Trees. Dutse Journal of Agriculture and Food Security. 2015; 2(1): 8-12

4. Byadgi SA, Kulloli SD, Venugopa CK. Phytochemical Screening and Antimicrobial Activity of Plant Extracts for Textile Applications. *International Journal of Biochememistry Research and Review*. 2017; 20(3): 1-10.
5. Karande AM, Kamble HV, Kumbhar VH, Kane SR, Magdum CS. Preliminary Phytochemical Screening of *Glochidion ellipticum*. *European Journal of Experimental Biology*. 2016; 6(4): 41-45
6. Osadebe PO, Okide GB, Akabogu IC. Study of Antidiabetic activity of Crude Methanolic Extracts of *Loranthus micranthus* (linn). *J. Ethnopharmacol*. 2004; 95: 133-138
7. Adebisi F, Oluwaseye A, Fausat A, Michael A, Adedayo A, Ikokoh P. Chemical Characterization and Antimicrobial Activity of *Loranthus micranthus* Linn Leaves. *Journal of Medicinal Plant Research*. 2013; 7(45): 3332-3336
8. Ezema BE, Eze FU, Ezeofor CC. Phytochemical and Antibacterial Studies of Eastern Nigerian Mistletoe (*Loranthus Micranthus*) Parasitic on *Pentacletra macrophylla* and *Parkia biglobosa*. *International Journal of Pharmaceutical Technological Research*. 2016; 9(5): 360-365
9. Azwanida NN. A Review on Extraction Methods Use in Medicinal Plants, Principle Strength and Limitation. *Medicinal and Aromatic Plants*. 2015; 4(3):1-6
10. Adawia K, Rawaa A, Ghalia S. Phytochemical Screening and Antioxidant Activity of Selected Wild Plants in Liliaceae Family Growing Syria. *International Journal of Pharmacognosy and Phytochemical Research*. 2016; 8(12): 2025-2032
11. Sabri FZ, Belarbi M, Sabri S, Alsayadi M.M. Phytochemical Screening and Identification of some Compounds from Mallow. *Journal of Natural Products and Plant Research*. 2012; 2(4):512-516
12. Satheesh KB, Suchetha KN, Vadisha BS, Sharmila KP, Mahesh PB. Preliminary Phytochemical Screening of Various Extracts of *Punica granatum* Peel, Whole Fruit and Seeds. *Nitte University Journal of Health Science*. 2012; 2(4): 34-38
13. Wahab OM, Ayodele AE, Moody JO. TLC phytochemical screening in some Nigerian Loranthacea. *Journal of Pharmacognosy and Phytotherapy*. 2010; 2(5): 64-70
14. James JT, Dubery IA. Identification and Quantification of Triterpenoid Centelloids in *Centella asiatica* (L.) Urban by Densitometric TLC. *Journal of Planar Chromatography- Modern TLC*. 2011; 24(1): 82-87
15. Nwankwo NE, Egbunu ACC. Phytochemical and Antimicrobial Activity of Petroleum Ether Extract of the African Mistletoe (*Loranthus micranthus* Linn) Leaves. *International Research of Pharmacy and Pharmacology*. 2011; 1(8): 211-214