

Socio-demographic and obstetric variations of *T. gondii* and HIV-1 co-infection among pregnant women in Cameroon

ABSTRACT

Aims: This study, aimed to identify the seroprevalence of *T. gondii* and HIV-1 co-infection in pregnant women in the Northwest Region of Cameroon.

Study design: This cross-sectional study was conducted among 606 pregnant women attending antenatal clinic.

Place and Duration of Study: This study was carried out at the Bamenda Regional Hospital from May 2017 to December 2017.

Methodology: Venous blood samples were collected for the detection of anti-*Toxoplasma* antibodies using rapid test kits. Data were analyzed using SPSS version 23 statistical package. *P*-value <0.05 was considered statistically significant.

Results: The mean (SD) age was 27.3 (5.3) years. The prevalence of *T. gondii* and HIV-1 was 139 (22.9%) and 70(11.6%) respectively, while that of *T. gondii* and HIV-1 co-infection was 31(5.1%). **With the exception of age group and gestational age that was significant for HIV**, socio-demographic and obstetrical characteristics of *T. gondii*, HIV-1 and *T. gondii* and HIV-1 co-infection prevalence did not show any significant differences (*p*>0.05). **Conclusion:** The high prevalence of *T. gondii* and HIV-1 co-infection seen in this study demonstrates the need for routine antenatal screening for both infections. In addition, data from this study will be useful in designing control and prevention strategies against diseases. Furthermore, the result will also be used as baseline data for further research on *T. gondii* and HIV-1 co-infection.

Keywords: Co-infection Human immunodeficiency virus, Pregnant women, Toxoplasmosis, Cameroon

1. INTRODUCTION

Antenatal care provides adequate measures against maternal-fetal transmission of several diseases, including toxoplasmosis and HIV [1, 2]. Toxoplasmosis caused by *Toxoplasma gondii* is a neglected zoonotic disease and is asymptomatic [3, 4]. Toxoplasmosis is prevalent worldwide whereby about one third to half of the global population is infected [5-7]. Human infections result from **foodborne** transmission (consumption of water, raw or undercooked meat or unpasteurized milk contaminated with cyst), animal to human transmission (ingestion of oocysts through close contact with infected cats or cat's **faeces**), mother-to-child transmission (from an infected woman to her unborn child) and through blood transfusion and organ transplants [6-8].

Infection with *T. gondii* has severe consequences in immune compromised hosts such as pregnant women, HIV patients and patients receiving chemotherapy or immunosuppressive drugs [4, 5, 9]. The prevalence of toxoplasmosis among pregnant women showed significant variation between continents and countries and ranges from 9 - 92.5% [6, 8, 9]. In Cameroon, the prevalence ranges from 48.5 - 70% [5, 10]. This variation depends on social and cultural habits, geographic factors, individual's hygiene, route of transmission and the immune status [5, 10, 11]. The high prevalence in pregnant women indicates a greater probability of congenital transmission with latent infection reactivated when immunity

32 is suppressed [5, 12]. Reactivation of latent *T. gondii* infection causes severe and fatal neonatal
33 complications such as stillbirth or abortion, anemia, petechiae due to thrombocytopenia, seizures,
34 neurological defect (epilepsy), ocular disease (blindness, chorioretinitis, strabismus, retinochoroiditis)
35 microcephaly, brain damage (intracranial calcifications, hydrocephalus), mental retardation, cardiac and
36 cerebral anomalies [8, 12, 13]. Congenital transmission of the infection during the first trimester is critical
37 and causes severe clinical conditions in the fetus, whereas infections during the third trimester lead to
38 rapid transmission [4, 14].

39
40 Worldwide, about 36.7 million people are infected with HIV, but very little is known about the prevalence
41 of HIV-1 co-infection with *T. gondii* parasites [15, 16]. Early HIV diagnosis and interventions among
42 pregnant women have shown to decrease the likelihood of mother to child transmission [10, 17]. HIV
43 prevalence is shown to increase among pregnant women as such screening all women during antenatal
44 care is important [17, 18]. The prevalence of HIV among pregnant women ranges from 0.5-61.6% in other
45 countries [11, 17, 19] and between 2.6 -22.1% within other towns in Cameroon [16, 20; 21]. With the
46 advent of highly active antiretroviral therapy, the rate of mother to child transmission has greatly reduced
47 to about 1.4-2.5% [18].

48
49 *T. gondii* and HIV co-infections cause serious complications in pregnant women and pose a serious
50 health threat [11]. Although screening practices of *T. gondii* and HIV during antenatal care are
51 standardized in developed countries, it is somehow limited in developing countries where the burden of *T.*
52 *gondii* infection among HIV infected pregnant women is greatly felt [20]. As such screening for *T. gondii*
53 and HIV infections among pregnant women may be an important primary prevention strategy. Studies
54 carried out elsewhere have shown co-infection rates between 12-40.8% [9, 12]. However, such data are
55 dearth in many developing countries including Cameroon.

56
57 This study is the first study to determine the prevalence of *T. gondii* and HIV co-infection among pregnant
58 women in the Northwest region of Cameroon. It is hoped that the outcome of this study will enable
59 policymakers to design effective strategies for controlling and preventing the disease which in turn will
60 curb the maternal-fetal transmission rate alongside its associated complications. In addition, it will set a
61 base for further studies to be carried out in this area.

62 63 2. MATERIAL AND METHODS

64 65 Study site and design

66 This study was a hospital-based cross-sectional study conducted at the Bamenda Regional Hospital from
67 May 2017 to December 2017. This hospital serves as a referral hospital for the entire Northwest region
68 (NWR). The NWR is characterized by wet and hot climates which have been documented to favor *T.*
69 *gondii* oocyst survival. Inhabitants in this region keep domestic animals like cats, sheep, dogs, goats,
70 fowls that have shown to transmit the disease [4, 12]. Roasted meat (beef, pork, fish, and chicken) is a
71 common delicacy eaten by most people on a daily basis and is a medium for ingesting infectious
72 parasites.

73 Ethical considerations

74 Ethical clearance and administrative authorization were obtained from the ethical review board of the
75 delegation of Public Health Bamenda and Bamenda Regional Hospital review board. Each subject gave
76 their consents before sample collection. Participation in the study was on a voluntary basis and study
77 participants were free to withdraw from the study before and after collection of blood samples without
78 losing any of the benefits they were supposed to obtain from the hospital.

79 80 Sample size determination and sampling technique

81 The sample size was calculated based on Toxoplasma morbidity using the Lorenz formula

$$82 \quad N = \frac{(Z_{1-\alpha})^2 P(1-P)}{i^2}$$

83 Where, $Z_{1-\alpha}$ = the normal distribution value = 1.96

84 P =Relative prevalence of HIV in the region= 54.5% [5]
 85 i = precision (sampling error) = 0.05
 86 The minimum sample size (N) was calculated to be 382.

87 **Data collection**

88 A structured closed-ended questionnaire was used to obtained information on socio-demographic and
 89 obstetric data.

90 **Sample collection and processing**

91 A total of 2ml venous blood was collected using labeled test Ethylene diamine tetraacetic (EDTA) tubes
 92 by the hospital laboratory technician and centrifuged to obtain plasma. Diagnosis of toxoplasmosis was
 93 done using the OnSite ToxolgG/IgM rapid test (CTK Biotech Inc, USA) as per the manufacturer's
 94 procedure. This rapid test kit simultaneously detects both IgG and IgM anti-*Toxoplasma gondii* antibodies.
 95 Whereas HIV test was done using the Alere Determine™HIV-1/2 test kit(Alere, Japan) and confirmatory
 96 test for those that were positive was done using OraQuick HIV 1/2 Rapid Antibody Test (for OraSure
 97 Technologies, Thailand) as described in the manufacturer's procedure.

98 **Data analysis**

99 The data were analyzed using the SPSS statistical software package version 23. Discrete variables were
 100 tested using the chi-square test. Chi-square was used for comparison between categorical variables
 101 through cross-tabulations. P-values of < 0.05 were considered statistically significant.
 102
 103

104 **3. RESULTS AND DISCUSSION**

106 **3.1 RESULTS**

107 A total of 683 pregnant women were approached and, 650 provided consent for the study. Of these 606
 108 women who had recorded HIV status were considered for the study. The age range was 14-45 years with
 109 a mean (SD) of 27.3 (5.3) years. Of the 606 participants, the age group 21-30years 397 (65.5%) were the
 110 most represented, 362 (59.7%) participants were married and 301(49.7%) participants had attained
 111 secondary education. A greater number of them were multigravidae 381 (62.9%) and were in their third
 112 trimester of pregnancy 350(57.8%) (Table 1).
 113
 114

115 **Table 1: General characteristic of the study participants**

Demographic characteristics	Number (%)
Age group in Years	
<21	53(8.7)
21-30	397(65.5)
>30	156(25.7)
Marital status	
Single	164(27.1)
Concubine	58(9.6)
Married	362(59.7)
Widow	22(3.6)
Level Educational	
None	17(2.8)
Primary	107(17.7)
Secondary	301(49.7)
Tertiary	181(29.9)
Gestational age classification	
First (<14weeks)	106(17.5)
Second (14-28weeks)	150(24.8)
Third (>28weeks)	350(57.8)
Gravidity (number of pregnancies)	
Primigravidae (1)	199(32.8)

Multigravidae(2-4)	381(62.9)
Grandmultigravidae (>4)	26(4.3)

116
117

118 **Prevalence of HIV-1**

119 The overall prevalence of HIV-1 amongst the participants was 70 (11.6%). The age group >35 years had
120 the highest HIV-1 prevalence (18.6%), while women <21 years had the lowest prevalence (5.7%). This
121 difference was statistically significant ($p = 0.001$). The youngest seropositive pregnant woman was aged
122 17 years and the oldest was 42 years of age. HIV prevalence in pregnant women was relatively high
123 among married women 9(15.5%) and among women who had attended primary schools though the
124 difference was not significant ($p = 0.45$). Furthermore, the prevalence of HIV-1 was significantly high ($P =$
125 0.02) among women who started antenatal care at first trimester 20(18.9) and insignificantly high ($P =$
126 0.74) among grand multigravidae 6(23.1) women (Table 2).

127
128
129
130

Table 2: Univariate analysis of HIV prevalence according to socio-demographic and obstetrical characteristics

Demographic characteristics	Number (%)	HIV neg (%)	HIV pos (%)	OR (95% CI)	P value
Age group in Years					
<21	53(8.7)	50(94.3)	3(5.7)	0.22 (1.38-3.54)	0.001
21-30	397(65.5)	359(90.4)	38(9.6)		
>30	156(25.7)	127(81.4)	29(18.6)		
Marital status					
Single	164(27.1)	143(87.2)	21(12.8)	0.92 (0.6.3-1.34)	0.65
Concubine	58(9.6)	324(89.5)	38(10.5)		
Married	362(59.7)	49(84.5)	9(15.5)		
Widow	22(3.6)	20(90.9)	2(9.1)		
Level Educational					
None	17(2.8)	15(88.2)	2(11.8)	0.86 (0.62-1.23)	0.45
Primary	107(17.7)	93(86.9)	14(13.1)		
Secondary	301(49.7)	267(88.7)	34(11.3)		
Tertiary	181(29.9)	161(89.0)	20(11.0)		
Gestational age classification					
First (<14weeks)	106(17.5)	86(81.1)	20(18.9)	0.69 (0.51-0.97)	0.021
Second (14-28weeks)	150(24.8)	132(88.0)	18(12.0)		
Third (>28weeks)	350(57.8)	318(90.9)	32(9.1)		
Gravidity (number of pregnancies)					
Primi gravidae (1)	199(32.8)	175(87.9)	24(12.1)	1.09 (0.65-1.85)	0.74
Muilti gravidae(2-5)	381(62.9)	341(89.5)	40(10.5)		
Grand multigravidae (>5)	26(4.3)	20(76.9)	6(23.1)		

131 **OR: odds ratio**

132
133

134 **Prevalence of *T. gondii***

135 Considering women with either anti-Toxoplasma IgG or anti-Toxoplasma IgM or both anti-Toxoplasma
136 IgG and IgM, 139(22.9%) women presented with toxoplasmosis. Of the 606 participants, 135 (22.3%)
137 were found seropositive for anti-Toxoplasma IgG antibodies, while 11(1.8%) had anti-Toxoplasma IgM.

138 This difference was statistically significant (p=0.00). Seven (5.2%) of the women tested positive for both
 139 IgG and IgM anti-Toxoplasma. Univariate analyses of demographic and obstetrical characteristics
 140 showed no significant difference. However, the prevalence was highest among age group <21years
 141 (24.5%; P = 0.37), married women (25.9%; P = 0.46), women who had attained tertiary level of education
 142 (26.0%; P = 0.16), 1st trimester women (29.2%; P= 0.65), and primigravidae women (28.1%; P = 0.07)
 143 (Table 3)

144
 145
 146
 147
 148
 149

Table 3: Univariate analyses of demographic and obstetrical characteristics of *T. gondii* antibodies

characteristics	<i>T. gondii</i> negative	<i>T. gondii</i> positive	Crude odds ratio	95% CI	P value
Age group (Year)					
<21	40(75.5)	13(24.5)	0.83	0.56-1.25	0.37
21-30	303(76.3)	94(23.7)			
>30	124(79.5)	32(20.5)			
Marital status					
Single	135(82.3)	29(17.7)	1.14	0.81-1.59	0.46
Married	271(74.9)	91(25.1)			
Concubine	43(74.1)	15(25.9)			
Widow	18(81.8)	4(18.2)			
Level Educational					
None	14(82.4)	3(17.6)	1.24	0.91-1.70	0.16
Primary	89(83.2)	18(23.6)			
Secondary	230(76.4)	71(23.6)			
Tertiary	134(74.0)	47(26.0)			
Gestational age classification					
First (<14weeks)	75(70.8)	31(29.2)	0.94	0.69-1.26	0.65
Second (14-28weeks)	112(74.7)	38(25.3)			
Third (>28weeks)	280(80.0)	70(20.0)			
Gravidity					
Primigravidae (1)	143(71.9)	56(28.1)	0.63	0.37-1.04	0.07
Multigravidae(2-5)	301(79.0)	80(21.0)			
Grandmultigravidae (>5)	23(88.5)	3(11.5)			

150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167

***T. gondii* and HIV-1 and co-infection**

Of the 606 women, 31(5.1%) were positive for both HIV and *T. gondii*. Co-infection rate was high among women of the age group 21-35 years (5.5%), single women 8(17.7%), women who never went to school (5.9%), women who started antenatal cars at a gestational age of < 14 weeks (8.7%) and women with more than 5 pregnancies (7.7%). This differences were however not significant p>0.05 (Table 4).

168 **Table 4: univariate analyses of socio-demographic and obstetrical data of HIV and *T. gondii* co-**
 169 **infection**

Demographic characteristics	Number (%)	Co-infection	P value
Age group (Year)			
<21	53(8.7)	1(1.9)	0.53
21-30	397(65.5)	22(5.5)	
>30	156(25.7)	8(5.1)	
Marital status			
Single	164(27.1)	8(17.7)	0.30
Concubine	362(59.7)	16(4.4)	
Married	58(9.6)	6(10.3)	
Widow	22(3.6)	1(4.5)	
Level Educational			
None	17(2.8)	1(5.9)	0.91
Primary	107(17.7)	4(3.7)	
Secondary	301(49.7)	16(5.3)	
Tertiary	181(29.9)	10(5.5)	
Gestational age classification			
First (<14weeks)	106(17.5)	9(8.5)	0.07
Second (14-28weeks)	150(24.8)	10(6.7)	
Third (>28weeks)	350(57.8)	12(3.4)	
Gravidity			
Primigravidae (1)	199(32.8)	10(5.0)	0.83
Multigravidae(2-5)	381(62.9)	19(5.0)	
Grandmultigravidae (>5)	26(4.3)	2(7.7)	

170

171 3.2 DISCUSSION

172 This study is one of the few studies carried out in Cameroon to explore the risk factors associated with *T.*
 173 *gondii* and the seroprevalence of *T. gondii* and HIV co-infection among pregnant women in Bamenda
 174 Health District.

175
 176 Despite the low HIV prevalence (4.3%) in Cameroon in 2016 [22], the prevalence of HIV (11.6%) among
 177 pregnant women was high compared to the 0.5-10.3% range reported in other countries of the world [14,
 178 17, 23] and from other towns in Cameroon [20, 21]. However, it was quite low compared to the 61.6%
 179 reported by Simporé *et al.*, [11] in a study carried out in Burkina Faso.

180
 181 There was no statistical significance between marriage, level of education, gravidity and HIV-1 infection
 182 ($p < 0.05$). However, the risk of contracting the disease was 78%, 8%, 14%, and 31% less likely to occur
 183 in women of the age group >30years, widows, those who attain secondary school, and women who
 184 started ANC at third trimester respectively, while the risk was 9% more likely to occur in grand
 185 multigravidae women. In this study, HIV-1 prevalence was highest among the age group >30 years
 186 contrary to the previous studies [12, 23] which state that HIV-1 prevalence was high in the 21–25years
 187 age range. Similar results have been reported in a different town in Cameroon [16]. This is most likely due
 188 to progressive increase duration of exposure to sexual activity in this age group compared to a lower age
 189 group. In addition, the majority of the women, in the >30years age range were multigravida or grand
 190 multigravida indicating that they have been exposed more to unprotected sexual intercourse which is a
 191 risk factor for HIV infection.

192 As reported in other studies [19, 23, 24] married women had a high HIV prevalence (15.5%). However, is
 193 contrary to another study from a different town in Cameroon where single women were more infected
 194 [16]. It has been reported that susceptibility and vulnerability to HIV/AIDS are attributed to marital and
 195 family status [25]. This high HIV prevalence in the group of women is associated with the fact that married
 196 women usually have unprotected sex and in addition, it was difficult to assess information on multiple
 197 partners in these women although extramarital affairs are common in the said setting.

198 Data from this study showed that woman who had attained primary education had the highest HIV
 199 prevalence (13.1%) followed by those who did not go to school (11.8%). This may be attributed to lack of
 200 adequate information on the mode of transmission and prevention of HIV and other STDs. This result is

201 similar to studies by [17, 23] and contrary to other studies where women with tertiary education had a
202 higher HIV seroprevalence [19, 26].

203 Prevalence of HIV in this study was insignificantly high among women who were multigravida similar to
204 report by Nayak *et al.*, [23] and contrary to the previous study done by Patil *et al.*, [25] where HIV was
205 common among primigravida. The high prevalence is associated with increased risk of unprotected
206 sexual intercourse in this group of women

207
208 The seroprevalence of *T. gondii* infection was 22.9% while seroprevalence for anti-Toxoplasma IgG and
209 IgM, antibodies were 22.3% and 1.8% respectively. The risk of contracting Toxoplasma was 93%, 16%,
210 and 37% less likely to occur in the age group >30years, women who started ANC at second trimester and
211 grand multigravidae women respectively, while the risk was 14% and 24% more likely to occur in women
212 in a concubine relationship and women who had attain tertiary education respectively. The
213 seroprevalence of *T. gondii* infection in this study was found to be lower than the 30-90% range reported
214 in different countries [6, 8, 12] and was higher compared to the 5.9- 18.5% range in other studies [4, 27,
215 28]. In Cameroon, previous studies have reported a range of 54.4-77.1% [5, 10, 29]. The differences seen
216 with other studies can be attributed to environmental or climatic conditions favoring the transmission and
217 infectivity of *T. gondii* oocysts, diagnostic methods, living styles, standards of the people, sampled
218 populations, cultural characteristics, personal hygienic practice, feeding habits and genetic background
219 [6, 10, 30]. This decrease in prevalence can be as a result of the awareness that is been created from the
220 result of previous studies.

221
222 Detection of both IgG and IgM simultaneously helps to establish the chronological status of *T. gondii* [31].
223 Toxoplasma IgG antibodies indicate a chronic infection while Toxoplasma IgM antibodies indicate an
224 acute infection [8, 12]. The high prevalence of Toxoplasma IgG compared to Toxoplasma IgM antibodies
225 seen in this study have been reported elsewhere [7, 30, 32]. The low IgM (5.2%) antibodies might indicate
226 that the IgM antibodies present is from a previous infection [33]. The presence of IgM antibody during
227 pregnancy indicates the presence of an acute *T. gondii* infection which is associated with a higher risk of
228 maternal-fetal transmission [7]. Thus the early diagnosis of Toxoplasmosis in pregnant mothers is of great
229 importance for early initiation of measures and therapy that reduce the risk of transmission and possible
230 consequence on the newborn. However, other studies did not report the presence of *T. gondii*-specific
231 IgM [8, 11, 12]. However, it has been reported that IgM antibody is usually detected within the first two
232 weeks of infection and reduces to negligible levels within 6 months after exposure. As such the presence
233 of IgM may not be an acute infection but for the fact that it can persist for prolonged times after infection
234 [28, 29].

235
236 Other studies have reported that the risk of contracting *T. gondii* infection increases with age unlike the
237 case in this study [34, 35]. Though age was not a risk factor to the *T. gondii* infection, younger women
238 <21 years were more infected compared to older women. This result contradicts studies by [4, 6, 28] that
239 identify age group > 21 years as a risk factor. In addition, the result is similar to studies by Njunda *et al.*,
240 [10] and Shimelis *et al.*, [36] which state that seroprevalence of *T. gondii* does not depend on age.
241 Nevertheless, another study in Cameroon indicates that women aged between 31-35years had a higher
242 prevalence [10]. The variation in age classification of the different studies can also account for the
243 variation of the results seen in the different studies. The high prevalence in younger women can be
244 attributed to their lifestyle. It has been reported that younger people are more exposed to activities like
245 grilled meat or fish which might be undercooked as well as raw food like fruits and salad which may be
246 contaminated with the parasites hence increased risks of infection [12, 13].

247
248 In this study no significant association was found between the seroprevalence of toxoplasmosis and
249 educational status as opposed to a study by da Silva *et al.*, [1] who reported low education or illiteracy as
250 a risk factor. A similar finding was recorded by Walle *et al.*, [31]. On the contrary women with tertiary
251 school education which suggests a better understanding of hygiene principles had the highest prevalence
252 of toxoplasmosis. The high prevalence in this group can be attributed with higher socioeconomic
253 standards such as eating of raw vegetables, fruit and roasted meats which have been identified sources
254 of disease transmission.

255 The degree of severity of the disease depends on the gestational age as severe fetal affection occurred
256 with early gestational age infection [32]. Gestational age did not show any significant association as also
257 reported by Frimpong *et al.*, [4] in another study. Contrary to this study, data presented by Shao *et al.*,
258 [13] showed that gestational age was a significant risk factor. The highest seroprevalence of Toxoplasma

259 antibodies (29.2%) was found in pregnant women at the first trimester is similar to the result of Alsammani
260 [35] contrary to second and third semesters [4, 24].
261 Despite the non-statistical significant association contrary to another study [6], data from this study
262 showed that the risk of toxoplasmosis decreases with increase in gravidity. Primigravidae recorded the
263 highest prevalence of 56(28.1%). This result is contrary to other studies by Awoke *et al.*, [28] and Negero
264 *et al.*, [6] which state that *T. gondii* is more likely to occur in multigravidae. The likely reason for this result
265 is that the test for *T. gondii* has been encouraged for more than 5 years in this setting. As such women
266 with multiple pregnancies are were knowledgeable with the method of prevention than primigravidae
267 women. Secondly, previously infected women must have been treated prior to the present pregnancy.
268
269

270 In this study, no significant difference was seen between seropositivity of *T. gondii* in HIV positive
271 31(44.3%) and negative 108(20.1%) women similar to studies in other countries [4, 14, 24]. The high
272 prevalence in this group is as a result of decreased immunity which leads to reactivation of latent
273 infection/tissue toxoplasmosis in HIV positive women [9, 30]. The reason for the non-significance in this
274 study can be as a result of the use of antiretroviral therapy (ART). ART suppresses HIV viral replication
275 and increased CD4⁺ T-cell counts, therefore, preventing the development of opportunistic infections. In
276 addition, since 2012, Bamenda health district in Cameroon has been implementing the test and treat
277 method (option B+) where all HIV pregnant or breastfeeding mothers are placed on ART irrespective of their
278 CD4⁺ T cell counts or clinical stage [18]. On the other hand studies by Siteo *et al.*, [37] and Walle *et al.*,
279 [31] showed a significant difference in the prevalence rate between HIV positive and negative women.
280

281 In this study we recorded a prevalence of 5.1% co-infection rates lower than the 12- 25% range in other
282 studies [5, 12] but higher compared to the 2.1% reported by Fernandes *et al.*, [14]. HIV and *T. gondii* co-
283 infection rate are common in pregnant women because both pregnancy and HIV weakens the immune
284 system that favors *T. gondii* and other opportunistic infection to occur [11, 12]. Furthermore, it is more
285 likely that these women with co-infection were recently diagnosed with HIV and were not on treatment or
286 were newly initiated on treatment. In addition, HIV-1 and *T. gondii* co-infection could be attributed to
287 common social lifestyle or associated risk factors common to both infections, such as exposure to sexual
288 contacts, consumption of undercooked meat or roasted meat and raw vegetable.
289

289 **Limitation**

290 The present study has certain limitations that need to be taken into account. No CD4⁺ T cell count was
291 measured, History on ART was not taken into consideration, or the year of HIV diagnosis was not known
292 by most women.
293
294

295 **4. CONCLUSIONS**

296 This study demonstrates that the prevalence of *T. gondii* infection among pregnant women is decreasing.
297 The high prevalence of *T. gondii* and HIV co-infection among pregnant women indicates a greater
298 probability of congenital transmission of *T. gondii*.
299

300 **RECOMMENDATION**

301 The high prevalence of *T. gondii* and HIV-1 co-infection indicate the need to intensify the education of the
302 associated risk factors of both *T. gondii* and HIV-1 infections and methods of prevention. This will reduce
303 the risk of mother to child transmission and thus prevent the consequences of toxoplasmosis and HIV in
304 children. In addition, serological screening for *T. gondii* infection should be considered as part of an
305 antenatal investigation during ANC follow-up.
306

307 **COMPETING INTERESTS**

308
309 We have no competing interest

310
311 **CONSENT**

312
313 We declare that oral informed consent was obtained from the patients for **publication without** disclosure of
314 identity

315
316
317 **ETHICAL APPROVAL**

318
319 Ethical clearance and administrative authorization were obtained from the ethical review board of the
320 delegation of Public Health Bamenda and Bamenda Regional Hospital review **board.**

321
322
323 **REFERENCES**

- 324 1. da Silva MG, Vinaud MC, de Castro AM. Prevalence of toxoplasmosis in pregnant women and
325 vertical transmission of *Toxoplasma gondii* in patients from basic units of health from Gurupi,
326 Tocantins, Brazil, from 2012 to 2014. *PLoS One*. 2015 Nov 11;10(11):e0141700.
- 327 2. Nguefack HL, Gwet H, Desmonde S, Oukem-Boyer OO, Nkenfou C, Téjiokem M, et al.
328 Estimating mother-to-child HIV transmission rates in Cameroon in 2011: a computer simulation
329 approach. *BMC infectious diseases*. 2015 Dec;16(1):11.
- 330 3. Dawit G, Shishay K. Epidemiology, public health impact and control methods of the most
331 neglected parasite diseases in Ethiopia: a review. *World J Med Sci*. 2014;10:94-102.
- 332 4. Frimpong C, Makasa M, Sitali L, Michelo C. Seroprevalence and determinants of toxoplasmosis
333 in pregnant women attending antenatal clinic at the university teaching hospital, Lusaka, Zambia.
334 *BMC infectious diseases*. 2017 Dec;17(1):10.
- 335 5. Wam EC, Sama LF, Ali IM, Ebile WA, Aghangu LA, Tume CB. Seroprevalence of *Toxoplasma*
336 *gondii* IgG and IgM antibodies and associated risk factors in women of child-bearing age in
337 Njinikom, NW Cameroon. *BMC research notes*. 2016 Dec;9(1):406.
- 338 6. Negero J, Yohannes M, Woldemichael K, Tegegne D. Seroprevalence and potential risk factors
339 of *T. gondii* infection in pregnant women attending antenatal care at Bonga Hospital,
340 Southwestern Ethiopia. *International Journal of Infectious Diseases*. 2017 Apr 1;57:44-9.
- 341 7. Yohanes T, Zerdo Z, Chufamo N. Seroprevalence and predictors of hepatitis B virus infection
342 among pregnant women attending routine antenatal care in Arba Minch Hospital, South Ethiopia.
343 *Hepatitis research and treatment*. 2016;2016.
- 344 8. Paul E, Kiwelu I, Mmbaga B, Nazareth R, Sabuni E, Maro A, et al. *Toxoplasma gondii*
345 seroprevalence among pregnant women attending antenatal clinic in Northern Tanzania. *Tropical*
346 *medicine and health*. 2018 Dec;46(1):39.
- 347 9. Nazari N, Bozorgomid A, Janbakhsh A, Bashiri F. *Toxoplasma gondii* and human
348 immunodeficiency virus co-infection in western Iran: A cross-sectional study. *Asian Pacific*
349 *Journal of Tropical Medicine*. 2018 Jan 1;11(1):58.
- 350 10. Njunda AL, Assob JC, Nsagha DS, Kamga HL, Nde PF, Yugah VC. Seroprevalence of
351 *Toxoplasma gondii* infection among pregnant women in Cameroon. *Journal of public health in*
352 *Africa*. 2011 Sep 5;2(2).
- 353 11. Simpore J, Savadogo A, Ilboudo D, Nadambega MC, Esposito M, Yara J, et al. *Toxoplasma*
354 *gondii*, HCV, and HBV seroprevalence and co-infection among HIV-positive and-negative
355 pregnant women in Burkina Faso. *Journal of medical virology*. 2006 Jun;78(6):730-3.
- 356 12. Okwuzu JO, Odunukwe NN, Ezechi OC, Gbajabiamila TA, Musa AZ, Ezeobi PM, et al.
357 *Toxoplasma gondii* infection in HIV/AIDS: prevalence and risk factors. *African Journal of Clinical*
358 *and Experimental Microbiology*. 2014;15(2):97-102.
- 359 13. Shao ER, Ndazana SG, Chacha W, Masenga G, Tolbert S, Mosha D, et al. Sero-Prevalence and
360 Factors Associated with *Toxoplasma Gondii* Infection among Pregnant Women Attending
361 Antenatal Care in the Referral Hospital in Tanzania: Cross Sectional Study *iMedPub Journals*.
362 *Annals of Clinical and Laboratory Research*. 2015;3(2):17.

- 363 14. Fernandes MA, Batista GI, Carlos JD, Gomes IM, Azevedo KM, Setúbal S, Oliveira SA, et al.
364 Toxoplasma gondii antibody profile in HIV-1-infected and uninfected pregnant women and the
365 impact on congenital toxoplasmosis diagnosis in Rio de Janeiro, Brazil. *Brazilian Journal of*
366 *Infectious Diseases*. 2012 Apr;16(2):170-174.
- 367 15. Wang ZD, Wang SC, Liu HH, Ma HY, Li ZY, Wei F, et al. Prevalence and burden of Toxoplasma
368 gondii infection in HIV-infected people: a systematic review and meta-analysis. *The Lancet HIV*.
369 2017 Apr 1;4(4):e177-88.
- 370 16. de Dieu Anoubissi J, Gabriel EL, Nde CK, Fokam J, Tseuko DG, Messeh A, Moussa Y, Nkenfou
371 CN, Bonono L, Billong SC, Nfetam JB. Factors Associated with Risk Of HIV-Infection Among
372 Pregnant Women in Cameroon: Evidence from the 2016 National Sentinel Surveillance Survey of
373 HIV and syphilis. *BioRxiv*. 2018 Jan 1:482901.
- 374 17. Yang S, Yang C, Liao Q, Zhai W, Yu G, Xiao L, et al. Analysis of HIV prevalence among
375 pregnant women in Liangshan Prefecture, China, from 2009 to 2015. *PloS one*. 2017 Sep
376 7;12(9):e0183418.
- 377 18. Kalua T, Barr BA, Van Oosterhout JJ, Mbori-Ngacha D, Schouten EJ, Gupta S, et al. Lessons
378 learned from option B+ in the evolution toward "test and start" from Malawi, Cameroon, and the
379 United Republic of Tanzania. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 2017
380 May 1;75: S43-50.
- 381 19. Kiptoo M, Mpoke S, Mueke J, Okoth F, Songok E. Survey on prevalence and risk factors on HIV-
382 1 among pregnant women in North-Rift, Kenya: a hospital-based cross-sectional study conducted
383 between 2005 and 2006. *BMC international health and human rights*. 2009 Dec;9(1):10.
- 384 20. Dionne-Odom J, Mbah R, Rembert NJ, Tancho S, Halle-Ekane GE, Enah C, et al. Hepatitis B,
385 HIV, and syphilis seroprevalence in pregnant women and blood donors in Cameroon. *Infectious*
386 *diseases in obstetrics and gynecology*. 2016;2016.
- 387 21. Sama CB, Fetei VF, Tindong M, Tanyi JT, Bihle NM, Angwafo III FF. Prevalence of maternal HIV
388 infection and knowledge on mother-to-child transmission of HIV and its prevention among
389 antenatal care attendees in a rural area in northwest Cameroon. *PloS one*. 2017 Feb
390 15;12(2):e0172102.
- 391 22. Teto G, Tagny CT, Mbanya D, Fonsah JY, Fokam J, Nchindap E, et al. Gag P2/NC and pol
392 genetic diversity, polymorphism, and drug resistance mutations in HIV-1 CRF02_AG-and non-
393 CRF02_AG-infected patients in Yaounde, Cameroon. *Scientific reports*. 2017 Oct 26;7(1):14136.
- 394 23. Nayak AK, Jain MK, Dhivya S, Hota S. A study on the prevalence of HIV infection among
395 pregnant women attending antenatal clinic in a tertiary care hospital, Cuttack, India. *International*
396 *Journal Of Community Medicine And Public Health*. 2017 Apr 24;4(5):1504-9.
- 397 24. Gelaye W, Kebede T, Hailu A. High prevalence of anti-toxoplasma antibodies and absence of
398 Toxoplasma gondii infection risk factors among pregnant women attending routine antenatal care
399 in two Hospitals of Addis Ababa, Ethiopia. *International Journal of Infectious Diseases*. 2015 May
400 1;34:41-5.
- 401 25. Patil VM, Moray AP, Patil SP. Ten years trend of HIV seroprevalence among Indian Pregnant
402 women attending antenatal clinic at a tertiary hospital in Dhule, Maharashtra. India. *IJRCOG*.
403 2016;5(5):1514-9.
- 404 26. Smith J, Nalagoda F, Wawer MJ, Serwadda D, Sewankambo N, Konde-Lule J, et al. Education
405 attainment as a predictor of HIV risk in rural Uganda: results from a population-based study.
406 *International journal of STD & AIDS*. 1999 Jul 1;10(7):452-9.
- 407 27. Cong W, Dong XY, Meng QF, Zhou N, Wang XY, Huang SY, et al. Toxoplasma gondii infection in
408 pregnant women: a seroprevalence and case-control study in Eastern China. *BioMed research*
409 *international*. 2015;2015.
- 410 28. Awoke K, Nibret E, Munshea A. Sero-prevalence and associated risk factors of Toxoplasma
411 gondii infection among pregnant women attending antenatal care at Felege Hiwot Referral
412 Hospital, northwest Ethiopia. *Asian Pacific journal of tropical medicine*. 2015 Jul 1;8(7):549-54.
- 413 29. Ndumbe PM, Andela A, Nkemnkeng-Asong J, Watonsi E, Nyambi P. Prevalence of infections
414 affecting the child among pregnant women in Yaounde, Cameroon. *Medical microbiology and*
415 *immunology*. 1992 Jul 1;181(3):127-30.
- 416 30. Yusuf AM, Yahaya S, Azeez-Akande O. Seroprevalence and risk factors of Toxoplasma gondii
417 infection (toxoplasmosis) among HIV seropositive pregnant women in a tertiary healthcare centre,
418 Kano, Northern Nigeria. *J. Med. Med. Sci*. 2016 Feb;7:001-5.

- 419 31. Walle F, Kebede N, Tsegaye A, Kassa T. Seroprevalence and risk factors for Toxoplasmosis in
420 HIV infected and non-infected individuals in Bahir Dar, Northwest Ethiopia. *Parasites & vectors*.
421 2013 Dec;6(1):15.
- 422 32. Kamal AM, Ahmed AK, Abdellatif MZ, Tawfik M, Hassan EE. Seropositivity of toxoplasmosis in
423 pregnant women by ELISA at Minia University Hospital, Egypt. *The Korean journal of*
424 *parasitology*. 2015 Oct;53(5):605.
- 425 33. Chaudhry SA, Gad N, Koren G. Toxoplasmosis and pregnancy. *Canadian Family Physician*. 2014
426 Apr 1;60(4):334-6.
- 427 34. Elsheikha HM, Aboul-Dahab MA, Abdel AM, El-Sherbini ET. Prevalence and risk factors of
428 *Toxoplasma gondii* antibodies in asymptomatic Egyptian blood donors. *Journal of the Egyptian*
429 *Society of Parasitology*. 2009 Apr;39(1 Suppl):351-61.
- 430 35. Alsammani MA. Sero-epidemiology and risk factors for *Toxoplasma gondii* among pregnant
431 women in Arab and African countries. *Journal of Parasitic Diseases*. 2016 Sep 1;40(3):569-79.
- 432 36. Shimelis T, Tebeje M, Tadesse E, Tegbaru B, Terefe A. Seroprevalence of latent *Toxoplasma*
433 *gondii* infection among HIV-infected and HIV-uninfected people in Addis Ababa, Ethiopia: A
434 comparative cross-sectional study. *BMC Research Notes*. 2009 Dec;2(1):213.
- 435 37. Siteo SP, Rafael B, Meireles LR, Andrade Jr HF, Thompson R. Preliminary report of HIV and
436 *Toxoplasma gondii* occurrence in pregnant women from Mozambique. *Revista do Instituto de*
437 *Medicina Tropical de São Paulo*. 2010 Dec;52(6):291-5.

438
439
440

441 **DEFINITIONS, ACRONYMS, ABBREVIATIONS**

442 ART: antiretroviral therapy, AOR: adjusted odds ratio, CI: confidence interval, HIV: human
443 immunodeficiency virus, *T: Toxoplasma*

444

445