

**EFFICACY OF POULTRY DROPPINGS, COW DUNG, SAW DUST AND NPK FERTILIZER ON THE GROWTH, YIELD COMPONENTS AND YIELD OF *Glycine max. L* (SOYA BEAN) IN THE NORTHERN GUINEA SAVANNA REGION OF NIGERIA**

**ABSTRACT**

This research study was conducted at the Federal College of Forestry, Jos demonstration farm to determine the efficacy of poultry droppings, cow dung, saw dust and Npk fertilizer on the growth, yield components and yield of *Glycine Max. L.* (Soya Bean). Randomised complete block design (RCBD) was used with five treatments replicated four times. Growth and yield characteristics were recorded on plant height, number of leaves, stem girth, number of branches, leaf area, days to 50% flowering, number of pods/plant, number of seeds/pod, 1000 seeds weight, yield and biomass weight. The result shows that there was significant difference at the application of the treatments. poultry droppings differed significantly as it gave the highest plant height (54.78cm), number of leaves (50.40), number of branches (3.80), leaf area (143.80cm<sup>2</sup>), stem girth (3.26cm), number of pods/plant (95.00), number of seeds/pod (2.62), 1000 seeds weight (420.80g), yield (24.84 tons/ha) and biomass (15.66 tons/ha). The least number of days it takes for 50% of the seeds to flowering was also recorded on the application of poultry manure. It can be concluded that the application of poultry droppings gave the highest growth and yield characteristics of soya bean in the study area. Thus, soya bean farmers are encourage to apply poultry droppings for better growth and yield.

**Key Words:** Soya bean, *Glycine max. L.*, Growth Components, Yield components, Fertilizers.

**1.0 INTRODUCTION**

Soya beans (*Glycine max. (L.) Merr.*) are one of the oldest cultivated crops of the temperate regions and one of the world's most important sources of oil and protein. Available records indicate that Soya beans originated from china and from there it spread to all parts of the world (Onwueme and Sinha, 1991). The annual average world soybean production was about 125.7 million metric tons, corresponding 50% of this amount to North America, 31.5% to Central and South America, 16% to Asia and the remaining 2.5% to Africa, Europe and Oceania (Islas-Rubio and Higuera-Ciapara, 2002). In Africa, Nigeria is the largest producer of soybean with an annual turnover of about 500,000 metric tons (Agronewsng, 2016). Soya bean seeds contain 43.2% protein, 19.5% fat, 20.9% carbohydrate and a good amount of other nutrients like calcium, phosphorus, iron and vitamins (Khaim et al., 2013). Soya bean has 3% lecithin which is helpful for brain development.

Organic manure is a cheap and readily available source of essential nutrients to the plants. It is used primarily as a source of plant nutrients (Mullins et al., 2002; Samia et al., 2015). Naturally, the use of organic manure can improve soil properties and maintain the quality of soil fertility. Organic manures act not only as a source of nutrients and organic matter, but also increase microbial biodiversity and activity in soil, influence structure, nutrients get turnover and

46 many other changes related to physical, chemical and biological parameters of the soil (Muzafer  
 47 et al., 2015). The soil having higher organic matter concentrations have been proved to enhance  
 48 the growth and yield of different crops (Sarwar, 2005; Muzafer et al., 2015) as well as soil  
 49 aeration, soil density and maximizing water holding capacity of soil for seed germination and  
 50 plant root development.

51  
 52 In spite of the increase in land areas under soya bean production, yield is still low. Some  
 53 of the major causes of low yields are declining soil fertility and insufficient use of fertilizers  
 54 resulting in severe nutrient depletion of soils. In the past, a long fallow period (5-10 years)  
 55 allowed natural restoration of soil fertility. However, because of pressure on land to  
 56 increase food production and other socio-economic activities, the fallow period is almost  
 57 nonexistent in many farming communities in Nigeria. Fertilizer/manure has been shown to be  
 58 an effective means of enhancing crop performance for more than a century. It has contributed  
 59 largely to the major increase in yields which have been achieved worldwide and for the  
 60 substantial improvement of human and animal health.

61  
 62 Soya bean being a high protein and energy crop its productivity is often limited due to  
 63 poor yield. The interest in soybean has recently been increased, and a lot of researches have been  
 64 conducted due to the increasing demand for soya bean both for domestic and industrial purposes.  
 65 Thus, this research work is carried out to determine the efficacy of poultry droppings, cow dung,  
 66 saw dust and npk fertilizer on the growth, yield components and yield of *Glycine max. L* (soya  
 67 bean) in the northern guinea savanna region of Nigeria.

68

69 **2.0 Materials and Methods**

70

71 The field experiment was carried out between June to August, 2018 at the Federal  
 72 College of Forestry demonstration farm located in Jos, Plateau state. The region lies between  
 73 latitude 7° and 11° north, longitude 7° and 25° east and at an altitude of about 1200km above sea  
 74 level. The area lies in the northern guinea savanna of Nigeria with an annual rainfall of 1460mm  
 75 and a temperature of 19°C to 32°C, (Olowolafe et al., 2004).

76

77 **2.1 Soil Analysis**

78 Soil samples from the study area were collected randomly at a depth of 0cm to 30cm to  
 79 determine the physical and chemical properties. A soil analysis was carried out at ASTC  
 80 (Agricultural Services and Training Center) KASSA, VOM, Jos, Plateau sate.

81

82 **Table 1: Physical and Chemical Properties of Soil in the Study Area**

Sample	PH	N (%)	P PPM	K PPM	Ca PM	MgPPM	O.M(%)	H+ mMol/ 100g	Clay (%)	Silt (%)	Sand (%)	Textural Class
0- 15cm	5.8	0.04	6.2	96.0	530	102	115	157x10	10.88	12	77.12	Sandy loam

83 **Source:** - Agricultural Services and Training Center KASSA/VOM, 2018.

The physical and chemical properties of the soil as presented in Table 1 showed that the soil PH was 5.8 which is slightly acidic. It is the preferred soil PH range for good growth and development of most crops. Organic matter had an average value of 115%, while the respective nutrient constituents of nitrogen, phosphorus, potassium, calcium and magnesium were 0.04%, 6.2, 96.0, 530 and 102ppm were in average quantities for optimum production of most crops. The soil can be classified as sandy loam. The percentage composition of sand, silt and clay (10.88% clay, 12% silt, and 77.12% sand) confirms that the presences of organic matter, which make the soil good for crop production.

The experiment was laid out on a randomised complete block design with five treatments (control, poultry droppings 2.5t/ha, cow dung 2.5t/ha, saw dust 2.5t/ha and NPK fertilizer 180Kg/ha) replicated four times. The seeds were obtained at IITA kano and planted at the rate of two seeds per hole. Growth and yield characteristics were recorded on plant height, number of leaves, stem girth, number of branches, leaf area, days to 50% flowering, number of pods/plant, number of seeds/pod, 1000 seeds weight, yield and biomass weight.

### 3.0 Results and Discussions

**Table 2: Efficacy Of Poultry Droppings, Cow Dung, Saw Dust And Npk Fertilizer On The Growth *Glycine max. L* (SOYA BEAN)**

Treatment	Plant Height (cm)	Number of Leaves	Number of Branches	Leaf Area (cm <sup>2</sup> )	Stem Girth (cm)
Control	32.14a	31.00a	1.60a	100.40a	1.92a
Saw Dust	36.18b	36.00b	2.60b	102.40a	2.64b
NPK Fertilizer	38.56c	43.80c	2.20ab	132.00c	2.52b
Cow Dung	45.92d	45.60d	2.80b	121.80b	2.56b
Poultry Dropping	54.78e	50.40e	3.80c	143.80d	3.26c
SE±	0.98	0.79	0.37	1.70	0.23
LSD	**	**	**	**	**

Source: Field Experiment 2018

Means within a column having same letters are not significantly different at  $P \leq 0.05$ .

LS = level of significance

\* = Significant at 0.05

\*\* = Significant at 0.001

**3.1 Plant Height:** The efficacy of poultry droppings, cow dung, saw dust and Npk fertilizer on plant height as presented in table 2 indicates the there is significant difference between the treatments at both 1% and 5% levels of probability. The highest mean plant height was 54.78cm given by the application of poultry droppings, the application of cow dung gave 45.92cm, Npk fertilizer gave 38.56cm while saw dust and the control gave 36.18cm and 32.14cm respectively.

**3.2 Number of Leaves:** The efficacy of poultry droppings, cow dung, saw dust and Npk

119 fertilizer on number of leaves as shown from table 2 indicates the there is significant difference  
 120 between the treatments at both 1% and 5% levels of probability. The highest (50.40) mean  
 121 number of leaves was obtained at the application of poultry droppings, followed by the  
 122 application of cow dung (45.60), then Npk fertilizer (43.80), with saw dust (36.00) and the  
 123 control (31.00) producing the least number of leaves.

124  
 125 **3.3 Number of Branches:** The result from table 2 shows that significant differences exists  
 126 between the treatments on the number of branches at 1% and 5% level of probability. The  
 127 application of poultry droppings produced the highest (3.80) mean number of branches while the  
 128 control produced the least (1.60) mean number of branches.

129  
 130 **3.4 Leaf Area:** The efficacy of poultry droppings, cow dung, saw dust and Npk fertilizer on leaf  
 131 area as shown in table 2 revealed that significant differences exist between the treatments.  
 132 Poultry droppings gave the highest mean leaf area of 143.80cm<sup>2</sup>. Although no significant  
 133 difference exist between the control and the application of saw dust, it has the least mean leaf  
 134 area of 100.40cm<sup>2</sup>.

135  
 136 **3.5 Stem Girth:** The application of poultry droppings has significant effect (3.26cm) on stem  
 137 girth at 1% and 5% level of probability compared to saw dust (2.64cm), cow dung (2.56), Npk  
 138 fertilizer (2.52cm) and the control (1.92cm).

139  
 140 This result is similar to Samia et al. (2015) who opined that chicken manure fertilizer had  
 141 significant effect on stem diameter, number of branches, plant height and number of leaves. The  
 142 result is also in agreement with Maheshbabu et al. (2008) that all the growth (plant height,  
 143 number of branches, number of leaves and leaf area index), were differed significantly due to the  
 144 application of organic manures. Falodun and Osaigbovo (2010) and Patwary (2003) in Khaim  
 145 (2013) conducted an experiment and reported that the growth characteristics of soya beans were  
 146 enhanced by organic and inorganic fertilizers. Organic manure is a reservoir of nutrients and  
 147 these nutrients are released during humification, thus supplying the necessary elements for plant  
 148 growth (Chiezey and Odunze, 2009).

149  
 150 **Table 3: Efficacy Of Poultry Droppings, Cow Dung, Saw Dust And Npk Fertilizer On The**  
 151 **Yield *Glycine max. L* (SOYA BEAN)**  
 152

Treatment	Days to 50% Flowering	Number of Pods/Plant	Number of Seeds/Pod	1000 Seeds Weight (g)	Yield (tons /ha)	Biomass (tons/ha)
Control	57.60d	72.20a	1.60a	328.00a	14.08a	9.34a
Saw Dust	55.80c	76.00b	2.00ab	353.00b	17.40b	11.62b
NPK	56.80cd	81.20c	1.40ab	356.00b	20.66c	12.96c
Fertilizer						
Cow Dung	52.20b	88.60d	2.20a	377.00c	20.34c	12.76bc
Poultry Dropping	48.20a	95.00e	2.62b	420.80d	24.84d	15.66d
SE±	0.70	0.99	0.36	8.65	0.81	0.61
LS	**	*	**	**	**	**

153 Source: Field Experiment 2018

154 Means within a column having same letters are not significantly different at  $P \leq 0.05$ .

155 LS = level of significance

156 \* = Significant at 0.05

157 \*\* = Significant at 0.001

158

159 **3.6 Days to 50% Flowering:** The efficacy of poultry droppings, cow dung, saw dust and Npk  
160 fertilizer on days to 50% flowering as presented in table 3 indicates the there is significant  
161 difference between the treatments at both 1% and 5% levels of probability. The least number of  
162 days to 50% flowering was obtained at the application of poultry droppings while the control  
163 takes the most number of days to 50% flowering.

164

165 **3.7 Number of Pods/Plant:** The efficacy of poultry droppings, cow dung, saw dust and Npk  
166 fertilizer on number of pods per plant as revealed in table 3 indicates the there is significant  
167 difference between the treatments at both 5% levels of probability. The highest (95.00) number  
168 of pods per plant were obtained on treating soya bean with poultry droppings while the least  
169 (72.00) was obtained when no treatment was given.

170

171 **3.8 Number of Seeds/Pod:** Significant differences were observed among the treatments on  
172 number of seeds per pod as shown in Table 3. The highest number of seeds per plant was  
173 recorded by poultry droppings (2.62) as compared to the other treatments.

174

175 **3.9 1000 Seeds Weight:** The efficacy of poultry droppings, cow dung, saw dust and Npk  
176 fertilizer on 1000 seeds weight as given in table 3 indicates the there is significant difference  
177 between the treatments at both 1% and 5% levels of probability. The largest weight (420.80g)  
178 was obtained at the application of poultry droppings while the control takes the least (328.00g).

179

180 **3.10Yield:** Soya bean yield significantly differed among the various treatments as shown in table  
181 3. The seed yield ranges between 28.84 tons/ha to 14.08tons/ha. All the application of fertilizer  
182 gave significantly higher grain yield over control. The highest soya bean grain yield (24.84  
183 tons/ha) was obtained at the application of poultry droppings, while the least (14.08 tons/ha) at  
184 the control.

185

186 **3.11 Biomass:** The application of poultry droppings gave better (15.66 tons/ha) biomass than the  
187 Saw dust (11.62 tons/ha), Npk fertilizer (12.96 tons/ha), cow dung (12.76 tons/ha) and the  
188 control (9.34 tons/ha). Significant difference was observed at the application of the different  
189 treatments given during the growing seasons in terms of biomass.

190

191 These results is similar to Samia et al. (2015) who opined that chicken manure had positive  
192 influence on growth and yield of soybean which gave the highest means in most growth and  
193 yield attributes. Poultry manure showed better performance in producing grain yield with respect  
194 to other organic manures. Yamika and Ikawati (2012) found that the combination of inorganic  
195 with organic fertilizers increased the seed yield. It can be concluded that a 50% substitution of  
196 inorganic fertiliser with poultry manure is recommended to reduce use of chemical fertilisers  
197 without sacrificing crop yield (Almaz eta al., 2017).

198

199 **4.0 Conclusion**

200 It can be concluded that the application of poultry droppings gave the highest growth and yield  
201 characteristics of soya bean in the study area. Thus, soya bean farmers are encourage to apply  
202 poultry droppings for better growth and yield.

203

204 **Reference**

- 205 1. Onwueme, I. C. and Sinha, T. D. Field Crop Production in Tropical Africa. Published by  
206 CTA, Ede, The Netherlands. 1991.ISBN NO. 9290810866.
- 207 2. AGRONEWSNG. 2016 Soybeans Production and Processing.  
208 [Http://edition.agronewsng.com/agro-business/soybeans-production-and-processing](http://edition.agronewsng.com/agro-business/soybeans-production-and-processing).  
209 Accessed 2/11/2018
- 210 3. Islas-Rubio,A.R. and Higuera-Ciapara, I. SOYBEANS: Post-harvest Operations  
211 Organization: Centro de Investigación y Desarrollo, A.C. (CIAD)  
212 2002.<http://www.fao.org/3/a-ax444e.pdf>. Accessed 2/11/2018
- 213 4. S. Khaim, M. A. H. Chowdhury and B. K. Saha Organic and inorganic fertilization on  
214 the yield and quality of soybean. J. Bangladesh Agril. Univ. 201311(1): 23–28, ISSN  
215 1810-3030
- 216 5. Samia, O. Y., Salam, K. A., Salam, A., Mohammed, M. H., Moaid, A. H.  
217 Effects of organic and mineral fertilizers on growth and yield of soybean ( Glycine max  
218 L. Merrill). International Journal of Agronomy and Agricultural Research (IJAAR) ISSN:  
219 2223-7054 (Print) 2225-3610 (Online) <http://www.innspub.net>. 2015Vol. 7, No. 1, p. 45-  
220 52.
- 221 6. Muzafer Ahmad Sheikh Pinky Dwivedi H. S. Dwivedi , Impact of Chemical  
222 Fertilizer and Organic Manure on the Germination and Growth of Soybean (Glycine max  
223 L.). Advances in Life Science and Technology. www.iiste.org 2015 .  
224 ISSN 2224-7181 (Paper) ISSN 2225-062X (Online) Vol 31
- 225 7. H. M. MAHESHBABU, RAVI HUNJE, N. K. BIRADAR PATIL AND H. B.  
226 BABALAD Effect of Organic Manures on Plant Growth, Seed Yield and Quality of  
227 Soybean. Karnataka J. Agric. Sci. 2008, 21( 2) : ( 219-221)
- 228 8. U. F. Chiezey and A. C. Odunze Soybean response to application of poultry manure and  
229 phosphorus fertilizer in the Sub-humid Savanna of Nigeria. Journal of Ecology and  
230 Natural Environment. 2009Vol. 1(2), pp. 025-031, Available online at  
231 <http://www.academicjournals.org/JENE>. Accessed 12/12/2008.
- 232 9. Yamika, W. S. D., & Ikawati, K. R.). Combination inorganic and organic fertilizer  
233 increased yield production of soybean. American-Eurasian Journal of Sustainable  
234 Agriculture, 2012. 6(1), 14–17.
- 235 10. Almaz, M. G., R. A. Halim and M. Y. Martini Effect of Combined Application of Poultry  
236 Manure and Inorganic Fertiliser on Yield and Yield Components of Maize Intercropped  
237 with Soybean. Pertanika Journal of Tropical Agricultural Science. 2017, 40 (1): 173 -  
238 184
- 239  
240  
241  
242