

A case-control study on the risk of Upper gastrointestinal bleeding in patients taking NSAIDs in Mashhad, Iran

ABSTRACT

Introduction: Gastrointestinal bleeding is one of the most common causes of patient admissions at emergency wards. Despite considering nonsteroidal antiinflammatory drugs (NSAIDs), aspirin and Helicobacter pylori as the leading causes, mortality from gastrointestinal (GI) bleeding is still high. So pattern of NSAID consumption and related conditions may help in preventative behavior.

Methods: This case-control study was conducted on 300 patients. Patients were divided into two groups: with and without gastrointestinal bleeding. Patient's information was extracted using their hospital records and the data eventually was statistically analyzed.

Results: The results of this study showed no significant difference between the two groups in terms of age, gender, marital status, distribution of weight, and education level ($P>0.05$). The frequency of NSAIDs use was significantly higher in patients with gastrointestinal bleeding ($P = 0.016$) with the most NSAID use as aspirin (32.66%). The prevalence of smoking, using drugs and alcohol consumption was significantly higher in the study group ($P < 0.05$).

Conclusion: A history of consuming NSAIDs increases the risk of GI bleeding. The frequency of cigarette, drug, and alcohol consumption in the case study group was significantly higher than that of the control group.

Keywords: Gastrointestinal bleeding, Nonsteroidal anti-inflammatory drugs, aspirin

Introduction:

Gastrointestinal (GI) bleeding is the most common reason for acute hospitalization of patients in gastroenterology wards(1). Different studies have evaluated the implemented cost of GI bleeding on both patients and the health care system, in addition to its impact on mortality and morbidity rates. These studies have concluded that GI bleeding, in fact,

31 incurs higher patient and system costs and raises mortality and morbidity
32 rates(2, 3).Several factors have been propounded as etiologic factors
33 behind GI bleeding, among which H.pylori, Non-Steroid Anti-
34 Inflammatory Drugs (NSAIDs), and aspirin have been deemed as the
35 most significant, especially in upper GI bleeding(4). Different
36 epidemiologic studies have suggested that a combination of several
37 different NSAIDs, or a high dose of any one of these drugs, can increase
38 the risk of GI bleeding up to seven and nine fold respectively(5). These
39 results emphasize not only the importance of NSAIDs but also their
40 sensible usage. Recently, due to the increased prevalence of arthritic
41 diseases and osteoarthritis, the use of NSAIDs has grown. The
42 prescription of multiple NSAIDs to patients by different physicians in
43 various fields has led to the increased simultaneous consumption of
44 several NSAIDs. This raises the risk of GI bleeding and other NSAID
45 side effects, especially among the older population.

46 In a study on Northeast of Iran Zeinali et al showed that about 20%
47 of all prescriptions included at least one NSAIDs. in comparison with
48 12.1% in USA (6). In consideration of the rising usage of NSAIDs, it is
49 imperative to study the relationship between the use of NSAIDs and
50 upper GI bleeding. Although this relationship is now mentioned in
51 textbooks, there has not yet been any related study conducted in

52 Mashhad, Iran to explore the high prevalence of NSAID consumption,
53 this despite numerous warnings about the side effects following the
54 unbridled use of these drugs. Furthermore, since accurate statistics
55 about NSAID side effects are critical for future prophylaxis
56 recommendations, it appears beneficial to conduct a study on the
57 correlation between upper GI bleeding and NSAID consumption (6).

58 **Method and Materials**

59 The present case control study was conducted in several steps. These
60 steps were performed simultaneously and by one researcher. The steps
61 were designed as a checklist for utilization in the present study, by which
62 samples were chosen, data extracted and collected, and statistical
63 calculations made.

64 Checklist Design:

65 The checklist was designed as two forms. The first form was intended for
66 patients hospitalized at the emergency ward of Qaem Hospital,
67 Mashhad, Iran due to upper GI bleeding and who had undergone
68 diagnostic and therapeutic measures. This checklist included
69 identification code, gender, age, weight, educational level, occupation,
70 marital status, and residence. Also listed was any history of digestive
71 diseases, GI bleeding, non-digestive diseases, smoking, alcohol
72 consumption, drug abuse, and medications. In addition, the following

73 information was provided: endoscopy results, primary hemoglobin,
74 primary platelet, primary PT, INR, and the possible need for a blood
75 transfusion.

76 The second form was designed for patients hospitalized at Khatam-
77 al_Anbia Hospital, Mashhad, Iran of Ophthalmology with a chief
78 complaint and reason for hospitalization that was unrelated to GI
79 bleeding (control group). This checklist included the following:
80 identification code, gender, age, weight, educational level, occupation,
81 marital status, and residence. Also listed was any history of digestive
82 diseases, GI bleeding, non-digestive diseases, smoking, alcohol
83 consumption, drug abuse. All the patients were asked for NSAID use
84 (continuously or based on need).

85 Selection of Cases and Control Samples:

86 The method of sampling in the current study was simple nonrandomized.
87 Two groups were studied. The first group (case group) consisted of
88 patients hospitalized for gastrointestinal bleeding according to clinical
89 manifestation and/or diagnostic endoscopic examination at the
90 emergency ward of Qaem Hospital. The second group (control group)
91 was made up of patients hospitalized at Khatam-al_Anbia Hospital of
92 Ophthalmology due to a chief complaint and reason for hospitalization
93 unrelated to gastrointestinal bleeding.

94 Data Extraction and Collection:

95 In this step of the present study, required data were collected and
96 registered onto the appropriate checklist. The case study group data
97 were collected from the patients' hospital files. The control group data
98 was obtained by conducting direct interviews and also by accessing the
99 patients' previous files. To accomplish this, at Khatam-al Anbia Hospital,
100 the researcher first explained the study and its purpose to the patients
101 and obtained their consent before reviewing files or interviewing.

102 Statistical Calculations:

103 First, the data **was entered** into SPSS ver.16 software. The **median** and
104 Interquartile range were utilized to describe the quantitative data
105 **indexes**. Frequency and frequency percentage served as the indexes for
106 explaining qualitative data. In order to compare qualitative variables in
107 the case study and control groups, the Chi-squared test or **where**
108 **appropriate**, exact fisher test were utilized. If the data had a normal
109 distribution, the independent t-test compared the quantitative data from
110 the two groups; otherwise, its nonparametric equivalent (Mann-Whitney)
111 performed this comparison. The confidence interval and the level of
112 significance were deemed as 95% and **0.05%, respectively**.

113 **Results**

114 The present work studied a total number of 300 patients aged above 35.
 115 Patients were divided into two groups: 1)those suffering from upper GI
 116 bleeding (case study) and 2) those without GI bleeding (control group).
 117 The mean age of the patients was 45.81±21.28 years (in the range of 40
 118 to 87 years).

119 The patient demographics of the two groups are compared in Tables 1
 120 and 2. Statistical tests showed no significant difference between the two
 121 groups in terms of age, gender, marital status, distribution of weight, and
 122 education level (P>0.05). In regard to residence, the results indicated a
 123 significant difference between that of the case study and the control
 124 group(p=0.002).

125 **Table 1:** Comparison of median and Interquartile Range of Demographic
 126 variables

Variable Group		Case Study Group (with GI bleeding) (IQR) Median (150n=)	Control Group (without GI bleeding) Median (IQR)(150n=)	P-value
§Age		(65-47) 59	(65-45) 54	0.116
Gender	male	88 (58.6%)	84 (56%)	0.726
	female	62 (41.3%)	66 (44%)	
	< 50	25 (16.6%)	31 (20.7%)	

Weight	50 -70	81 (54%)	74 (49.3%)	0.615
(kg)	>70	44 (29.4%)	45 (30%)	

127 §: Mann-Whitney statistical test

128

129 **Table 2:** Comparison of Qualitative demographic Variable Frequency

Variable Group		Case Study Group Frequency & Frequency Percentage	Control Group Frequency & Frequency Percentage	*P-value
Marital Status	single	45 (28.7%)	40 (26.7%)	0.796
	married	105 (71.3%)	110 (73.3%)	
Educational Level	illiterate	12 (8%)	12 (8%)	0.107
	only reading & writing	36 (24%)	20 (13.3%)	
	up to elementary school	32 (21.3%)	36 (24%)	
	high school diploma	53 (35.3%)	51 (34%)	
	associate degree	13 (8.7%)	25 (16.7%)	


	Bachelor degree&higher	4 (2.7%)	6 (4%)	
Residence	city	94 (62.7%)	124 (82.7%)	0.002
	village	56 (37.3%)	26 (17.3%)	

130

131 *: Chi-Square statistical test

132 Figure-1 compares the two groups' frequency of cigarette and alcohol
 133 consumption and drug abuse.

134 **Figure 1:** Comparison of the Consumption of Cigarettes, Drugs, and
 135 Alcohol



136

137 **Table 3** provides the frequency of NSAID consumption in the two
 138 groups. As indicated, NSAID consumption in patients with GI bleeding
 139 was significantly higher compared to those not suffering from this

140 condition (p=0.022). Furthermore, the comparison among the types of
 141 NSAID consumed by patients of the two groups showed a significant
 142 difference in regard to type(p<0.001). The most commonly used NSAID
 143 in the case study group was aspirin 80 mg per day, either as an ongoing
 144 consumption or at least for a past period of time. In fact, except for six
 145 patients, all subjects in the control group using NSAIDs were taking
 146 aspirin. After aspirin, the most common NSAID in the study group was
 147 ibuprofen. However, in the control group, there was greater consumption
 148 of ibuprofen followed by aspirin and diclofenac, respectively. In both
 149 study groups, other NSAIDs, such as indomethacin and naproxen, were
 150 less commonly used.

151 **Table 3:** Comparison of the frequency of NSAID consumption and its
 152 subtypes
 153

		Control	Case	P-value
NSAID Consumption	yes	35 (23.4%)	55 (36%)	0.022**
	no	115 (76.6%)	95 (64%)	
Type of NSAID	Aspirin	13 (8.7%)	49 (32.66%)	*<0.001
	Ibuprofen	14 (9.3%)	5 (3.34%)	

	Diclofenac	7 (4.7%)	0 (0%)
	Indomethacin	0 (0%)	1 (0.7%)
	Naproxen	1 (0.7%)	0 (0%)
	others	0(0%)	0(0%)

154

155 *: Chi-Square statistical test

156 **: Fisher Exact statistical test


157

158 The frequency of other medications taken by patients is presented in
 159 figure 2. A significant difference in the type of drugs taken is evident
 160 between the two groups ($p=0.035$). In the case study group, the most
 161 common were corticosteroids (15.3%), warfarin (14%), and plavix
 162 (13.3%). However, in the control group, the most prevalent drugs were
 163 specific serotonin receptor inhibitors (SSRIs) (10.6%), plavix (4%), and
 164 corticosteroids (4%). In both groups, the consumption of heparin was
 165 less than any other of the drugs.

166

167
168


Figure 2: Comparison of the Frequency of Other Drugs Consumption



169
170
171

172 Figure 3 provides patient endoscopy results. As seen, out of 150
173 patients with GI bleeding, 97 had undergone an endoscopy while the
174 other 53 patients had not because of various reasons, such as patient
175 unwillingness or medical conditions. The most common pathologic
176 finding following endoscopy was ulcers (38.1%), while a mass was the
177 least commonly observed pathology (4.2%). In 20.6 % of the patients,
178 the endoscopy results were normal.

179 **Figure 3: Frequency of Endoscopic Findings in Patients with Upper GI**
180 **Bleeding**



181

182

185

184 The logistic regression test was employed to assess the extent of the
 185 studied variables' prediction effect on GI bleeding. As seen, in a
 186 comparison between **the study** and the control group, cigarette use,
 187 alcohol consumption, and, finally, a history of NSAID consumption can
 188 lead to an increased risk of upper GI bleeding incidence with an OR of
 189 1.81, 4.241, **and 1.838**, respectively.

190 From these variables, drug abuse, gender, and age did not have any
 191 effect on raising or lowering the risk of upper GI bleeding incidence.

192 **Table 4** lists the results for each studied variable.

193 **Table 4:** Evaluating the Predictive Effect of Variables under Study on
 194 the Establishment of GI Bleeding

195

Variable	CI 95% for OR (lower-upper)	Odds Ratio(OR)	P- value**
----------	--------------------------------	-------------------	---------------

Age	0.942-1.004	0.965	0.175
Gender (risk of males compared to females)	0.634-1.664	1.029	0.92
Cigarette	1.071-3.151	1.838	0.024
Drug abuse	0.941-5.459	2.72	0.06
Alcohol Use	1.415-13.29	4.241	0.02
NSAID Use	1.076-3.067	1.812	0.021

196

Logistic regression test**

197

Discussion

198

199 In the present study, statistical tests revealed that there was no
 200 significant difference between the two groups in regard to gender
 201 frequency, age, marital status, weight group frequency, and educational
 202 level. The study revealed that patients in the group lived more frequently
 203 in urban areas, a finding that had barely been investigated previously.
 204 For instance, the Button et al. study conducted in 2010 showed that a
 205 higher number of patients with upper GI bleeding lived in urban areas(7).
 206 Likewise, in the 2012 study by Whiskey et al., the prevalence of variceal
 207 and non-variceal upper GI bleeding was reported to be greater among
 208 the urban population(8). Yet, these two above-mentioned studies did not

209 further explore the possible reasons behind their findings. In any case, it
210 seems that the stronger presence of risk factors for upper GI bleeding in
211 urban areas has led to a greater number of patients in these areas.

212 The present research suggest that the prevalence of cigarette smoking,
213 alcohol consumption, and drug abuse was significantly higher in the
214 study group when compared to the control group. These findings have
215 also been reported in other similar studies. For example, the Crooks et
216 al. 2013 study found that cigarette use (whether active or passive) and
217 alcohol consumption increased the risk of upper GI bleeding. The study
218 also reported that the risk of bleeding incidence grew following a rise in
219 alcohol consumption (9). Another US study in 2016 revealed that
220 drinking more than 30 gr of alcohol per day or more than 5 times per
221 week was deemed to be an independent risk factor increasing the
222 incidence of GI bleeding (10). The study also suggested that cigarette
223 use is not related to GI bleeding. Alcohol related mucosal damage can
224 be caused by a rise in the production of oxygen-free radical species, a
225 fall in the level of prostaglandins, and also the release of mucosal
226 leukotrienes (11, 12).

227 As for cigarette smoking and upper GI bleeding, different results have
228 been reported by various studies. This factor requires larger population
229 size for evaluation. Similar to the current work, some researchers have

230 propounded cigarettes as a risk factor for GI bleeding, while some others
231 have not(10, 13, 14).

232 In the present study's comparison of NSAID consumption between **the**
233 **study** and control group, there was a significant difference in the type of
234 NSAID used. Except for six patients, all of the study patients had used
235 NSAIDs,of which aspirin was the most commonly consumed followed by
236 ibuprofen. Patients in the control group, however, had comparatively
237 higher ibuprofen consumption, with aspirin being the second most
238 common drug consumed. **Aspirin was used at a dose of 80 mg per day**
239 **in both groups and unfortunately the dose of NSAIDs were not available**
240 **which was one limitation of the present study.** After aspirin, the control
241 group used **diclofenac** at a higher rate than that of the case study
242 patients. Both groups had a lower consumption of other types of
243 NSAIDs, such as indomethacin and naproxen. In conclusion, the present
244 study generally associates aspirin consumption with greater GI bleeding.
245 This finding has also been noted in several previous studies. For
246 example, review article by Castellsague et al. **at 2012** concluded that
247 ibuprofen, the most commonly used drug in the control group, is the
248 safest NSAID from the aspect of upper GI bleeding(15). Also, in their
249 2012 study, De Abajo et al. investigated the relation of NSAIDs and
250 other **drugs consumption** with upper GI bleeding. It was revealed that
251 aspirin poses a higher risk of upper GI disease than the consumption of

252 other drugs(16).These results are in accordance with the present work's
253 findings.

254 Another result of the present study addresses the frequency of other
255 drugs consumption by the patients of the two groups. In both the study
256 and the control group, a large spectrum of drugs were taken by patients,
257 out of which the current work attempted to discern which are closer
258 related to upper GI bleeding. The findings show a significant difference
259 in the amount of drugs consumed by the two groups. Among the study
260 subjects, the most common were corticosteroids (15.3%), warfarin
261 (14%), and clopidogrel(13.3%). However, the most prevalent
262 medications for control group subjects were SSRIs (10.6%), clopidogrel
263 (4%), and corticosteroids (4%). In general, the drug consumption in the
264 study group was significantly higher. In both groups, heparin was the
265 least used. Previous studies have also investigated the relation between
266 the use of various drugs and upper GI bleeding. For instance, the 2014
267 review by Narum et al. finally concluded that corticosteroids use is
268 associated with an increased risk of upper GI bleeding and gastric
269 ulcers(17). As mentioned in the present study, using these drugs in the
270 study group was more than in the control group. Regarding warfarin and
271 clopidogrel and their relation to upper GI bleeding, previously conducted
272 studies concur with the current paper's findings. In 2013, De Abajo et al.
273 concluded that Clopidogrel consumption can heighten the risk of upper

274 GI bleeding in comparison to healthy individuals(16). The subjects in the
275 present paper's **study** group also took more multiple NSAIDs than did
276 the control group. Previous studies have stressed that the consumption
277 of multiple NSAIDs **increases** the risk of upper GI bleeding. After
278 exploring the risk factor of upper GI bleeding in their 2010 research,
279 Scarpiganto and Hunt concluded that taking multiple NSAIDs or
280 anticoagulant drugs, such as warfarin and corticosteroids, all can
281 increase the risk of gastric bleedings, a finding with which the present
282 paper is in accordance(18).

283 The current paper's other results deal with patient endoscopies. **Ninety**
284 **seven** patients with GI bleeding underwent endoscopy while the other 53
285 patients did not for reasons such as medical issues or unwillingness to
286 consent to the procedure. The most common pathologic finding **was**
287 **ulcers. In the 2011Hearnshaw et al. study of 6,750 patients with upper**
288 **GI bleeding, the most commonly observed pathology was ulcers**
289 **(19),findings** similar to those of the current research.

290 **Conclusion**

291 The results of the present study indicate that greater consumption of
292 NSAIDs in patients with upper GI bleeding is significantly higher in
293 comparison with patients not suffering from this condition. Aspirin and
294 Ibuprofen were the most two common drugs used. Moreover,a history of
295 consuming NSAIDs increases a patient's risk of acquiring GI bleeding.

296 Concurrent consumption of corticosteroids, Warfarin and clopidogrel
297 increased the risk of GI bleeding in case group. Likewise, the frequency
298 of cigarette, drug, and alcohol consumption in the case study group was
299 significantly higher than that of the control group, thus signifying that
300 variables, such as alcohol and cigarettes, increase the risk of GI
301 bleeding in patients.

302 We suggest to consider the factors that increase the risk of upper GI
303 bleeding in patients with NSAID use and prescribe prophylaxis to high
304 risk patients.

305 Ethical Approval Disclaimer:

306 This research was approved at ethical committee of Mashhad University
307 of Medical Sciences with ethical code:922817

308

309 Consent Disclaimer:

310 As per international standard or university standard, patient's consent
311 has been collected and preserved by the authors.

312 References

- 313 1. Williams JG, Roberts SE, Ali MF, Cheung WY, Cohen DR, Demery G, et al. Gastroenterology
314 services in the UK. The burden of disease, and the organisation and delivery of services for
315 gastrointestinal and liver disorders: a review of the evidence. *Gut*. 2007;56 Suppl 1:1-113.
- 316 2. Peery AF, Crockett SD, Barritt AS, Dellon ES, Eluri S, Gangarosa LM, et al. Burden of
317 Gastrointestinal, Liver, and Pancreatic Diseases in the United States. *Gastroenterology*.
318 2015;149(7):1731-41.e3.
- 319 3. Abougergi MS, Travis AC, Saltzman JR. The in-hospital mortality rate for upper GI
320 hemorrhage has decreased over 2 decades in the United States: a nationwide analysis.
321 *Gastrointestinal endoscopy*. 2015;81(4):882-8.e1.

- 322 4. Yeomans ND. The ulcer sleuths: The search for the cause of peptic ulcers. *Journal of*
323 *gastroenterology and hepatology*. 2011;26 Suppl 1:35-41.
- 324 5. Grosser T, Smyth E, FitzGerald G. Anti-inflammatory, Antipyretic, and Analgesic Agents. In:
325 Brunton L, Chabner B, Knollmann B, editors. *Goodman & Gilman's The Pharmacological Basis of*
326 *Therapeutics*. 12th ed. New York: McGraw-Hill 2011. p. 959-1005.
- 327 6. Zeinali M, Tabeshpour J, Maziar SV, Taherzadeh Z, Zirak MR, Sent D, et al. Prescription
328 pattern analysis of nonsteroidal anti-inflammatory drugs in the northeastern iranian population.
329 *Journal of Research in Pharmacy Practice*. 2017;6(4):206.
- 330 7. Button LA, Roberts SE, Evans PA, Goldacre MJ, Akbari A, Dsilva R, et al. Hospitalized
331 incidence and case fatality for upper gastrointestinal bleeding from 1999 to 2007: a record linkage
332 study. *Alimentary pharmacology & therapeutics*. 2011;33(1):64-76.
- 333 8. Wysocki JD, Srivastav S, Winstead NS. A nationwide analysis of risk factors for mortality and
334 time to endoscopy in upper gastrointestinal haemorrhage. *Alimentary pharmacology & therapeutics*.
335 2012;36(1):30-6.
- 336 9. Crooks CJ, West J, Card TR. Comorbidities affect risk of nonvariceal upper gastrointestinal
337 bleeding. *Gastroenterology*. 2013;144(7):1384-93, 93.e1-2; quiz e18-9.
- 338 10. Strate LL, Singh P, Boylan MR, Piawah S, Cao Y, Chan AT. A Prospective Study of Alcohol
339 Consumption and Smoking and the Risk of Major Gastrointestinal Bleeding in Men. *PloS one*.
340 2016;11(11):e0165278.
- 341 11. Tarnawski A, Hollander D, Stachura J, Klimczyk B, Mach T, Bogdal J. Alcohol injury to the
342 normal human gastric mucosa: endoscopic, histologic and functional assessment. *Clinical and*
343 *investigative medicine Medecine clinique et experimentale*. 1987;10(3):259-63.
- 344 12. Oates PJ, Hakkinen JP. Studies on the mechanism of ethanol-induced gastric damage in rats.
345 *Gastroenterology*. 1988;94(1):10-21.
- 346 13. Udd M, Miettinen P, Palmu A, Heikkinen M, Janatuinen E, Pasanen P, et al. Analysis of the
347 risk factors and their combinations in acute gastroduodenal ulcer bleeding: a case-control study.
348 *Scandinavian journal of gastroenterology*. 2007;42(12):1395-403.
- 349 14. Yamada A, Sugimoto T, Kondo S, Ohta M, Watabe H, Maeda S, et al. Assessment of the risk
350 factors for colonic diverticular hemorrhage. *Diseases of the colon and rectum*. 2008;51(1):116-20.
- 351 15. Castellsague J, Riera-Guardia N, Calingaert B, Varas-Lorenzo C, Fourrier-Reglat A, Nicotra F,
352 et al. Individual NSAIDs and upper gastrointestinal complications: a systematic review and meta-
353 analysis of observational studies (the SOS project). *Drug safety*. 2012;35(12):1127-46.
- 354 16. de Abajo FJ, Gil MJ, Bryant V, Timoner J, Oliva B, Garcia-Rodriguez LA. Upper gastrointestinal
355 bleeding associated with NSAIDs, other drugs and interactions: a nested case-control study in a new
356 general practice database. *European journal of clinical pharmacology*. 2013;69(3):691-701.
- 357 17. Narum S, Westergren T, Klemp M. Corticosteroids and risk of gastrointestinal bleeding: a
358 systematic review and meta-analysis. *BMJ open*. 2014;4(5):e004587.
- 359 18. Scarpignato C, Hunt RH. Nonsteroidal antiinflammatory drug-related injury to the
360 gastrointestinal tract: clinical picture, pathogenesis, and prevention. *Gastroenterology clinics of*
361 *North America*. 2010;39(3):433-64.
- 362 19. Hearnshaw SA, Logan RF, Lowe D, Travis SP, Murphy MF, Palmer KR. Acute upper
363 gastrointestinal bleeding in the UK: patient characteristics, diagnoses and outcomes in the 2007 UK
364 audit. *Gut*. 2011;60(10):1327-35.