

1 **Modern fungicides: mechanisms of action, fungal resistance and phytotoxic effects**

2
3
4 **ABSTRACT**

5
6 The establishment of safe and effective methods for controlling fungal diseases is an urgent issue
7 in agriculture and forestry. Fungicide research has provided a wide range of products with new
8 modes of action. Extensive use of these compounds in agriculture enhances public anxiety due to
9 the harmful potential for the environment and human health. Moreover, the phytotoxic effects of
10 some fungicides are already recognized but still little is known about their influence on the
11 photosynthetic apparatus and plant physiology. This review provides an understanding of the
12 mechanisms of action of fungicides, mechanisms of fungicide resistance development, and the
13 phenomenon of phytotoxicity.

14 **KEYWORDS** – contact fungicides, systemic fungicides, disease control, resistance,
15 phytotoxicity
16

17 **1. INTRODUCTION**

18 Fungicides are chemical substances used for control and treatment of fungal diseases of
19 plants. The employment of fungicides has become widespread in recent decades in agriculture
20 since it was estimated that fungal infections reduce yields of the crops worldwide by nearly 20%
21 (Rohr *et al.*, Brown, Battaglin, McMahon, & Relyea, 2017). Fungicides have become the primary
22 means of fungal disease control due to their relatively low cost, ease of use and efficiency (Xia *et*
23 *al.*, 2006).

24 Disease management is an essential component of production for all crops, often having a
25 significant economic impact on their yield and quality. There are three main reasons for using
26 fungicides:

- 27 – To control the infection during the establishment and growth of a grain crop;
 - 28 – To enhance the productivity of cereal and to decrease defects.
- 29 Infection may result in a decrease in productivity due to the damage to photosynthetic parts.
30 Defects in the edible parts of the crop or leaves of ornamentals affect their attractiveness, and
31 consequently the market prices;
- 32 – To improve the shelf life and quality of produced and harvested plants.

33 Some of the significant disease damage occurs post-harvest. Harmful fungi often worsen
34 stocks of grain crops, vegetables, and tubers. Several grain-infecting species of *Fusarium*,
35 *Penicillium* or *Aspergillus* produce important mycotoxins which can cause serious illness or even
36 death in humans and animals after eating contaminated food (Marin, Ramos, Cano-Sanche, &
37 Sanehis, 2012). Fungicides have been used to decrease mycotoxin contamination of wheat
38 affected by *Fusarium* head blight, but most fungicides developed so far have not been entirely
39 adequate for the regulation of mycotoxin production associated with other diseases (Forrer *et al.*,
40 2014). This is due to insufficient knowledge of the protectants mechanisms action and the
41 response of the plant.

42 The appearance of new strains of fungal pathogens and their resistance to the available
43 commercial products is often associated with extensive use of these compounds (Pablo *et al.*,
44 Garcia, Rosa-M. Rivero, Juan-M. Ruiz, 2003). What is more, the widespread and frequent use of
45 fungicides in plant protection generates a long-term accumulation of residues in food and the
46 environment (*Report on the pesticide residues monitoring programme: Quarter 1 2017*, 2017), (Anne-
47 Nolle Petit *et al.*, Fontaine, Ement, & Vaillant-Gaveau, n.d.). In the Report on the pesticide residues
48 monitoring programme in 2017, analyzing vegetables and fruits from 27 countries for contamination
49 with pesticides has shown that dithiocarbamates are among the most common residual
50 contaminants. Accordingly, the excessive use of such compounds in agriculture gave rise to public
51 concerns because of the detrimental effects on the environment and risk for human health (*Report on*
52 *the pesticide residues monitoring programme: Quarter 1 2017*, 2017).

53 For example, the fungicide chlorothalonil - the most common synthetic fungicide in the
54 United States - was shown to be toxic to aquatic animals such as tadpoles, oysters, or fish (Vincelli
55 P., 2002).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

56 In some cases, fungicides derived from "natural" sources are much safer than synthetic. The
57 primary sources include copper, sulphur, plant oils and bicarbonates. But even copper can be skin
58 irritating, eyes and the respiratory and digestive tracts, while sulphur can result in dermatitis and
59 diarrhea (Southern-AG, 2015). To use any fungicide safely and efficiently, one needs to correctly
60 diagnose the problem and choose the best treatment strategy.

61 2. CLASSIFICATION OF FUNGICIDES

62 Fungicides are often classified as protective or system. Protective fungicides are usually
63 effective against a range spectrum of fungi and protect the plant from infection on leaf surface and
64 stems. They often require repeated application during the growing season to provide coverage as
65 new plants appear. Systemic fungicides can be absorbed by the plant without damage and be
66 transported to other tissues where they are toxic to fungi. These compounds can control and fight
67 infections, but they are also vulnerable to resistance to fungi, as they usually target only one step,
68 to kill the fungus. To reduce resistance due to excessive use of chemicals, the fungicides are
69 classified according to their chemical class. By alternating between different classes of fungicides
70 the fungal population is less likely to develop resistance to a particular chemical. (Add references)

71 Chemically, organic molecules always contain carbon atoms in their structure while most
72 inorganic molecules do not. Initially, first fungicides were inorganic compounds based on sulphur or
73 metal ions (copper, tin, cadmium, mercury) that are known to be toxic to fungi. Currently, fungicides
74 based on copper and sulphur are still widely used. Copper sulphate has been registered for use in
75 the United States since 1956. The copper atom binds to proteins, changing their structure. This
76 may break the membranes around the cells, causing the cells to die. Thus, copper sulfate is
77 effective in the destruction of fungi, algae and even snails. However, most fungicides used today
78 are organic synthetic compounds (Lesemann, *et al.*, Schimpke, Dunemann, & Deising, 2006).

Formatted: Font: Italic

79 2.1. Non-systemic (contact) fungicides

80 This type of fungicides has a preventive impact by killing or inhibiting fungi and fungal spores
81 before the mycelia can grow and develop within the plant tissues (Oliver & Hewitt, 2014), but have
82 little or no effect once the fungus has entered or colonized host tissue. Additionally, while non-
83 systemic fungicides generally remain on the surface of plants, they are potentially phytotoxic and
84 can damage the plant when absorbed (Lesemann, *et al.*, 2006). Contact action has derivatives
85 dithiocarbamates acid, agents based on sulphur, copper, etc. Thus, this kind of fungicides can be
86 used only as protectants. It is therefore also important to apply them on given plants before known
87 infection period begins to decrease the chance of infection. Contact agents – such as zineb,
88 polycarbonate, copper oxychloride, sulfur, mancozeb, bordeaux liquid and others are not able to cure
89 already diseased plants. Despite their potential harm to plants, non-systemic pesticides are thought
90 to be okay as they can be removed or flushed from the plant before harvest. This makes the
91 produce clean from pesticide chemical tainting and thus better for human consumption.

Formatted: Font: Italic

92 Typical examples of the primary contact fungicides are inorganic copper compounds such as
93 Bordeaux mixture, copper carbonate, and inorganic sulphur in the form of elemental sulphur and
94 lime sulphur (Pablo G. García, Rosa M. Rivero, Juan M. Ruiz, *et al.*, 2003). The organic contact
95 fungicides (e.g., thiram, ferbam, and ziram) play an important role in the comprehensive control of
96 plant diseases since they are more efficient and less toxic than the inorganic compounds (Aynalem
97 & Assefa, 2017), (Nason, Farrar, & Bartlett, *et al.*, 2007).

Formatted: Font: Italic

98 Contact fungicides are products suited for preventive (prophylactic) use as they work by contact
99 action on the surface of the plant. Therefore, to protect new plant growth and renewal of the material
100 washed off by rain or irrigation, or degraded by such environmental factors as wind and the amount of
101 UV, repeated applications are necessary. The protective action of these fungicides does not exceed 10-
102 12 days before the first heavy rain, after which the treatment is repeated. The number of treatments with
103 a fungicide of contact action is 3 to 6 treatments per season. During processing, it is necessary to spray
104 not only the surface of the leaves but the underside too, since many types of fungi begin to grow from
105 the underside of the leaves. For example, for processing potatoes the rate of application may be every
106 7 days during the month (Johnson, Hamm, & Sunseri, n.d.????).

Formatted: Font: Italic

107 Contact fungicides do not penetrate deeply in the plant tissue and are easily removed,
108 leaving a clean product for consumption. They are effective with timely treatment and following
109 instructions. Because of this, and due to relatively low prices (but it should be remembered that
110 their consumption is much higher than systemic fungicides), they are still extensively used for
111 plant protection even though new, more potent fungicides are developed.

112 2.2. Systemic Fungicides

113 Systemic fungicides are absorbed by the plant and transported to the site of infection. These
114 compounds can, therefore, kill the fungus after the mycelia have penetrated the parenchyma of the
115 plant tissue, stopping the spread of infection (Oliver & Hewitt, 2014). Some systemic fungicides move
116 within the plant only a short distance from the site of penetration. This is local-systemic fungicides.
117 The dicarboximide fungicides are one example of this group (González M., and Caetano P., 2017).
118 The dicarboximide fungicides, iprodione, procymidone, vinclozolin, chlozolinate, and metomeclan
119 are especially promising for the control of plant diseases caused by species of *Botrytis*, *Sclerotinia*,
120 *Monilinia*, *Alternaria*, *Sclerotium*, and *Phoma* [56????]. The mode of action of these compounds is
121 apparently related to the inhibition of triglyceride biosynthesis in fungi [17????].

122 Some locally systemic fungicides cross the leaf plate from one leaf surface to the other but do
123 not spread inside the plant. Those fungicides are called translaminar, i.e. trifloxystrobin(reference?).
124 Systemic fungicides, which are called xylem-mobile or acropetal systemics, move inside the water-
125 conducting tissue (xylem), which raises them up in the transpiration flow, however, mobility within the
126 plant is limited. For example, DMI fungicides are moderately mobile within plants. Others are very
127 mobile and easily move around the xylem. The examples of systemic fungicides which are mobile in
128 xylem are thiophanate-methyl and mefanox (Paul Vincelli and, Bruce Clarke, 2017). The third type of
129 systemic fungicide is a phloem-mobile system, compound circulates in phloem out of the sheet
130 where deposited upwards to the other leaves and downwards to the roots (Lesemann, et al., 2006).
131 Only one example of this type of systemic exists among turfgrass fungicides: the phosphonates,
132 which include fosetyl-Al and the phosphites (reference?).

133 Systemic fungicides can be used as protectants, eradicants, or both, and are the most
134 recently developed and the most promising type of fungicides at the moment (Pablo et al., G.
135 Garcia, Rosa M. Rivero, Juan M. Ruiz, 2003). Though systemic fungicides usually have a particular
136 location of action, fungi may quickly develop resistance to them if they are managed inappropriately
137 (reference).

138 Highly specific modern fungicides block only one target in the pathogen (monospecific
139 fungicides or single-site inhibitors). Deising et al. (2008) state that "examples of single-site
140 inhibitors are the benzimidazoles, phenylamides and strobilurins, released to the market in the late
141 1970es and the mid 1990es" (Miguez et al., Reeve, Wood, & Hollomon, 2004).

142 Extensively used in agriculture are also benzimidazoles, a group of organic fungicides with
143 systemic action. These types of compounds control a wide range of fungi at a comparatively low cost
144 of treatment (Bernauer, Gaines-Day, & Steffan, et al., 2015). For example, benomyl is one of the most
145 effective and extensively used benzimidazoles in crop protection (Pablo C. Garcia, Rosa M. Rivero et
146 al., Juan M. Ruiz, 2003). The benzimidazoles benomyl, carbendazim, and thiabendazole and the
147 phenylcarbamate diethofencarb specifically interfere with the formation of microtubules, which
148 function in a variety of cellular processes, including mitosis and maintenance of the cell shape
149 (Saladin Gaëlle, Magné Christian, & Clément, et al., 2003); (Eislahi, et al., Osman, Sherif, &
150 Elhussein, 2014). These fungicides bind specifically to protein subunits called tubulin and prevent
151 their assembly from forming microtubules.

152 The main difference between the effects of systemic and contact fungicides is that the first one
153 sometimes suppresses the fungus after infection of the plant, whereas the second one must be
154 present on the plants surface before infecting. Gradually, since the 1960s, systemic fungicides
155 replaced non-systemic non-systemic preparation, providing higher levels of plant protection (Dias
156 Maria Celeste, 2012). However, compared with the non-systemics, systemic fungicides are roughly
157 twice as expensive regarding sales (McGrath, 2004).

158 3. BREADTH OF ACTIVITY

159 Depending on the scope of their targets, fungicides can be classified as single-site or multi-
160 site. Single-site fungicides active against one point in one metabolic pathway of the fungus (D.
161 Mueller, n.d.) (reference). Examples of such fungicides can be various different drugs with one
162 active ingredient, such as prothioconazole, pyraclostrobin, fludioxonil, the benzimidazoles
163 (benomyl, thiophanatemethyl) and others. However, there are connections that are not very
164 desirable to use alone (reference). For example, azoxystrobin is recommended to use as a mixture
165 with other fungicides having a different mechanism of action (reference). The probability of the
166 pathogen's development resistance, in this case, is significantly reduced because resistant isolates
167 to one fungicide will be killed by another fungicide. The effectiveness of this method can be
168 demonstrated by Metalaxyl, phenylamide fungicide. When used as the sole compound in Ireland to
169 combat pollution in potatoes (*Phytophthora infestans*) resistance developed within one growing
170 season. However, in countries such as the UK where it was sold only as a mixture, resistance
171 problems developed more slowly (reference).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

172 On the other hand, because of this specific activity, fungi are more likely to develop
173 resistance to the fungicide (Lesemann *et al.*, 2006).

Formatted: Font: Italic

174 Multi-site fungicides can target multiple locations (different metabolic pathways). But single-
175 site fungicides are considered less toxic to plants. Older contact fungicides such as mancozeb,
176 fluazinam etc have multi-site activity and affect many fungal species in different classes
177 (*Sclerotinia*, *Botrytis*, *Alternaria*, *Phytophthora*, *Peronospora*) (reference). Due to the rise in the
178 stringency and number of normative tests required to register a new active ingredient, fungicide
179 manufacturers have found it easier to develop single-site systemics recently (reference).
180 Consequently, fungicide resistance has become a more critical issue in disease regulation.
181 Examples of narrow-spectrum fungicides can be Folplan and Karatan (reference).

182 The active ingredient of Folplan — folpet derived phthalimide. Folplan, has a narrow
183 spectrum of activity, suppresses the development of pathogens peronospora and other fungi,
184 except for muchnationalmuch national (reference). To broaden the spectrum of action can be
185 mixed with other systemic fungicides, insecticides, which have no alkaline reaction (reference).
186 Folplan registered and approved for use on potatoes and grapes. Suppresses the development of
187 *Phytophthora*, *Peronospora*, *Oidium*, *Botrytis*. The flow rate - about 3.0 kg/ha. Maximum number of
188 treatments – two for season (reference).

189 The active substance of Karatan – dinocap derived nitrophenol. It suppresses the
190 development of powdery mildew pathogens and has acaricidal action. Ineffective against
191 peronosporic fungi. Can be mixed with other fungicides and insectoacaricides, which have no
192 alkaline reaction. The duration of the protective effect in the optimal concentrations of 10-15 days.
193 It is advisable to use prophylactic. The fungicide does not penetrate the leaves and fruit, so it's
194 easy to rinse them. Karatan is registered and approved for use on cucumbers the closed and open
195 soil, grapes, Apple, pear. The flow rate of the drug is 0.5-2.0 l/ha. The maximum number of
196 treatments – three for season (Add references).

197 4. APPLICATION METHODS

198 Fungicides can be produced in the form of dust, granules, gas, but most often fluid.
199 Depending on the type there are different methods of application:

200 1. Treat of planting material (mordanting). Fungicides can be applied in various solutions or
201 incrustation of seeds, dry method or humidification, encapsulating or pelleting.

202 2. Application to the soil. This process is suitable when dealing with soil-borne pathogens.
203 Most of these fungicides have low selectivity and thus eliminate not only bacteria and fungi but also
204 the larvae of insect pests which could be of concern for environmental protection.

205 3. Spraying. The manual sprayers are used, as well as a specialized automobile or aircraft
206 vehicles. Spraying can be carried out repeatedly in the rate of appearance of the young vegetative
207 organs of the plant, the duration (Woodward *et al.*, Russell, Baring, Cason, & Baughman, 2015) of
208 action of a fungicide, and the risk of re-infection (E. Lee Butler, 2006).

Formatted: Font: Italic

209 Great importance in the success of seed protection is the correct timing of fungicide
210 treatment. Thus, seed disinfectants are commonly used in packing material deposited in the late
211 summer or autumn, and fungicides are used for spraying perennial plants during dormancy in late
212 fall, winter or early spring, as they can be dangerous to growing plants (Hasan *et al.*; Ahmed,
213 Tofazzal, Mian, & Haque, 2013); (Shuping & Eloff, 2017). Currently, in addition to the use of the
214 described methods to prevent spoilage during storage, fruit treatment by fungicides is also
215 practiced (Clayton *et al.*, A. Hollier, Jeffrey W. Hoy, Christopher A. Clark, Charles Overstreet,
216 Jaspreet Sidhu, Melanie L. Lewis Ivey, Raghuwinder Singh, Trey Price III, Mary Helen Ferguson,
217 G. Boyd Padgett, 2016).

Formatted: Font: Italic

Formatted: Font: Italic

218 5. ROLE OF FUNGICIDES IN DISEASE MANAGEMENT

219 Forecasting systems are developed for many diseases based on an understanding of the
220 environmental conditions favourable for pathogen development. Typically, these are based on
221 temperature and relative humidity or leaf wetness in the area with a growing crop (reference).
222 Threshold-based fungicide programs involve routinely scouting the crop for symptoms, then
223 applying fungicides when the number of signs reaches a critical level beyond which the disease
224 cannot be controlled adequately (reference). In general, the most crucial aspect of developing and
225 using forecasting systems is the knowledge of the disease cycle of the pathogen. The disease
226 cycle determines whether the disease is monocyclic (one generation per year) or polycyclic
227 (multiple generations) and latent period (time between infection and symptom expression) is also
228 essential aspect [58????].

229 There are examples of an artificial neural network (ANN) capable of predicting diseases
 230 based on existing data. They perform extraordinarily complex calculations imitating biological in the
 231 real world without about course to exact quantitative. Back-propagation neural network (BPNN) is
 232 the most important and widely used one (reference). The RBF network is used in Ming-wang Shi
 233 research, which is one of the new effective neural networks and is realized through a linear
 234 combination of nonlinear primary functions from the space RN into a spatial RM through nonlinear
 235 transformation (reference). He applied the GM Model (1,1) to predict plant diseases collected
 236 during the simulations. The results of the experiments show that the coincidence of the GM model
 237 parameter (1,1) coincides with the standard deviation of the disease index and incidence. This
 238 indicates that the GM system (1,1) is effective for the analysis of morbidity, and the parameters GM
 239 (1,1) may well reflect the change in the incidence of plants (Ming-wang Shi, 2011).

240 Another interesting example of plant diseases prediction is the using of electric fields (Benelli
 241 Jesse J, 2013); (Kuna-Broniowski et al., Makarski, & Kuna-Broniowska, 2015). In the work of
 242 Marek Kuna-Broniowski and etc., this method is used to predict the spread of plant diseases from
 243 the *Septoria* by determining the splashing of raindrops. Most existing methods use climate
 244 conditions, calendar measurements, and disease cycles to predict infections (Donatelli et al.,
 245 2017). However, it is important to take into account the spraying of rain droplets as a method of
 246 transporting spores to higher parts of plants and neighbouring plants. Measurements of the
 247 scattering range and the number of spray particles using an electric field are achieved using a
 248 measuring system that allows accurate and reliable measurement of the dispersion range of
 249 sprayed droplets (reference).

250 Economic factors often influence the choice of fungicide and application timing. The most
 251 expensive fungicides and numerous applications are used on valuable plantings that might suffer a
 252 significant economic loss in the absence of treatment, for example, fruit trees (reference). The crop
 253 tolerance level, or detriment threshold, can change depending upon the stage of the crop
 254 development when attacked, crop management practices, climatic and location conditions
 255 (reference).

256 It is important to use the correct type of fungicide at the right time of year because one of the
 257 fungicide side-effects is phytotoxicity, i.e. a toxic effect on (beneficial) plants. For example,
 258 trifloxystrobin, which is often applied to *Vitis vinifera* vines, can damage and even kill some trees of
 259 the genus *Malus*. However, trifloxystrobin is dangerous for particular grape cultivars but not others
 260 (can cause injury to *Vitis labrusca*) (Vincelli P., 2002). Some fungicides are even more specific,
 261 such as triazole + Qols that cannot be applied to glycine max later than during a growth stage
 262 known as R5 (reference).

263 6. THE MAIN CLASSES OF FUNGICIDE AND PLANT PHYSIOLOGICAL RESPONSES

264 There are five main chemical classes of fungicides (Table 1). The largest group of them is
 265 triazoles. Fungicides of this class have been using against pathogens of various diseases of fruit
 266 and vegetable crops. Substances differ in the degree of activity, the spectrum of effects on
 267 pathogens, the rate of consumption, the grade of risk to ecosystems, the population and working
 268 personnel, the payback of the costs of their use. Despite the wide range of action, triazoles have
 269 disadvantages. The systematic use of preparation based on triazoles leads to the emergence of
 270 resistant fungal strains. For example, triadimefon does not completely inhibit the fungal germination
 271 of the genus *Puccinia*.

272 The widely accepted assumption that fungicide has low phytotoxicity has started to be
 273 outdated with the publication of more detailed analyses at the cell level that demonstrated several
 274 damages to the photosynthetic apparatus (Anne-Nolle Petit et al., n.d., ???); (Saladin Gaëlle et
 275 al., 2003).

276 Table 1 – The major classes of fungicides and their effects

Chemical class	Fungicides	Mechanism of action	Fungi	Resistance	Phytotoxicity	References
Triazoles	tebuconazole, prothioconazole, diphenocnazole, ciproconazole, propiconazole,	Inhibit sterol biosynthesis	<i>Botrytis</i> , <i>Ustilago</i> , <i>Cercospora</i> , <i>Tilletia</i> , <i>Zymoseptoria</i> , <i>Fusarium</i> , <i>Cochliobolus</i> , <i>Erysiphe</i> , <i>Altemaria</i> ,	The systematic use of drugs based on triazoles causes resistance. The triadimefon does not completely inhibit the germination of conidia and rust <u>urediospores</u> .	there is a violation of the synthesis of gibberellins (retardant	(Cools, Hawkins, & Fraaije, 2013), (Dias Maria Celeste, 2012),

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

	epoxiconazole, flutriafol, triadimefon, triticonazole, diniconazole		<i>Puccinia</i> , <i>Septoria</i> , <i>Pythium</i> , <i>Drechslera</i> , <i>Pyrenophora</i> , <i>Rhynchosporium</i> , <i>Cladosporium</i> , <i>Epicoccum</i> , <i>Phoma</i>		effect), the synthesis of sterols, a decrease in transpiration of plants	(D. S. Mueller, 2006), (Ahmad & Khan, 2012), (Costa et al., 2017)
Phenylpyrroles	fluodioxonyl	Inhibit micellar growth, reduce glucose phosphorylation during cell respiration, disrupt the function of cell membranes	<i>Tilletia</i> , <i>Fusarium</i> , <i>Ascochyta</i> , <i>Alternaria</i> , <i>Fusarium</i> , <i>Aspergillus</i> , <i>Rhizoctonia</i> , <i>Helminthosporium</i> ,	Low risk of resistance due to the mechanism of action	decrease CO ₂ assimilation, transpiration, stomatal conductance and intercellular CO ₂ concentration	(Anne-Nolle Petit et al., n.d.), (Saladin Gaëlle et al., 2003), (Kilani & Fillingier, 2016), (Lew, 2010), (Ren, Shao, Han, Zhou, & Chen, 2016)
Strobilurins	picoxystrobin, fluoxastrobin, azoxystrobin, trifloxystrobin, pyraclostrobin, krezoxim-methyl	Inhibit mitochondrial respiration by blocking electron transport in the cytochrome b and c ₁ chain	<i>Puccinia</i> , <i>Septoria</i> , <i>Pyrenophora</i> , <i>Alternaria</i> , <i>Cladosporium</i> , <i>Epicoccum</i> , <i>Botrytis</i> , <i>Rhynchosporium</i> , <i>Drechslera</i> , <i>Fusarium</i> , <i>Rhizoctonia</i> , <i>Ustilago</i> , <i>Erysiphe</i>	Field resistance was recorded in <i>Oidium erysiphoides</i> , <i>Erysiphe graminis</i> , <i>Botrytis cineria</i> . When strobilurins inhibit the activity of cytochrome b, alternative pathways of electron transport can easily be activated	in the plant are rapidly hydrolyzed by ether linkage. During periods of drought, damage is exacerbated	(Balba, 2007), (Reddy, 2012), (Vincelli P., 2002), (Wojdyła, 2007)
Benzimidazoles	prochloraz, thiabendazole, thiophanate-methyl, benomyl, carbendazim	Inhibit the synthesis of ergosterol in the fungal cell and disrupt its life activity	<i>Fusarium</i> , <i>Botrytis</i> , <i>Sclerotinia</i> , <i>Septoria</i> , <i>Uncinula</i> , <i>Erysiphe</i>	Stable pathogenic strains: <i>Pseudocercospora</i> , <i>Pseudocercospora</i> , <i>Septoria</i> , <i>Fusarium</i> , <i>Erysiphe</i> ,	decrease plant biomass. induces a considerable reduction on the chlorophyll a, chlorophyll b, caroteno	(Dias Maria Celeste, 2012), (Isaac, 1992), (Deising, Reimann, & Pascholati, 2008)

					ids, and the total pigments content	
Morpholines (cinnamic acid derivatives)	spiroxamine, dimethomorph	Prevent the formation of mycelium and block the reduction of the double compound C-C and ergosterol synthesis	<i>Erysiphe</i> , <i>Uncinula</i> , <i>Septoria</i> , <i>Puccinia</i>	Stable fungal strains form slowly, fungicides block the reduction reactions in the process of sterol biosynthesis and isomerization	decrease of the sterols synthesis	(Biol et al., 2013), (Isaac, 1992)

277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316

Triazoles also have phytotoxicity to protected plants. In a significant amount, fungicides cause a retardant effect (impaired synthesis of gibberellins); violate the synthesis of sterols, reduce transpiration of plants (Tom Allen, 2013). Triadimenol and propiconazole delay the removal of the primary leaf and violate its geotropism in the processing of cereal seeds. Tebuconazole can pass into the retardant under unfavourable conditions (waterlogging of the soil, lack of moisture, low germination energy, etc.). The same properties are inherent in triticonazole, to a lesser extent - to other azoles. But as the review "Constraints on the evolution of azole resistance in plant pathogenic fungi" says, today, the azoles still apply in the fight against pathogens of many culture, including grains, fruits and vegetables, canola and soybeans, despite numerous reports of azole-resistant fungal strains (Cools, Hawkins, & Fraaije *et al.*, 2013).

The next well-known group of fungicides (over 30 years old) is phenylpyrrole. They are chemical analogues of the natural antifungal compound pyrrolnitrin (Kilani & Fillinger, 2016). Currently, fungicoxon is used as the active substance of fungicides. Phenylsilyl inhibits all stages of fungal development, germination of spores, lengthening of the embryonic tubes and mycelium growth. The observed consequences are swollen hyphae with increased branching and apical lysis, which indicate that phenylpyrrols can act on the biosynthesis of the intragenic turgor and cell wall (Lew, 2010).

Recently strains resistant to fludioxonil have been isolated from *B. cinerea* populations in China at low levels (<3%). They represent typical osmosensitivity and developmental defects of fludioxonil resistant mutants (Ren, Shao, Han, Zhou, & Chen, *et al.*, 2016), which raises the question of their ability to compete with sensitive and severe strains and the selective pressure of fungicide treatments on these specific populations. Globally, there is no specific resistance to fludioxonil among gray mold populations that support the high efficacy of this fungicide (Walker *et al.*, 2013).

To avoid the emergence of resistance to phenylpyrroles, combined preparations should be used or alternate with different mechanisms of action. In addition to problems with possible resistance, there is a risk of phytotoxic effects in relation to protecting plants (reference). For example, in research of Petit A.N, Fontaine F, Clement and Vaillant-Gaveau N (Anne-Nolle Petit *et al.*, n.d.) and also Saladin G, Magne C, Clement C (Saladin Gaëlle *et al.*, 2003) about effects of fludioxonil in *Vitis vinifera L.* These reports have shown that application of fungicides has consequences for plant physiology, such as a plant growth reduction, perturbation of reproductive organ development, alteration of nitrogen, and/or carbon metabolism and limit photosynthetic activity (reference).

Saladin *et al.* reported that *in vitro* application of some fungicides, i.e. fludioxonil, and a systemic fungicide pyrimethanil, promoted different physiological responses of plants. Firstly, both fungicides decreased net CO₂ assimilation, transpiration rate, stomatal conductance, and intercellular CO₂ concentration; secondly, in the fruiting cuttings, the fungicides affected CO₂ exchange neither transpiration rates (Saladin Gaëlle *et al.*, 2003).

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374

There are paragraphs without references, add references to sentences

Strobilurin group includes synthetic substances similar in structure to natural fungicidal toxins - strobilurins A and B, isolated from the culture of microorganisms *Strobilurus tenacellus* (Balba, 2007). Strobilurins are recommended to be used first in the growing season because they rapidly reduce the ability of resistant to triazole forms to their development on leaves. In addition, the selection pressure is reduced, since the level of the inoculum is the lowest at the beginning of the growing season. Due to the wide range of action and practical safety for the environment, strobilurins are considered to be the most significant group of fungicides that appeared after the preparations of the triazole classes. These substances can be attributed to biofungicides, since they are of natural origin (Reddy, 2012). High resistance to strobilurins (for example, 200 times less sensitive to them in powdery mildew of wheat) is due to a one-point mutation in that part of the cytochrome b molecule, which determines the binding of this enzyme to fungicides. At the same time, the active centre of the enzyme does not change, and the resistant (mutant) forms of fungi do not lose their viability as a result of mutation and the acquisition of resistance to strobilurins. It is also possible the cross-resistance between strobilurins-methoxyacrylates, oximinoacetates and non-strobilurins with a similar mechanism of action-oxazolidinediones. Resistance is registered in *Oidium erysipoides*, *Erysiphe graminis*, *Botrytis cinerea*. (reference) To prevent resistance, only 1-2 treatments (in some cases, three) at intervals of 14-16 days are permitted during the season and only preparation in the fungicide alternation system with a different mechanism of action for strobilurins (Benelli Jesse ^J, 2013) are allowed. For vegetable and fruit, it is triazoles, ethylenebisdithiocarbamates, preparations based on copper and sulfur. When processing annuals on the treated area, it is necessary to practice changing cultures (Reddy, 2012).

Some reports suggested that the systemic fungicide strobilurin may improve the water status and stress management of plants under conditions of drought stress (K. Paranjape, V. Gowariker, V.N. Krishnamurthy, S. Gowariker, 2014.); (Barr *et al.*; Neiman, & Taylor, 2005). Nason *et al.* (D. S. Mueller, 2006) showed that the application of beta-methoxyacrylate, a strobilurin fungicide, improve the water use efficiency only in well-watered *Triticum aestivum* and *Hordeum vulgare* plants. However, when these plants were under drought stress, strobilurin strongly reduced net CO₂ assimilation, intercellular CO₂ concentration, transpiration rate, and rate of stomatal conductance to water. In this study, net CO₂ assimilation reduction seems to be related to stomatal conductance decrease. It is possible that stomata respond to strobilurin-induced changes in mesophyll photosynthesis either by sensing changes in the intercellular CO₂ concentration or by responding to the pool size of an unidentified C-fixing substrate. It is also possible that the effects of strobilurin fungicides are mediated via ABA-based chemical signalling (D. S. Mueller, 2006).

The analysis of several chlorophylls a fluorescence parameter of plants treated with fungicides (Xia *et al.*, 2006), 14, (D. S. Mueller, 2006), (Deising *et al.*; Reimann, & Pascholati, 2008) demonstrated that light reactions of photosynthesis are also sensitive to fungicide exposure. Bader and Abdel-Basset showed, for the first time, that fungicides of the triforine type (a systemic and contact fungicide) strongly inhibit electron-transport reactions of chloroplasts. Moreover, the application of systemic fungicides, benzimidazoles and triazole, and a dithiocarbamate contact fungicide affected the effective quantum yield of PSII as well as the maximal quantum efficiency of PSII (Fv/Fm). This reduction was attributed to the decrease in photochemical quenching (qP) (Xia *et al.*, 2006), (Deising *et al.*, 2008). In *Glycine max*, strobilurin fungicides application reduced the ratio of Fv/Fm. Strobilurin fungicides seem to block the transport of electrons between PSII and PSI by binding to the Qi site of the chloroplast cytochrome bf complex (D. S. Mueller, 2006).

Benzimidazole formulations were among the first systemic fungicides to appear on the market. Benzimidazole derivatives are effective against diseases of vegetative organs, as well as a complex of phytopathogens transmitted between seeds, so they find wide application as seed disinfectants (reference). Over time, interest in benzimidazole fungicides has fallen, in part, this is due to the emergence of resistant strains to them. Now it is difficult to evaluate how much this is related to the characteristics of the fungicides, and how much with the unpreparedness to such a consequence of their application (reference). Today, in many countries, the scope of their application has declined due to a rapid decrease in their effectiveness (reference). The narrow selectivity of the action contributes to a sufficiently rapid selection of resistant genotypes and the formation of a resistant population after a systematic (within 3-4 years) use of substantive of this group (reference). Several reports show a decrease in biomass production in fungicide-treated plants: benomyl, a systemic fungicide, reduced the growth of *Gossypium hirsutum*, *Helianthus annuus*, *Cucumis sativus*,

Formatted: Font: 20 pt

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

375 | *Lactuca sativa*, and *Pinus taeda* (Pablo C. García, Rosa M. Rivero, Juan M. Ruiz, 2003);
376 | (Hunsche *et al.*, Damerow, Schmitz-Eiberger, & Noga, 2007). Moreover, the application of
377 | carbendazim (systemic benzimidazole fungicide) in *Nicotiana tabacum* affected negatively plant
378 | biomass (Pablo C. García, Rosa M. Rivero, Juan M. Ruiz *et al.*, 2003).

379 | Pigment biosynthesis is reported by Ahmed *et al.* (Hunsche *et al.*, 2007) to be inhibited by
380 | benomyl. This fungicide induces a considerable reduction on the chlorophyll a, chlorophyll b,
381 | carotenoids, and the total pigments content of *Helianthus annuus* plants (Hunsche *et al.*, 2007).
382 | Similarly, the treatment of *Vitis vinifera* with fludioxonil and *Nicotiana tabacum* with carbendazim also
383 | decreases the chlorophyll and carotenoid content (Pablo C. García, Rosa M. Rivero, Juan M. Ruiz,
384 | 2003), (Saladin Gaëlle *et al.*, 2003). Mihuta-Grimm *et al.* (Changjun Chen, *et al.*, Jianxin
385 | Wang, Qingquan Luo, 2007) and Van Iersel and Bugbee reported leaf chlorosis after benomyl
386 | application on *Impatiens walleriana*, *Cucumis sativus*, *Celosia plumosa* *Petunia* hybrid, and
387 | *Lycopersicon esculentum* (Deising *et al.*, 2008).

388 | There is also a phenomenon of cross-resistance. Fungi that are resistant to one fungicide
389 | are often also resistant to other fungicides from the same chemical class. Sometimes between
390 | fungicides from different chemical classes, there is a negative cross-resistance. For example, one
391 | such case was identified in the study of two major pathogens (*Mycosphaerella graminicola* and
392 | *Tapesia acuformis*) of winter wheat in France. Negative cross-resistance to edifenphos and several
393 | sterol biosynthesis inhibitors, such as prochloraz and fenpropimorph, was observed in strains
394 | resistant to fenhexylamide (Leroux *et al.*, EROUX, CHAPELAND, ARNOLD, & GREDT, 2000). The
395 | reason for this phenomenon may be that a genetic modification that occurs under the action of a
396 | single fungicide and imparts resistance to it, makes the resistant isolate more susceptible to
397 | another fungicide (McGrath, 2004).

398 | Morpholines are a class of low-toxic and highly effective fungicides, one of the first groups of
399 | sterol synthesis inhibitors. They are part of the combined preparations. Although other inhibitors of
400 | sterol synthesis outperform the group of morpholines by economic parameters, these substances
401 | again acquire importance for the problem of the resistance to fungicides (Lamberth, 2012). In
402 | contrast to triazoles, morpholines block the isomerization and reduction reactions in the process of
403 | sterols biosynthesis, therefore the populations of fungi that are resistant to them are formed much
404 | more slowly. According to the spectrum of action on pathogens, morpholines do not differ from
405 | triazoles but require higher application rates. Despite the slow development of resistant strains,
406 | there is a potential for dimethomorph to develop resistant strains of pathogens that do not have
407 | cross-resistance to phenylamides.

408 | There are cases of phytotoxicity with substances from other chemical classes. In study Yuba
409 | R. Kandel, Daren S. Mueller and etc. (Kandel *et al.*, 2018) says that preemergence herbicides and
410 | seed treatment fluopyram each has led to increased phytotoxicity in the VC-V1 growth stage in
411 | soybean compared to the untreated control. Physiological studies after fungicide application on
412 | several species reported modifications of both photosynthetic activity and chlorophyll a
413 | fluorescence [(Saladin Gaëlle *et al.*, 2003). Decreased CO₂ assimilation in fungicide-treated plants
414 | is attributed to both stomatal (due to stomatal closure) (Xia *et al.*, 2006) and nonstomatal effects
415 | due to a disruption in the capacity of RuBisCO carboxylation, decrease of RuBisCO content, and/or
416 | reduction of the ribulose 1.5 biphosphate regeneration (Anne-Nolle Petit *et al.*, n.d. (D. S.
417 | Mueller, 2006).

418 | Modifications of dark respiration were reported after mancozeb (contact fungicide) and
419 | flusilazol (systemic fungicide) application in *Malus domestica*. The increase in dark respiration can
420 | be explained by additional energy requirement, metabolic breakdown of the compound, and/or
421 | activation of the alternative, cyanide-insensitive, respiration. Curiously, the treatment with
422 | strobilurin fungicides induced different responses: while in *Triticum aestivum* and in *Spinacia*
423 | *oleracea* plants respiration was inhibited (K. Paranjape, V. Gowariker, V.N. Krishnamurthy, S.
424 | Gowariker, 2014), (Pantazopoulou & Diallinas, 2007) in *Triticum aestivum* dark respiration was
425 | reduced (D. S. Mueller, 2006).

426 | The most crucial aspect of work of fungicides is their efficiency against fungal pathogens or
427 | their residues in crops (Report on the pesticide residues monitoring programme: Quarter 1 2017,
428 | 2017), (Saladin Gaëlle *et al.*, 2003)]. Several reports found that some fungicides can improve plant
429 | defences through phytoalexin synthesis and cell wall lignification or stimulate enzymes involved in
430 | the synthesis of phenolic compounds [(Saladin Gaëlle *et al.*, 2003); (War *et al.*, 2012). Others
431 | describe the supposed protective role of fungicides for crops against various types of stress
432 | factors. Wu and Von Tiedemann (Anne-Noëlle Petit, Fontaine, Clément, & Vaillant-Gaveau, 2008),
433 | (Untiedt & Blanke, 2004) described the protective function of triazoles in *Hordeum vulgare* and
434 | *Arachis hypogaea* against ozone exposure or salt stress by stimulating antioxidative enzymes.

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

435 Furthermore, azoxystrobin and epoxiconazole were shown to retard senescence of *Triticum*
436 *aestivum* primarily due to an expansion of the antioxidative potential protecting the plants from
437 damage by active oxygen species (Untiedt & Blanke, 2004). Muthukumarasamy and
438 Panneerselvam described the induction of the synthesis of photosynthetic pigments and proteins in
439 treated plants (Indian Council Of Agricultural Research, 2011). However, only small number of
440 studies have considered the question of whether these products boost or inhibit physiological and
441 metabolic activities in the plant tissues (Pablo *C. Garcia, Rosa M. Rivero, Juan M. Ruiz et al.*,
442 2003), and the negative impact of fungicides on photosynthesis, pigment content, growth, and
443 alterations in the reproductive organs was poorly analyzed (Anne-Nolle Petit et al., n.d.), (Saladin
444 Gaëlle *et al.*, 2003).

Formatted: Font: Italic

Formatted: Font: Italic

445 The decrease in photosynthesis rate intensely influences plant biomass production and
446 growth rates. Information about fungicide effects on plant physiology (especially on photosynthesis)
447 is decisive for the understanding of the primary regulatory mechanisms and the phytotoxicity of a
448 given compound ([reference](#)).

449 8. MYCORRHIZAL FUNGI RESPONSES

450 Fungicidal compositions for seeds containing a multi-ingredient system are targeted at
451 multiple metabolic processes. And many researchers in this field are concerned with the question:
452 can these fungicides to inhibit inappropriate soil fungi, such as obligate plant symbiotic arbuscular
453 mycorrhizal (AM) fungi (AMF).

454 Arbuscular mycorrhizal fungi are symbionts of plants, which interrelate with approximately 80% of
455 plant species (J. Cameron, 2016). For example, multilateral interactions between roots and
456 mycorrhizal fungi can have a synergistic effect on the growth and systemic priming of wheat
457 (Pérez-de-Luque *et al.*, 2017). These symbionts often have a beneficial effect on the host plant,
458 increasing nutrient intake and tolerance to biotic and abiotic stresses, improving soil quality in
459 cropping systems.

Formatted: Font: Italic

460 The study of Huan Jing Ke Xue ([year????](#)) says that in the treatment with benomyl, the
461 content of K in the shoot and the Fe in the root decreased significantly in mycorrhizal plants; in the
462 treatment with difenoconazole, the total N and K content in the shoot also decreased, Ca in the
463 roots; mycorrhizal colonization, total P, K and Cu content in the shoot, the total amount of N, Ca,
464 Zn and Fe in the root was significantly reduced with flusilazole. The inhibitory effect of flusilazole
465 on the colonization of *Glomus mosseae* and the growth of *Scutellaria baicalensis* were higher than
466 with difenoconazole and benomyl (He *et al.*, Wang, Ma, & Meng, 2012).

Formatted: Font: Italic

467 But in other studies, in the analysis of corn (*Zea mays* L.), soybean (*Glycine max* L.) and
468 oats (*Avena sativa* L.) treated with azoxystrobin, fludioxonil, mefenoxam, trifloxystrobin, and
469 pyraclostrobin, no found significant effect on AM fungal colonization (J. C. Cameron *et al.*,
470 Lehman, Sexton, Osborne, & Taheri, 2017). Fungicides were applied according to
471 the recommended dosages. In small amounts, the following negative effects were observed. Corn
472 treated by Cruiser Extreme had significantly lower ($P < 0,05$) colonization of AM fungi compared to
473 the other two fungicides (Trilex, Stamina) and tended to decrease the colonization of AM corn roots
474 as compared to controls ($P = 0,08$). The Cruiser Extreme consists of a locally systemic fungicide
475 (azoxystrobin) inhibiting respiration, a systemic fungicide (mekenoxane) inhibiting the synthesis of
476 nucleic acids, and a contact fungicide (fludioxonil), which prevents the transduction of cells
477 ([reference](#)).

Formatted: Font: Italic

478 However, in the analysis of soy, the same relation was not found. In oats, the results were
479 lower than the rest, but not lower than the controls ([reference](#)). The differences in the colonization of
480 AM fungal between fungicidal medication, apparently, are not related to a particular mode of action.
481 There was no relationship between the treatment of fungicide and plant genotype during
482 colonization of AM fungi or the content of plant nutrients ([reference](#)). The plant genotype has a
483 consistent effect on the colonization of AM fungi and the nutrient content of plants.

484 Schreiner and Bethlenfalvai have shown that a higher variety of AMF can better withstand
485 the negative effects of fungicides (Schreiner & Bethlenfalvai, 1997). The essential role of fungicidal
486 action on AMF can be played by their movement in the plant. As a rule, contact fungicides are less
487 harmful than systemic fungicides when using seeds measured by sporulation, glomalin and
488 biomass of the host plant (Hongyan, Germida, & Walley, 2013).

489 Murillo-Williams and Pedersen found that fludioxonil in treated seed had a positive effect
490 on the AMF colonization in soy (*Glycine max* L.) due to a decrease in competition with the
491 aggressive pathogen *Rhizoctonia* spp. (Murillo-Williams & Pedersen, 2008). But in another case,
492 fludioxonil had no significant effect on the colonization of AMF in onions (Hernández-Dorrego &
493 Mestre-Parés, 2010). Thus, the potential negative effects of systemic and contact fungicides on

Formatted: Font: Italic

Formatted: Font: Not Italic

494 | non-targeted, useful AMF are not fully understood and studied (reference). With the recent
495 introduction of commercial modified AMF for large-scale crop production, understanding the effects
496 of fungicides on these beneficial organisms can help minimize the unintentional interactions
497 between fungicides and AMF.

498 7. CONCLUSION

499 Fungicides are widely used and have become the main means of inhibiting the growth of
500 fungi and fungal spores due to their relatively low cost, high efficiency and ease of use.

501 However, despite the wide variety of existing products and various routes of use, the
502 problem of the emergence of new fungicide-resistant strains of pathogens remains open. Available
503 studies have demonstrated that fungicide application may impair photosynthesis, the synthesis of
504 sterols, gibberellins, transpiration, reduce CO₂ assimilation and biomass, influence on the total
505 pigments content. However, reports on phytotoxicity are generally based on a few physiological
506 parameters using a large variety of plant species and different types and concentrations of
507 fungicides, leading in some cases to contradictory results. This significantly jeopardizes a
508 comprehensive knowledge on the primary effects of fungicides on the photosynthesis and certainly
509 deserves further investigation.

510 It may be worthwhile to study in more detail methods for predicting the spread of diseases
511 and testing theories during the development of fungicides using machine learning (i.e. artificial
512 neural network). And as an attractive aspect for further fungicide study are such aspects as cross-
513 resistance and negative cross-resistance of different chemical classes fungicides. This knowledge
514 would be extremely useful when developing new preparations.

515 Furthermore, the problem of the negative impact of fungicides on the environment due to
516 their high toxicity still remains unresolved. However, the situation can be improved with the use of
517 new technologies and a deeper understanding of the fungicides mechanism of action. Because it
518 allows to create preparations with a lower content of active substance, but not less effective. The
519 solution to that problem will provide benefits not only for plants yield but also for the environment
520 and human health.

521 Concerns about the non-targeted effects of fungicides on AMF are mainly focused on the
522 potential impact on natural AMF in integrated management systems. However, understanding the
523 compatibility of fungicides used for seeds, not only with natural but with modified useful AMF, is
524 important if we want to maximize the benefits of both, obtained from sowing crops.

525 COMPETING INTERESTS

526 Authors have declared that no competing interests exist.

527 AUTHORS' CONTRIBUTIONS

528 All authors read and approved the final manuscript.

529 REFERENCES **Check all references**

- 530 Ahemad, M., & Khan, M. S. (2012). Alleviation of fungicide-induced phytotoxicity in greengram
531 [*Vigna radiata* (L.) Wilczek] using fungicide-tolerant and plant growth promoting
532 *Pseudomonas* strain. *Saudi Journal of Biological Sciences*, 19(4), 451–459.
533 <https://doi.org/10.1016/J.SJBS.2012.06.003>
- 534 Aynalem, B., & Assefa, F. (2017). Effect of Glyphosate and Mancozeb on the Rhizobia Isolated
535 from Nodules of *Vicia faba* L. and on Their N₂-Fixation, North Showa, Amhara Regional
536 State, Ethiopia. *Advances in Biology*, 2017, 1–7. <https://doi.org/10.1155/2017/5864598>
- 537 Balba, H. (2007). Review of strobilurin fungicide chemicals. *Journal of Environmental Science and*
538 *Health, Part B*, 42(4), 441–451. <https://doi.org/10.1080/03601230701316465>
- 539 Barr, C. M., Neiman, M., & Taylor, D. R. (2005). Inheritance and recombination of mitochondrial
540 genomes in plants, fungi and animals. *New Phytologist*, 168(1), 39–50.
541 <https://doi.org/10.1111/j.1469-8137.2005.01492.x>
- 542 Benelli Jesse J. (2013). *Non-target effects of strobilurin fungicide applications on creeping*
543 *bentgrass putting greens during summer stress*. University of Tennessee. Retrieved from
544 http://trace.tennessee.edu/utk_gradthes/1595
- 545 Bernauer, O., Gaines-Day, H., & Steffan, S. (2015). Colonies of Bumble Bees (*Bombus impatiens*)
546 Produce Fewer Workers, Less Bee Biomass, and Have Smaller Mother Queens Following
547 Fungicide Exposure. *Insects*, 6(4), 478–488. <https://doi.org/10.3390/insects6020478>
- 548 Biol, T. J., Jităreanu, A., Pădureanu, S., Tătăringă, G., Tuchiluş, C., & Stănescu, U. (2013).
549 Evaluation of phytotoxic and mutagenic effects of some cinnamic acid derivatives using the

Formatted: Font: 16 pt

550 Triticum test. *Turkish Journal of Biology*, 748–756. <https://doi.org/10.3906/biy-1304-39>

551 Cameron, J. (2016). *Effects of Seed Applied Fungicide on Arbuscular Mycorrhizal Colonization of*

552 *South Dakota Cultivars of Oat, Soybean, and Corn*. South Dakota State University. Retrieved

553 from <http://openprairie.sdstate.edu/etd>

554 Cameron, J. C., Lehman, R. M., Sexton, P., Osborne, S. L., & Taheri, W. I. (2017). Fungicidal Seed

555 Coatings Exert Minor Effects on Arbuscular Mycorrhizal Fungi and Plant Nutrient Content.

556 *Agronomy Journal*, 109(3), 1005–1012. <https://doi.org/10.2134/agronj2016.10.0597>

557 Changjun Chen, Jianxin Wang, Qingquan Luo, S. Y. andMingguo Z. (2007). Characterization and

558 fitness of carbendazim-resistant strains of *Fusarium graminearum* (wheat scab). *Pest*

559 *Management Science*, 63(12), 1201–1207. <https://doi.org/10.1002/ps.1449>

560 Clayton A. Hollier, Jeffrey W. Hoy, Christopher A. Clark, Charles Overstreet, Jaspreet Sidhu,

561 Melanie L. Lewis Ivey, Raghuwinder Singh, Trey Price III, Mary Helen Ferguson, G. Boyd

562 Padgett, D. G. (2016). *Louisiana Plant Disease Management Guide. Plant Disease*.

563 Louisiana. Retrieved from file:///C:/Users/ekate_000/Desktop/Pub1802-

564 plant_management_guide.pdf

565 Cools, H. J., Hawkins, N. J., & Fraaije, B. A. (2013). Constraints on the evolution of azole

566 resistance in plant pathogenic fungi. *Plant Pathology*, 62, 36–42.

567 <https://doi.org/10.1111/ppa.12128>

568 Costa, A. V., Oliveira, M. V. L. de, Pinto, R. T., Moreira, L. C., Gomes, E. M. C., Alves, T. de A., ...

569 Júnior, W. C. de J. (2017). Synthesis of Novel Glycerol-Derived 1,2,3-Triazoles and

570 Evaluation of Their Fungicide, Phytotoxic and Cytotoxic Activities. *Molecules*, 22(10), 1666.

571 <https://doi.org/10.3390/molecules22101666>

572 Deising, H. B., Reimann, S., & Pascholati, S. F. (2008). Mechanisms and significance of fungicide

573 resistance. *Brazilian Journal of Microbiology*, 39(2), 286–295. [https://doi.org/10.1590/S1517-](https://doi.org/10.1590/S1517-83822008000200017)

574 [83822008000200017](https://doi.org/10.1590/S1517-83822008000200017)

575 Dias Maria Celeste. (2012). Phytotoxicity: An Overview of the Physiological Responses of Plants

576 Exposed to Fungicides. *Journal of Botany*, 2012, 1–4. <https://doi.org/10.1155/2012/135479>

577 Donatelli, M., Magarey, R. D., Bregaglio, S., Willocquet, L., Whish, J. P. M., & Savary, S. (2017).

578 Modelling the impacts of pests and diseases on agricultural systems. *Agricultural Systems*,

579 155, 213–224. <https://doi.org/10.1016/J.AGSY.2017.01.019>

580 E. Lee Butler, L. P. T. (2006). Method and Timing of Fungicide Applications for Control of Spring

581 Dead Spot In Hybrid Bermudagrass. *Plant Health Progress*. [https://doi.org/10.1094/PHP-](https://doi.org/10.1094/PHP-2006-0901-01-RS)

582 [2006-0901-01-RS](https://doi.org/10.1094/PHP-2006-0901-01-RS)

583 Elslahi, R. H., Osman, A. G., Sherif, A. M., & Elhoussein, A. A. (2014). Comparative study of the

584 fungicide Benomyl toxicity on some plant growth promoting bacteria and some fungi in pure

585 cultures. *Interdisciplinary Toxicology*, 7(1), 12–16. <https://doi.org/10.2478/intox-2014-0002>

586 Forrer, H.-R., Musa, T., Schwab, F., Jenny, E., Bucheli, T. D., Wettstein, F. E., & Vogelgsang, S.

587 (2014). *Fusarium* head blight control and prevention of mycotoxin contamination in wheat with

588 botanicals and tannic acid. *Toxins*, 6(3), 830–849. <https://doi.org/10.3390/toxins6030830>

589 González M., Caetano P., S. M. E. (2017). Testing systemic fungicides for control of *Phytophthora*

590 oak root disease. *Forest Pathology*, 47(4), 1–3. <https://doi.org/10.1111/efp.12343>

591 Hasan, M. A., Ahmed, J. U., Tofazzal, H., Mian, M. A. K., & Haque, M. M. (2013). Evaluation of the

592 physiological quality of wheat seed as influenced by high parent plant growth temperature.

593 *Journal of Crop Science and Biotechnology*, 16(1), 69–74. [https://doi.org/10.1007/s12892-](https://doi.org/10.1007/s12892-010-0056-1)

594 [010-0056-1](https://doi.org/10.1007/s12892-010-0056-1)

595 He, X.-L., Wang, P., Ma, L., & Meng, J.-J. (2012). Effects of three fungicides on arbuscular

596 mycorrhizal fungal infection and growth of *Scutellaria baicalensis* Georgi. *Huan Jing Ke Xue=*

597 *Huanjing Kexue*, 33(3), 987–91. Retrieved from

598 <http://www.ncbi.nlm.nih.gov/pubmed/22624398>

599 Hernández-Dorrego, A., & Mestre-Parés, J. (2010). Evaluation of some fungicides on mycorrhizal

600 symbiosis between two *Glomus* species from commercial inocula and *Allium porrum* L.

601 seedlings. *Spanish Journal of Agricultural Research*, 8(S1), 43.

602 <https://doi.org/10.5424/sjar/201008S1-1222>

603 Hongyan, Germida, J. J., & Walley, F. L. (2013). Suppressive effects of seed-applied fungicides on

604 arbuscular mycorrhizal fungi (AMF) differ with fungicide mode of action and AMF species.

605 *Applied Soil Ecology*, 72, 22–30. Retrieved from [http://agris.fao.org/agris-](http://agris.fao.org/agris-search/search.do?recordID=US201400165252)

606 [search/search.do?recordID=US201400165252](http://agris.fao.org/agris-search/search.do?recordID=US201400165252)

607 Hunsche, M., Damerow, L., Schmitz-Eiberger, M., & Noga, G. (2007). Mancozeb wash-off from

608 apple seedlings by simulated rainfall as affected by drying time of fungicide deposit and rain

609 characteristics. *Crop Protection*, 26(5), 768–774. <https://doi.org/10.1016/j.cropro.2006.07.003>

610 Indian Council Of Agricultural Research. (2011). *Handbook of agriculture* (6th ed.). New Delhi:
611 Directorate of Information and Publications of Agriculture, Indian Council of Agricultural
612 Research.

613 Isaac, S. (1992). *Fungal-plant interactions* (1. ed.). London: Chapman & Hall. Retrieved from
614 <http://www.worldcat.org/title/fungal-plant-interactions/oclc/231386069>

615 Johnson, D. A., Hamm, P. B., & Sunseri, M. (n.d.). Fungicide Application for Management of Potato
616 Late Blight. Retrieved from http://public.wsu.edu/~djohnsn/index_files/Paper4.pdf

617 K. Paranjape, V. Gowariker, V.N. Krishnamurthy, S. Gowariker, K. P. (2014). *The Pesticide*
618 *Encyclopedia* (UK ed. edi). London: CABI.

619 Kandel, Y. R., Mueller, D. S., Legleiter, T., Johnson, W. G., Young, B. G., & Wise, K. A. (2018).
620 Impact of fluopyram fungicide and preemergence herbicides on soybean injury, population,
621 sudden death syndrome, and yield. *Crop Protection*, *106*, 103–109.
622 <https://doi.org/10.1016/j.cropro.2017.12.009>

623 Kilani, J., & Fillingner, S. (2016). Phenylpyrroles: 30 Years, Two Molecules and (Nearly) No
624 Resistance. *Frontiers in Microbiology*, *7*, 2014. <https://doi.org/10.3389/fmicb.2016.02014>

625 Kuna-Broniowski, M., Makarski, P., & Kuna-Broniowska, I. (2015). Application of Electric Fields as
626 a Method for Plant Disease Forecasting. *Agriculture and Agricultural Science Procedia*, *7*,
627 146–151. <https://doi.org/10.1016/J.AASPRO.2015.12.009>

628 Lamberth, C. (2012). Morpholine Fungicides for the Treatment of Powdery Mildew. In *Bioactive*
629 *Heterocyclic Compound Classes* (pp. 119–127). Weinheim, Germany: Wiley-VCH Verlag
630 GmbH & Co. KGaA. <https://doi.org/10.1002/9783527664412.ch10>

631 LEROUX, P., CHAPELAND, F., ARNOLD, A., & GREDET, M. (2000). New Cases of Negative
632 Cross-resistance between Fungicides, Including Sterol Biosynthesis Inhibitors. *Journal of*
633 *General Plant Pathology*, *66*(1), 75–81. <https://doi.org/10.1007/PL00012925>

634 Lesemann, S. S., Schimpke, S., Dunemann, F., & Deising, H. B. (2006). Mitochondrial
635 heteroplasmy for the cytochrome b gene Controls the level of strobilurin resistance in the
636 apple powdery mildew fungus *Podosphaera leucotricha* (Ell. & Ev.) E.S. Salmon. *Journal*
637 *of Plant Diseases and Protection*, *113*(6), 259–266. <https://doi.org/10.1007/BF03356191>

638 Lew, R. R. (2010). Turgor and net ion flux responses to activation of the osmotic MAP kinase
639 cascade by fludioxonil in the filamentous fungus *Neurospora crassa*. *Fungal Genetics and*
640 *Biology*, *47*(8), 721–726. <https://doi.org/10.1016/J.FGB.2010.05.007>

641 Marín, S., Ramos, A. J., Cano-Sancho, G., & Sanchis, V. (2012). Reduction of mycotoxins and
642 toxigenic fungi in the Mediterranean basin maize chain. *Phytopathologia Mediterranea*, *51*(1),
643 93–118. Retrieved from www.fupress.com/pm

644 McGrath, M. T. (2004). What are Fungicides? *The Plant Health Instructor*.
645 <https://doi.org/10.1094/PHI-I-2004-0825-01>

646 Miguez, M., Reeve, C., Wood, P. M., & Hollomon, D. W. (2004). Alternative oxidase reduces the
647 sensitivity of *Mycosphaerella graminicola* to QOI fungicides. *Pest Management Science*, *60*(1),
648 3–7. <https://doi.org/10.1002/ps.837>

649 Ming-wang Shi. (2011). Based on time series and RBF network plant disease forecasting. *Procedia*
650 *Engineering*, *15*, 2384–2387. <https://doi.org/10.1016/J.PROENG.2011.08.447>

651 Mueller, D. (n.d.). Field Crop Fungicides For the north Central united states. Retrieved from
652 <https://www.ncipmc.org/action/FungicideManual4.pdf>

653 Mueller, D. S. (2006). Fungicides: Terminology. *Integrated Crop Management*, 120–123. Retrieved
654 from <http://lib.dr.iastate.edu/cropnews>

655 Murillo-Williams, A., & Pedersen, P. (2008). Arbuscular Mycorrhizal Colonization Response to
656 Three Seed-Applied Fungicides. *Agronomy Journal*, *100*(3), 795.
657 <https://doi.org/10.2134/agronj2007.0142>

658 Nason, M. A., Farrar, J., & Bartlett, D. (2007). Strobilurin fungicides induce changes in
659 photosynthetic gas exchange that do not improve water use efficiency of plants grown under
660 conditions of water stress. *Pest Management Science Pest Manag Sci*, *63*, 1191–1200.
661 <https://doi.org/10.1002/ps.1443>

662 Oliver, R. P., & Hewitt, H. G. (2014). *Fungicides in crop protection* (2th ed.). CAB International.
663 Retrieved from
664 [https://books.google.pl/books/about/Fungicides_in_Crop_Protection_2nd_Editio.html?id=4x0x](https://books.google.pl/books/about/Fungicides_in_Crop_Protection_2nd_Editio.html?id=4x0xBQAAQBAJ&redir_esc=y)
665 [BQAAQBAJ&redir_esc=y](https://books.google.pl/books/about/Fungicides_in_Crop_Protection_2nd_Editio.html?id=4x0xBQAAQBAJ&redir_esc=y)

666 Pablo C. García, Rosa M. Rivero, Juan M. Ruiz, L. R. (2003). The Role of Fungicides in the
667 Physiology of Higher Plants: Implications for Defense Responses. *The Botanical Review*,
668 *69*(2), 162–172. [https://doi.org/10.1663/0006-8101\(2003\)069\[0162:TROFIT\]2.0.CO;2](https://doi.org/10.1663/0006-8101(2003)069[0162:TROFIT]2.0.CO;2)

669 Pantazopoulou, A., & Diallinas, G. (2007). Fungal nucleobase transporters. *FEMS Microbiology*

670 *Reviews*, 31(6), 657–675. <https://doi.org/10.1111/j.1574-6976.2007.00083.x>

671 Paul Vincelli, Bruce Clarke, G. M. (2017). *Chemical Control of Turfgrass Diseases 2017*. University
672 of Kentucky College of Agriculture, Food and Environment Chemical Control of Cooperative
673 Extension Service. Retrieved from <http://www2.ca.uky.edu/agcomm/pubs/ppa/ppa1/ppa1.PDF>

674 Pérez-de-Luque, A., Tille, S., Johnson, I., Pascual-Pardo, D., Ton, J., & Cameron, D. D. (2017).
675 The interactive effects of arbuscular mycorrhiza and plant growth-promoting rhizobacteria
676 synergistically enhance host plant defences against pathogens. *Scientific Reports*, 7(1).
677 <https://doi.org/10.1038/s41598-017-16697-4>

678 Petit, A.-N., Fontaine, F., Clément, C., & Vaillant-Gaveau, N. (2008). Photosynthesis Limitations of
679 Grapevine after Treatment with the Fungicide Fludioxonil. *Journal of Agricultural and Food
680 Chemistry*, 56(15), 6761–6767. <https://doi.org/10.1021/jf800919u>

681 Petit, A.-N., Fontaine, F., Ement, C. C., & Vaillant-Gaveau, N. (n.d.). Photosynthesis Limitations of
682 Grapevine after Treatment with the Fungicide Fludioxonil. <https://doi.org/10.1021/jf800919u>

683 Reddy, P. P. (2012). Strobilurin Fungicides. In *Recent advances in crop protection* (pp. 185–200).
684 New Delhi: Springer India. https://doi.org/10.1007/978-81-322-0723-8_12

685 Ren, W., Shao, W., Han, X., Zhou, M., & Chen, C. (2016). Molecular and Biochemical
686 Characterization of Laboratory and Field Mutants of *Botrytis cinerea* Resistant to Fludioxonil.
687 *Plant Disease*, 100(7), 1414–1423. <https://doi.org/10.1094/PDIS-11-15-1290-RE>

688 *Report on the pesticide residues monitoring programme: Quarter 1 2017*. (2017). London.
689 Retrieved from
690 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/655056/pesticide](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/655056/pesticide-residues-quarter1-2017-report.pdf)
691 [e-residues-quarter1-2017-report.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/655056/pesticide-residues-quarter1-2017-report.pdf)

692 Rohr, J. R., Brown, J., Battaglin, W. A., McMahon, T. A., & Relyea, R. A. (2017). A pesticide
693 paradox: fungicides indirectly increase fungal infections. *Ecological Applications*, 27(8), 2290–
694 2302. <https://doi.org/10.1002/eap.1607>

695 Saladin Gaëlle, Magné Christian, C. C., & Clément, C. (2003). Effects of fludioxonil and
696 pyrimethanil, two fungicides used against *Botrytis cinerea*, on carbohydrate physiology in *Vitis*
697 *vinifera* L. *Pest Management Science*, 59(10), 1083–1092. <https://doi.org/10.1002/ps.733>

698 Schreiner, R. P., & Bethlenfalvai, G. J. (1997). Plant and soil response to single and mixed species
699 of arbuscular mycorrhizal fungi under fungicide stress. *Applied Soil Ecology*, 7, 93–102.
700 Retrieved from
701 <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.579.3176&rep=rep1&type=pdf>

702 Shuping, D. S. S., & Eloff, J. N. (2017). THE USE OF PLANTS TO PROTECT PLANTS AND
703 FOOD AGAINST FUNGAL PATHOGENS: A REVIEW. *African Journal of Traditional,
704 Complementary, and Alternative Medicines: AJTCAM*, 14(4), 120–127.
705 <https://doi.org/10.21010/ajtcam.v14i4.14>

706 Southern AG. (2015). *LIQUID COPPER FUNGICIDE* (No. 10465-3–829). New Jersey . Retrieved
707 from <https://www.domyown.com/msds/SDSCopper.pdf>

708 Tom Allen. (2013). Not Everything is as it Seems: Fungicide Phytotoxicity and Plant Diseases |
709 Mississippi Crop Situation. Retrieved January 9, 2018, from [http://www.mississippi-](http://www.mississippi-crops.com/2013/08/09/not-everything-is-as-it-seems-fungicide-phytotoxicity-and-plant-diseases/)
710 [crops.com/2013/08/09/not-everything-is-as-it-seems-fungicide-phytotoxicity-and-plant-](http://www.mississippi-crops.com/2013/08/09/not-everything-is-as-it-seems-fungicide-phytotoxicity-and-plant-diseases/)
711 [diseases/](http://www.mississippi-crops.com/2013/08/09/not-everything-is-as-it-seems-fungicide-phytotoxicity-and-plant-diseases/)

712 Untiedt, R., & Blanke, M. M. (2004). Effects of fungicide and insecticide mixtures on apple tree
713 canopy photosynthesis, dark respiration and carbon economy. *Crop Protection*, 23(10), 1001–
714 1006. <https://doi.org/10.1016/j.cropro.2004.02.012>

715 Vincelli P. (2002). QoI (Strobilurin) Fungicides: Benefits and Risks. *The Plant Health Instructor*.
716 <https://doi.org/10.1094/PHI-I-2002-0809-02>

717 Walker, A.-S., Micoud, A., Rémuson, F., Grosman, J., Gredt, M., & Leroux, P. (2013). French
718 vineyards provide information that opens ways for effective resistance management of
719 *Botrytis cinerea* (grey mould). *Pest Management Science*, 69(6), 667–678.
720 <https://doi.org/10.1002/ps.3506>

721 War, A. R., Paulraj, M. G., Ahmad, T., Buhroo, A. A., Hussain, B., Ignacimuthu, S., & Sharma, H.
722 C. (2012). Mechanisms of plant defense against insect herbivores. *Plant Signaling &
723 Behavior*, 7(10), 1306–20. <https://doi.org/10.4161/psb.21663>

724 Wojdyła, A. T. (2007). Influence of strobilurin compounds on the development of *Puccinia horiana*.
725 *Communications in Agricultural and Applied Biological Sciences*, 72(4), 961–6. Retrieved
726 from <http://www.ncbi.nlm.nih.gov/pubmed/18396835>

727 Woodward, J. E., Russell, S. A., Baring, M. R., Cason, J. M., & Baughman, T. A. (2015). Effects of
728 Fungicides, Time of Application, and Application Method on Control of Sclerotinia Blight in
729 Peanut. *International Journal of Agronomy*, 8. <https://doi.org/10.1155/2015/323465>

730 Xia, X. J., Huang, Y. Y., Wang, L., Huang, L. F., Yu, Y. L., Zhou, Y. H., & Yu, J. Q. (2006).
731 Pesticides-induced depression of photosynthesis was alleviated by 24-epibrassinolide
732 pretreatment in *Cucumis sativus* L. *Pesticide Biochemistry and Physiology*, 86(1), 42–48.
733 <https://doi.org/10.1016/J.PESTBP.2006.01.005>
734

UNDER PEER REVIEW