

**Antibiotic Susceptibility profile and prevalence Pattern of of
Gram negative pathogens in Tertiary Care Hospital**

**Pradnya Joshi¹, Z A Khan², Reema Tandle³, Amol Harshe⁴, Amruta Bhutada⁵,
Sunita Gogavale⁶**

¹Consultant Microbiologist, Noble Hospital

²Director ICU, Noble Hospital

³HOD Medicine Department, Noble Hospital

⁴Consultant Pathologist, HOD Pathology & Microbiology, Noble Hospital

⁵Technician Microbiology Department, Noble Hospital

⁶Technician Microbiology Dept, Noble Hospital

Pradnya Joshi (Email:drvenrem@gmail.com)

Corresponding author

Pradnya Joshi

Consultant Microbiologist, Noble Hospital

Pune, Maharashtra 411013

Email ID: joshiprandya@gmail.com

25 **Abstract**

26 ***Background and objective***

27 Large amounts of antibiotics consumed by the human population have resulted in the
28 culmination of pathogenic bacteria resistant to multiple drugs. The resistance profile of
29 pathogens differ from one geographical location to another and keeps on changing
30 continuously.

31 ***Methods***

32 A retrospective observational analysis of antibiogram data was performed to characterize
33 the susceptibility pattern of different pathogen isolates from various clinical sources. A
34 total of 213 clinical isolates identified from the period June 2015 to June 2016 were
35 included in the study.

36 ***Results***

37 Of the 213 Gram-negative isolates, 36.6% were from urine, 23.9% from respiratory
38 specimens, 11.74% from blood, 10.33% from pus whereas 17.37% were from other
39 sources. *E. coli* (42.25%) was most predominant pathogen isolated followed by *K.*
40 *pneumoniae* . (25.35%) and *Pseudomonas spp.* (15.96%) while other Gram-negative
41 pathogens contributed 16.4%. Antibiogram analysis has shown CSE-1034 as the most
42 susceptible drug exhibiting 91.1%, 77.8%, 82.4% and 82.3% susceptibility against *E. coli*,
43 *K. pneumoniae*, *A. baumannii* and *P. aeruginosa*. Among carbapenems, both meropenem
44 and imipenem-Cilastin were most effective against *E. coli*. Meropenem was least effective
45 against *K. pneumoniae* (50%) and imipenem against *P. aeruginosa* (32.35%).
46 Like imipenem, Piperacillin-Tazobactam was highest effective against *E. coli* (20%) and
47 lowest against *P. aeruginosa* (26.47%).

48

49 ***Conclusion***

50 Susceptibility profile indicates CSE-1034 (a novel antibiotic resistance breaker) as the
51 most effective susceptible drug among all the classes of antibiotics against the Gram-
52 negative pathogens. A high resistance to piperacillin-tazobactam and penems, advocates
53 use of CSE-1034 as empiric drug of choice in the treatment of bacterial infectious
54 diseases where the pathogen isolates are suspected resistant towards β -lactam and β -
55 lactamase inhibitor combinations.

56 **Keywords:** Antibiotic, Clinical isolates, CSE-1034, Prevalence, Susceptibility,
57 Resistance.

58

59 **1. Introduction**

60 The emergence of resistance among pathogenic bacteria towards potent antimicrobial
61 agents has become a critical problem in modern medicine [1]. WHO has warned that the
62 level of resistance to drugs used to treat common infectious diseases is arriving at a crisis
63 point and if not controlled, the entire population could be wiped out by these superbugs
64 [2]. The developing resistance towards currently available drugs increases the economic
65 burden on the community by increasing the rates of hospitalization, length of hospital
66 stays and cost of treatment [3] [4][5]. The rising antimicrobial resistance among the most
67 common opportunistic Gram-negative pathogens are also associated with increased
68 mortality and morbidity rates [2].

69 β -lactam antibiotics used to be the most common treatment for bacterial infections but
70 the constant exposure of bacteria to β -lactams drugs has created a selective pressure
71 leading to ESBL and ~~MBL~~ carbapenemase-producing strains including MBLs. [6]. In past
72 few years, a significant increase in the prevalence of ~~ESBL, MBL and~~ and
73 Carbapenemase producing strains including MBLs has been observed throughout the

74 globe [7]. These beta-lactamase producing Gram-negative pathogens are reported
75 resistant to other classes of antibiotics also [12,13].

76 Taking into account such a situation, there is a need to optimize the antibiotic therapy
77 against multidrug-resistant pathogens which may vary from one geographical locale to
78 another. Surveillance data and hospital antibiogram profiles help clinicians in the
79 prescription of appropriate antimicrobial therapy. Therefore, we aimed to study the
80 susceptibility profile of clinical isolates collected from Noble Hospital, Pune towards
81 commonly used 2nd line antibiotics including Ceftriaxone/Sulbactam/EDTA, β -lactam and
82 β -lactamase inhibitor combination (Piperacillin-tazobactam) and Carbapenems
83 (meropenem and imipenem-cilastatin) drugs.

84 **2. Materials and Methods**

85 ***2.1 Sample collection***

86 Various clinical specimens used for pathogen isolation included urine, stool, blood, pus,
87 endotracheal tube secretions (ETT), tracheal tube (TT) secretions, sputum, wound, , gall
88 bladder specimens, abscess, drain, ear swab, vitreous eye, abdominal fluid, vitreous fluid,
89 semen, peritoneal fluid and tissue specimens collected from 347 infected patients at Noble
90 Hospital, Pune (India), during the period of July 2016 to February 2017. The collection
91 and processing of the samples were done as per common standard operating procedures of
92 the hospital.

93 ***2.2 Isolation and Identification of microbes***

94 All the samples were collected aseptically in sterile containers and inoculated on the
95 different selective and non-selective culture media as per the standard microbiological
96 techniques. Details of the culture media used for the isolation of pathogens from various
97 clinical samples are given in Table 1. Blood samples were collected in Bactec bottles and
98 incubated in Bactec machine. These samples were further sub-cultured on the selective or

99 non-selective media and incubated aerobically overnight at 37°C. Organisms were
100 identified on the basis of colony morphology, gram staining, motility and biochemical
101 reactions. Biochemical reactions were performed by inoculating the bacterial colony in a
102 nutrient broth at 37°C for 2– 3 hours [15].

103 ***2.3 Antibiotic susceptibility testing***

104 Antimicrobial susceptibility study was performed by Kirby–Bauer disk diffusion method
105 as recommended by the Clinical Laboratory Standards Institute (CLSI) guidelines [16]. In
106 brief, an inoculum of 0.5 McFarland standards turbidity was prepared in a Mueller-Hinton
107 broth (MHB, Hi-Media, Mumbai, India) from the isolated colony of pathogens selected
108 from 18–24 hour agar plates. A sterile cotton swab was dipped into the inoculum and
109 streaked many times on the dried surface of a Mueller-Hinton agar (MHA) plate. After 5
110 minutes, antibiotic discs were applied and pressed down to check absolute contact with
111 agar surface. The discs were apportioned in a minimum distance of 24 mm from the
112 centre. The plates were then incubated for 16-18 hrs aerobically at 37° C. The discs of
113 meropenem (10 µg), imipenem-cilastatin (20 µg) and piperacillin-tazobactam (110 µg)
114 were obtained from Microexpress Goa, India and CSE-1034 (45 µg) was obtained from
115 third party.

116 For the sensitivity of Imipenem-Cilastatin combination, we refer to the zone diameter
117 chart given for Imipenem in CLSI guidelines.

118 Breakpoints for CSE-1034: Enterobacteriaceae; >23mm - S, 20–22-I, and ≤19-R and
119 Gram-negative bacilli; >21 mm - S, 14–20-I, and ≤13-R.

120 **3. Results**

121 A total of 347 clinical specimens were obtained from the suspected patients out of
122 which 213 (61.38%) clinical samples tested positive for Gram-negative pathogens. Out of
123 these 213 Gram-negative isolates, the maximum isolates were obtained from urine

124 specimens (36.62%) followed by respiratory specimens (13.62%), blood (11.74%), pus
125 (10.33%) and wound (6.10%) while all other samples contributed a total of 6.58% (Table
126 2).

127 On the basis of morphological and biochemical screening, eight bacterial species were
128 obtained including *E. coli*, *K. pneumoniae*, *P. aeruginosa*, *A. baumannii*, along with other
129 less prevalent Gram-negative bacilli such as *Proteus spp.*, *Salmonella spp.*, *Serratia spp.*
130 and *Enterobacter spp.* which contributed 8.45% (% cumulatively) to the total clinical
131 isolates. The detailed profile of various pathogens isolated from clinical specimens is
132 shown in Table 3.

133 Table 2 represents the prevalence of different clinical isolates in different samples.
134 Data revealed the maximum prevalence of *E. coli* in urine samples, pus and stool samples.
135 *K. pneumoniae* was mostly isolated from blood and respiratory specimens whereas
136 *P. aeruginosa* isolates were mostly retrieved from the wound and respiratory specimens.
137 *A. baumannii* was least prevalent in all the specimens.

138 Susceptibility profile of pathogens isolated from clinical specimens is presented in
139 Table 4. Overall, 85.4% (182) of the total number of isolates were reported susceptible to
140 CSE-1034, 59.6% (127) to Pip-taz, 66.2% (141) to Meropenem and 64.8% (138) to
141 Imipenem. The susceptibility rates of CSE-1034 were *E. coli* (91.9%), *K. pneumoniae*
142 (77.8%), *A. baumannii* (82.4%) and *P. aeruginosa* (82.3%). Among all the antibiotics
143 tested, the least susceptibility was reported to Pip/Taz. *E. coli* exhibited the highest
144 susceptibility (80%) to Pip-taz whereas the lowest was reported by *P. aeruginosa*
145 (26.47%). Among Carbapenems, the almost similar activity of meropenem and imipenem-
146 cilastin was reported against *E. coli* (77-82%) and *K. pneumoniae* (48-50%). The
147 meropenem was marginally better than imipenem-cilastin against *A. baumannii* (58.8% vs

148 52.9%) whereas was reported significantly better than imipenem-cilastin against *P.*
149 *aeruginosa* (58.8% vs 32.3%).

150

151

152 **4. Discussion**

153 The predominant species isolated was *E. coli* (42.2%) followed by *K. pneumoniae*
154 (25.3%). A good number of studies have reported *E. coli* and *K. pneumoniae* as the most
155 common and opportunistic clinical pathogens [17, 18]. Similar results with a high
156 prevalence of *E. coli* (54.9%) were reported by Sikka *et al.* [19]. Sachdeva [20] has also
157 reported the prevalence of *E. coli* to a tune of 51.7%. A similar prevalence of *K.*
158 *pneumoniae* has been reported by Makkar *et al.* [21] who demonstrated 22% of *K.*
159 *pneumoniae* from clinical isolates. Sahu *et al.* [22] reported the prevalence of *K.*
160 *pneumoniae* to a tune of 32% which sustains our data. *Pseudomonas spp.* (15.9%) also
161 contributed significantly to the isolated pool of pathogens. As compared to other studies,
162 less number of *A. baumannii* isolates were identified in this study. The similar prevalence
163 pattern of *Proteus spp.*, *Salmonella spp.*, *Serratia spp.* and *Enterobacter spp.* is also
164 reported by many other studies. [17, 24, 25]

165 Similar to our observations, Ruppe *et al.* [14] have also reported 90% prevalence of *E.*
166 *coli* in stool samples. Majority of *E. coli* (54%) isolates were recovered from urine during the
167 study performed by Kumar *et al.* [17]. Ibrahim *et al.* [26] have also reported 40-50%
168 prevalence of *E. coli* in pus samples. *K. pneumoniae* isolates were mostly isolated from
169 blood and respiratory specimens whereas *P. aeruginosa* was mostly isolated from the wound
170 and respiratory specimens.

171 Among all the antibiotics tested, the least susceptibility was reported to Pip/Taz and
172 highest was reported towards CSE-1034. A high rate of resistance observed to Pip/taz which

173 is normally recommended the second line of treatment in our hospital could be possibly the
174 indiscriminate consumption of pip-taz. The AMR surveillance study conducted in India has
175 shown resistance against pip-taz has risen to 65-70% [19]. Among carbapenems, the average
176 susceptibility rates were 65% against all the pathogen isolates. The emergence of
177 carbapenem-resistant strains, which ranges from 18-68% in different isolates is a matter of
178 big concern as carbapenems are considered as the last resort drugs for MDR bacterial
179 infections. Singh *et al.* [11] have reported that MBLs to a tune of 15-22% among the Gram-
180 negative isolates in their study.

181 The high rate of carbapenem resistant strains reported in this surveillance study is a
182 matter of grave concern and needs to be addressed on priority at the global level. One of the
183 approaches that the clinicians have adopted to reduce selective pressure on last resort drugs is
184 by pumping in the use the antibiotic resistance breakers “ARBs” along with antibiotics to
185 revive them for clinical purposes. CSE-1034 is one such combination of beta-lactam/beta-
186 lactamase inhibitor (BL/BLI) combination with ARB “EDTA”. Interestingly, a significant
187 number of isolates were sensitive to CSE-1034 i.e., *E. coli* (98.8%), *K. pneumoniae* (90.5%),
188 *P. aeruginosa* (89.9%) and *Acinetobacter spp* (81%). Surprisingly, 131 isolates reported as
189 Meropenem resistant were susceptible to CSE-1034 (Table 4). The higher susceptibility of
190 Gram-negative pathogens to CSE-1034 has been reported by several other studies also. CSE-
191 1034 is a novel combination of Ceftriaxone, Sulbactam and disodium edetate and the high
192 susceptibility of CSE-1034 could be attributed to the synergistic effect of Ceftriaxone,
193 disodium edetate and Sulbactam. The non-antibiotic adjuvant, EDTA mediates various
194 antimicrobial effects by enhancing the penetration of antibiotic into cell membrane, decreases
195 over-expression of efflux pumps, bio-film eradication, de-activates carbapenemases-MBL by
196 chelating Zinc ions.

197 About last line therapy agents for MDR infections, our study has shown Carbapenems
198 as the most active agent only against *E.coli* (82%). Around 36-45% of *P. aeruginosa* and 45%
199 of *Acinetobacter* spp. were Carbapenem resistant. Resistance to meropenem was found
200 highest in *Klebsiella* spp. (54%). Chauhan *et al.* [19] have reported a Carbapenem resistance
201 of 14.6% in *E. coli* and 29.6% in *Klebsiella* spp. in hospital isolates from various in and
202 outpatient areas. Gupta *et al.* [21] have reported a Carbapenem resistance ranging from 17-
203 22% in different strains of Enterobacteriaceae from North India.

204 Based on pathogen type, *E. coli* exhibited the highest susceptibility rate whereas the
205 lowest was reported against *P. aeruginosa*. *E. coli* was found to be the most susceptible
206 clinical isolate among major pathogens which displayed 80%, 75.5% and 82.2% sensitivity
207 against piperacillin-tazobactam, meropenem and imipenem-cilastatin respectively.
208 Correspondent results were observed by many authors who reported 100%, 90% and 96.5%
209 sensitivity of *E. coli* against meropenem, piperacillin-tazobactam and imipenem-cilastatin
210 respectively [35,36]. *Klebsiella* spp. exhibited intermediate susceptibility i.e. 44.4%, 50% and
211 48.1% towards piperacillin-tazobactam, meropenem and imipenem-cilastatin. Similar results
212 were noted by many authors who revealed 40-60% sensitivity of *Klebsiella* spp. against
213 piperacillin-tazobactam, meropenem and imipenem-cilastatin [37,38]. As reported earlier
214 also, *Acinetobacter* spp. experienced highest susceptibility (96.3%) towards CSE-1034CSE-
215 1034 only while extreme resistance (96.3% each) against rest of the antibiotics which is due
216 to sulbactam (a β -lactamase inhibitor) which owns intrinsic whole-cell activity against
217 *Acinetobacter* spp. [39]. Surprisingly, contrary to expectations, *Pseudomonas* spp.
218 documented 73.5%, 73.5% and 67.6% resistance against piperacillin-tazobactam, meropenem
219 and imipenem-cilastatin respectively. Mohammadi and Feizabadi [37] reported >60%
220 resistance of piperacillin+tazobactam against gram negative bacilli isolated from clinical
221 samples which supports our data. Similarly, Hout *et al.* [40] revealed 70-100% resistance of

222 meropenem towards *Acinetobacter spp.* and *Pseudomonas spp.*. Likewise, Shour and El-
223 Sharif, [36] and Eldomany and Abdelaziz noticed significant resistance (>50%) of
224 imipenem-cilastatin in *Acinetobacter spp.* and *Pseudomonas spp.* which is in accordance with
225 our present data [41].

226 The emergence of antimicrobial resistance against BL-BLI and carbapenem drugs is due to
227 numerous elements which assists the scattering of resistance among clinical pathogens which
228 includes the production of MBL enzymes, biofilm formation, over expression of efflux
229 pumps and accumulation of the drug [42,43]. None of these mechanisms is dressed by either
230 piperacillin-tazobactam, meropenem or imipenem-cilastatin and probably this could be one
231 reason for CSE-1034 super performance that it is supplemented with EDTA as ARB. The
232 progressive and relentless resistance towards BL-BLI and carbapenem antibiotics is probably
233 the result of overuse of antibiotics, improper processing and inappropriate prescribing [44]. In
234 the light of above discussion, it is evident that Antibiotic adjuvant therapy which has ARB
235 can be used as the prime choice of therapeutics to overcome the resistance raised among
236 gram negative pathogens towards β -lactam and β -lactamase inhibitor combinations and
237 penems in the treatment of bacterial infectious diseases.

238 **5. Conclusion**

239 This retrospective study indicates the rise in resistance among the most prevalent and
240 opportunistic gram negative pathogens against β -lactam and β -lactamase inhibitor
241 combinations and penems. Present data strongly advocates precedence of CSE-1034 over β -
242 lactam and β -lactamase inhibitor combinations and penems as CSE-1034 has scored 85-100%
243 susceptibility which excels the antimicrobial activity of rest of the drugs. Therefore, CSE-
244 1034, a novel product with antibiotic resistance breaker can be used as an empirical and
245 alternate choice of the drug over potent therapeutics in encountering multidrug resistance
246 among healthcare-associated pathogens.

247 **Ethical Approval:**

248

249 As per international standard or university standard was written ethical approval has been collected
250 and preserved by the author(s).

251

252

253

254

255 **References**

256 1. Gelband H, Miller-Petrie M, Pant S, Gandra S, Levinson J, Barter D, et al. The State
257 Of The World's Antibiotics 2015. Center For Disease Dynamics, Economics & Policy
258 Washington DC and New Delhi. https://cddep.org/sites/default/files/swa_2015_final.pdf

259 2. Neill JO. Review on antibiotic resistance. Tackling drug resistant infections globally.
260 2016.

261 3. Peleg AY, Hooper DC. Hospital-Acquired Infections Due to Gram-Negative Bacteria.
262 N Engl J Med. 2010;362(19):1804-13.

263 4. Zalewska-Piatek BM. Urinary tract infections of Escherichia coli strains of the
264 chaperone-usher system. Pol J Microbiol. 2011;60(4):279-85.

265 5. Neill JO. Review on antibiotic resistance. Tackling drug resistant infections globally.
266 Tackling a crisis for the health and wealth of nation. 2014.

267 6. Sethuraman S, Arunachalam A, Karthikeyan M, Kumar SA, Manidipa S, Senthilraj R.
268 Antimicrobial Sensitivity Profile of Serratia Marcescens strain isolated in

- 269 Government General hospital, Nagapattinam, Tamilnadu, India. Int J Preclin Pharm
270 Res. 2011;’2(1):7-11.
- 271 7. Meeta S, Sati P, Preeti S. Prevalence and antibiogram of extended spectrum β -
272 lactamase (ESBL) producing Gram-negative bacilli and further molecular
273 characterization of ESBL producing *Escherichia coli* and *Klebsiella* spp. J Clin Diag
274 Res. 2013;7(10):2168–2172.
- 275 8. Singh N, Pattnaik D, Neogi DK, Jena J, Mallick B. Prevalence of ESBL in
276 *Escherichia coli* Isolates Among ICU Patients in a Tertiary Care Hospital. J Clin
277 Diagn Res. 2016;10(9):DC19–DC22.
- 278 9. Bora A, Sanjana R, Jha BK, Mahaseth SN, Pokharel K. Incidence of metallo-beta-
279 lactamase producing clinical isolates of *Escherichia coli* and *Klebsiella pneumoniae*.
280 BMC Res Notes. 2014;7:557.
- 281 10. Kontopidou F, Plachouras D, Papadomichelakis E, Koukos G, Galani I, Poulakou G,
282 et al. Colonization and infection by colistin-resistant Gram-negative bacteria in a
283 cohort of critically ill patients. Clin Microbiol Infect. 2011;17(11):E9–E11.
- 284 11. Parveen RM, Khan MA, Menezes GA, Harish BN, Parija SC, Hays JP. Extended-
285 spectrum β -lactamase producing *Klebsiella pneumoniae* from blood culture in
286 Puducherry, India. Indian J Med Res. 2011;134(3):392-395.
- 287 12. Datta S, Wattal C, Goel N, Oberoi JK, Raveendran R, Prasad KJ. A ten year analysis
288 of multi-drug resistant bloodstream infections caused by *Escherichia coli* &
289 *Klebsiella pneumoniae* in a tertiary care hospital. Indian J Med Res.
290 2012;135(6):907–12.
- 291 13. Center for Disease Dynamics, Economics & Policy (CDDEP). 2015b. ResistanceMap.
292 Accessed on August 20, 2015.

- 293 14. Ruppe E, Woerther P-L, Barbier F. Mechanisms of antimicrobial resistance in gram-
294 negative bacilli *Ann Intensive Care*. 2015;5:21.
- 295 15. Cheesbrough M. Biochemical tests to identify bacteria. In: *District laboratory practice*
296 *in tropical countries*. 2nd edition. Cambridge University Press, UK; 2000: 178-187.
- 297 16. Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial*
298 *Susceptibility Testing; Twenty-sixth Informational Supplement*. Wayne, PA 19087
299 USA: CLSI Document M100-S26; 2016.
- 300 17. Kumar M, Chaudhary S, Makkar DJ, Garg N, Chugh S. Comparative antimicrobial
301 efficacy evaluation of a new product CSE-1034 against Meropenem of gram negative
302 isolates. *Asian J Pharm Clin Res*. 2015;8(4):251-54.
- 303 18. Akter J, Chowdhury AMMA, Al Forkan M. Study on Prevalence and Antibiotic
304 Resistance Pattern of Klebsiella Isolated from Clinical Samples in South East Region
305 of Bangladesh. *American Journal of Drug Discovery and Development*. 2014;4:73-79.
- 306 19. Sikka R, Mann JK, Deep, Vashist MG, Chaudhary U, Deep A. Prevalence and
307 antibiotic sensitivity pattern of bacteria isolated from nosocomial infections in a
308 surgical ward. *Indian J Clin Pract*. 2012;22:519-25.
- 309 20. Sachdeva N. Antibiotic sensitivity pattern of bacterial pathogens in Rajeev Gandhi
310 Cancer Hospital, Delhi. *Int J Rec Sci Res*. 2016;7(1):8480-8485.
- 311 21. Makkar DK, Kumar M, Chaudhary S, Goyal S, Aggarwal P, Garg N. Prevalence and
312 Susceptibility Analysis of Gram negative Pathogens. *J Pharm Bio Sci*. 2015;10(3):58-
313 65.
- 314 22. Sahu M, Sanjith S, Bhalekar P, Keny D. *Waging War Against Extended Spectrum*

- 315 Beta Lactamase and Metallobetalactamase Producing Pathogens- Novel Adjuvant
316 Antimicrobial Agent Cse1034- An Extended Hope. J Clin Diagn Res.
317 2014;8(6):DC20-DC23.
- 318 23. Shaikh S, Fatima J, Shakil S, Mohd S, Rizvi D, Kamal RA. Prevalence of multidrug
319 resistant and extended spectrum beta- lactamase producing *Pseudomonas aeruginosa*
320 in a tertiary care hospital. Saudi J Biol Sci. 2015;22:62-64.
- 321 24. Ejaz H, Zafar A, Anwar N, Cheema TA, Shahzad H. Prevalance of bacteria in urinary
322 tract infections among children. Biomedica. 2006:22.
- 323 25. Mohammed MA, Alnour TMS, Shakurfo OM, Aburass MM. Prevalence and
324 antimicrobial resistance pattern of bacterial strains isolated from patients with urinary
325 tract infection in Messalata Central Hospital, Libya. Asian Pac J Trop Med.
326 2016;9(8):771-76.
- 327 26. Ibrahim ME, Bilal NE, Hamid ME. Increased multi-drug resistant *Escherichia coli*
328 from hospitals in Khartoum state, Sudan. African Heal Sci. 2012;12(3):368-75.
- 329 27. Sharma S, Srivastava P, Kumar A. Respiratory Tract Infection (RTI) and its
330 Treatment. Sch J App Med Sci. 2015;3(8E):3118-3122.
- 331 28. Subhedar V, Bandoowala S, Jain SK. Prevalence of multi drug resistant *Klebsiella*
332 *pneumoniae* infections, causing a HAVOC in intensive care units of a tertiary care
333 hospital. Sch J App Med Sci. 2016;4(11):1054-1059.
- 334 29. Nahar A, Anwar S, Saleh AA, Miah MRA. Isolation of *Acinetobacter* species and
335 their antimicrobial resistance pattern in an Intensive care unit (ICU) of a tertiary care
336 hospital in Dhaka, Bangladesh. Bangladesh J Med Microbiol. 2012; 06(01):03-06.
- 337 30. Gupta N, Gandham N, Jadhav S, Mishra RN. Isolation and identification of
338 *Acinetobacter* species with special reference to antibiotic resistance. J Nat Sci Biol
339 Med. 2015;6(1):159–62.

- 340 31. Patel H, Garala RN. Antibiotic susceptibility pattern of *Pseudomonas aeruginosa*
341 isolated at SSG hospital Baroda. *J Res Med Dent Sci*. 2014;2(1):84-87.
- 342 32. Senthamarai S, Reddy SK, Sivasankari S, Anitha C, Somasunder V, Kumudhavethi
343 MS et al. Resistance Pattern of *Pseudomonas aeruginosa* in a Tertiary Care Hospital
344 of Kanchipuram, Tamilnadu, India. *J Clin Diagn Res*. 2014;8(5):DC30–DC32.
- 345 33. Prasad RR, Shree V, Sagar S, Kumar S, Kumar P. Prevalence and Antimicrobial
346 Susceptibility Pattern of *Proteus* Species in Clinical Samples. *Int J Curr Microbiol*
347 *App Sci*. 2016;5(4):962-968.
- 348 34. Menon D. A case report of community acquired Pneumonia due to multidrug
349 resistance *Pseudomonas aeruginosa* treated with Eloxeres. *Int J Med Pharm case reports*.
350 2016;6(1):1-5.
- 351 35. Guillou M-L, Kempf M, Cavallo J-D, Chomarar M, Dubreuil L, Maugein J, et al.
352 Comparative *in vitro* activity of Meropenem, Imipenem and Piperacillin/tazobactam
353 against 1071 clinical isolates using 2 different methods: a French multicentre study.
354 *BMC Infect Dis*. 2010;10:72 (1-9).
- 355 36. Abuhandan M, Guzel M, Oymak Y, Ciftci H. Antibiotic sensitivity and resistance in
356 children with urinary tract infection in Sanliurfa. *Turk J Urol*. 2013;39(2):106–10.
- 357 37. Mohammadi MM, Feizabadi MM. Antimicrobial resistance pattern of Gram-negative
358 bacilli isolated from patients at ICUs of Army hospitals in Iran. *Iranian J Microbiol*.
359 2011;3(1):26-30.
- 360 38. Eldomany R, Abdelaziz NA. Characterization and antimicrobial susceptibility of gram
361 negative bacteria isolated from cancer patients on chemotherapy in Egypt. *Arch Clin*
362 *Microbiol*. 2011; 2(6):2.
- 363 39. Penwell WF, Shapiro AB, Giacobbe RA, Gu RF, Gao N, Thresher J et al. Molecular

- 364 Mechanisms of Sulbactam Antibacterial Activity and Resistance Determinants in
 365 *Acinetobacter baumannii*. Antimicrob Agents Chemother. 2015;59(3):1680–1689.
- 366 40. Hout B, Oum C, Men P, Vanny V, Supaprom C, Heang V et al. Drug resistance in
 367 bacteria isolated from patients presenting with wounds at a non-profit Surgical Center
 368 in Phnom Penh, Cambodia from 2011–2013. Trop Dis, Travel Med Vacc. 2015;1(4):1-
 369 11.
- 370 41. Shour HM, EL-Sharif A. Species distribution and antimicrobial susceptibility of gram
 371 negative aerobic bacteria in hospitalized cancer patients. J Transl Med. 2009;7:14.
- 372 42. Fair RJ, Tor Y. Antibiotics and Bacterial Resistance in the 21st Century. Perspect
 373 Medicin Chem. 2014;6:25–64.
- 374 43. Singh J, Shukla SK. A New Threat of Bacterial Resistance towards Life Saving
 375 Carbapenem Antibiotics. Res J Chem Sci. 2015;5(4):85-88.
- 376 44. Ventola CL. The Antibiotic Resistance Crisis Part 1: Causes and Threats. Pharm
 377 Therapeut. 2015;40(4):277-83.

378

379 **Table 1: Selective culture medium used for isolation of different pathogens.**

Pathogen	Selective media
<i>Klebsiella spp.</i>	Hicrome Klebsiella selective agar base medium
<i>E. coli</i>	Eosine Methylene Blue (EMB) agar medium
<i>Acinetobacter spp.</i>	Leeds acinetobacter agar base medium
<i>Pseudomonas spp.</i>	Citrimide agar medium
<i>Proteus spp.</i>	EMB agar medium
<i>Salmonella spp.</i>	Wilson and Blair bismuth sulphite medium
<i>Serratia spp.</i>	Caprylate-thallos agar medium
<i>Enterobacter spp.</i>	EMB agar medium

380

381

382
383
384
385
386
387
388
389
390

Table 2: A profile of clinical samples used as a source of the pathogenic isolates.

Sr. No.	Name of clinical samples	Total no. of samples collected	Number of samples showing growth of pathogens (%)	Number of samples not showing growth of pathogens
1	Urine	94	78 (36.62%)	16
2	Respiratory specimens	84	51	
3	Blood	39	25 (11.74%)	14
4	Pus	34	22 (10.33%)	12
5	Wound	22	13 (6.10%)	9
6	Stool	19	10 (4.69%)	9
7	Other samples	55	14 (6.58%)	41
	Total	347	213 (61.38%)	134

391
392
393
394
395
396
397
398
399
400

Table 3: Prevalence of different clinical isolates in different samples.

	Clinical Isolates					
Samples	No. of isolates	<i>E. coli</i> (%)	<i>Klebsiella spp.</i> (%)	<i>Acinetobacter baumannii</i> (%)	<i>Pseudomonas aeruginosa</i> (%)	Other pathogens (%)
Urine	78	59 (75.64)	11 (14.10)	0	7 (8.97)	1 (1.28)
Respiratory specimens	51	6 (3.45)	17 (31.03)	10 (44.83)	12 (23.52)	6 (11.76)
Blood	25	2 (8)	10 (40)	4 (36)	5 (20)	4 (16)
Pus	22	10 (45.45)	6 (27.27)	2 (9.09)	2 (9.1)	2 (9.1)
Wound	13	2 (15.38)	3 (23.08)	1 (7.69)	5 (38.46)	2 (15.38)
Stool	10	8 (80)	0	0	0	2 (20)
Other samples	14	3 (21.43)	7 (50.0)	0	3 (21.43)	1 (7.14)
Total	213	90	54	17	34	16
Total (%)		42.25 %	25.35 %	7.98 %	15.96 %	8.45%

401

402

403

404

Table 4: Susceptibility pattern of clinical isolates.

405

Susceptibility (%)									
Clinical isolates	No. of isolates	Antibiotic adjuvant entity		BL-BLI		Carbapenem			
		CSE-1034		Piperacillin-tazobactam		Meropenem		Imipenem-cilastatin	
		S	R	S	R	S	R	S	R
<i>E. coli</i>	90	91.1 (82)	8.9 (8)	(80) 72	(20) 18	77.7 70	22.3 22	82.22 74	17.78 16
<i>Klebsiella spp.</i>	54	77.8 (42)	22.2 (12)	(44.44) 24	55.56 30	50 27	50 27	48.15 26	51.85 28

<i>Acinetobacter baumannii</i>	17	82.4 (14)	17.6 (3)	35.3 6	64.7 11	58.8 10	41.2 7	52.9 9	47.1 8
<i>Pseudomonas aeruginosa</i>	34	82.3 (28)	17.7 (6)	26.47 9	73.53 25	58.8 20	41.2 14	32.35 11	67.65 23
Other pathogens	18	(88.9) 16	(9.91) 2	88.9 16	9.91 2	88.9 16	9.91 2	100 18	0
Total	213	(85.4) 182	(14.5%) 31	(59.6) 127	(40.4) 86	(66.2) 141	(33.8) 72	(64.8) 138	(35.2) 75

406