

1 The effect of solid (granular) and liquid (foliar) fertilizers application on the growth and yield of
2 maize (*Zea mays L*) in soils of Obubra, Cross River State, Nigeria

3

4 **ABSTRACT**

5 The research was carried out in the month of April, 2018 to determine the effect of solid
6 (granular) and liquid (foliar) fertilizers application on the growth and yield of maize in soils of
7 Obubra. Five (5) composite soil samples were collected at the depth of 0-20 cm for the analysis
8 of physical and chemical properties before application of the fertilizers. The experimental layout
9 was randomized Complete Block Design (RCBD) with three treatments and five replications in a
10 plot area of 25 m x 20 m (500 m²) which corresponds to 0.05 ha⁻¹. The plot was designed and
11 blocked into subplots, each measuring 6 m x 4 m (24 m²). Each block was separated from the
12 other with a distance of one meter (1 m) apart and between subplots 0.5 m apart. Three
13 treatments made up of treatment one (T₁) zero application at 0 kg ha⁻¹, treatment two (T₂) liquid
14 (foliar) N.P.K 20:20:20 at the rate of 100 ml of N, 50 ml of P₂O₅, 33.3 ml of Mp ha⁻¹ and
15 treatment three (T₃) solid (granular) N.P.K 20:20:20 at the rate of 44.4 kg of N, 40 kg of P₂O₅
16 and 33.3 kg of Mp ha⁻¹ were replicated five times making a total of fifteen (15) subplots.
17 Parameters of plant heights and number of leaves were observed at 6 and 8 weeks after planting.
18 Plant heights, number of leaves, number of cobs, weight of 1000 seeds in each subplot and
19 weight of grain after shelling were analyzed respectively. Results on soil analyses showed that
20 the soil texture was sandy loam with deficiencies in primary nutrients and other nutrients. On the
21 plant heights, the result was significant (P<0.05) and on the number of leaves, the result for 6
22 weeks was not significant (P>0.05) while that of 8 weeks was significant (P<0.05). On the
23 number of cobs, 1000 seeds and weight of grain after shelling were also significant (P<0.05).
24 The solid (granular) fertilizer showed to be more effective than liquid (foliar) fertilizer and
25 should therefore be recommended for the growth and yield of maize in the area.

26 **Keywords: Maize, Growth and yield, Fertilizer**

27

28

29

30 INTRODUCTION

31 Maize, other names corn, Indian corn, mealis (English), mais (French), milho
32 (Portuguese), maize (Spanish), Dura ash shahami (Opabic), makai, butta (Hindi) belongs to the
33 family *poaceae*. Tribe - *maydeae*, Genus – *Zea* and Specie - *mays*. However, there are a number
34 of theories regarding the origin of maize but it seems most probable that it originated in Mexico
35 or Central America [6] where it has been in cultivation for more than 700 years [16]. Maize was
36 brought to Europe by Columbus and was introduced into Africa by the Portuguese. Maize today
37 is probably the next most important grain cereal after wheat in the world [15]. It is now found all
38 over the world and its natural habitat is the tropics. In Nigeria, Maize is one of the major staple
39 foods, fodder and industrial crop for commercial and subsistence level where it is grown in all
40 agro ecological zones [14; 7]. Maize is predominantly the cereal crop of Southern Nigeria, just as
41 sorghum and millet are in the Northern Nigeria [14]. The crop to some extent is cultivated
42 practically throughout the country. Maize is one of the oldest and widely cultivated World' s
43 cereals and strong annual crop/grass, usually producing one stem and growing to a height of 1-
44 4.5m. Older/local varieties of the crop mature after 100 – 120 days but more rapidly maturing
45 varieties are now available. The male inflorescence called a tassel for hybrid varieties is produce
46 after 50 – 60 days as a continuation of the main stem. The female inflorescence, called the ear or
47 cob is a modified spike formed on a short branch in the axils of the largest foliage leaves.

48 Maize grows in a wide range of zones with altitude ranging from 100- 2900 m above sea
49 level and evenly distributed rainfall of 400 – 1200 mm during the growing period. Require well
50 drained light loam or silty loams of alluvial soils with a pH of 5.5 -7.0 and warm temperatures
51 between 15-30⁰C for good yields. It does not tolerate water logging. Cold conditions extends the
52 maturity period whereas high temperatures lower the yields. Its ability to thrive under different

53 ecological conditions in Nigeria has led to increased production. The major pests of maize are
54 fall army worm stalk borer, maize aphids and cutworms [22]. Major diseases include Maize
55 Lethal Necrosis Disease (MLND), Maize smut and Northern leaf blight [23].

56 To obtain maximum growth and yield of maize, many research findings have shown that
57 neither organic nor inorganic fertilizers alone can result in sustainable productivity [19]. Liquid
58 (foliar) fertilizer is a form of fertilizer obtained by dissolving NPK 20:20:20 or NPK 15:15:15 in
59 water to form soluble substance [12]. This dissolution can be made in a can bottle or any
60 container. The fertilizer is spread to the leaf of the plant where quantities of the major plants food
61 can be absorbed through the leaf at one time. The solid (granular) fertilizers have different
62 nutrient elements required by plants in its composition, but the most essential ones are nitrogen,
63 phosphorus and potassium. The nitrogen contain 1- 5 % weight by plant and exist as nitrate
64 (NO_3^-), ammonium ion (NH_4^+) and urea ($\text{Co}(\text{NH}_2)_2$). The nitrate form dominates in moist warm
65 and aerated soils and it is the preferred form of nitrogen in plants. The phosphorus varies in
66 concentration from 0.1 – 0.4 % in plant and available as phosphate ion (H_2PO_4^-), orthophosphate
67 (HPO_4^{2-}). Here the phosphate ion dominates in soil with optimal pH values. Other forms like
68 phosphate are component of fertilizers and form orthophosphate during hydration. These
69 phosphate ions are involved in the major soil chemical reactions and numerous metabolic
70 pathways in plant nutrition with the most essential being the storage and transfer of energy [9].

71 The soils of the Tropical Rain Forest are heavily leached of plant nutrients due heavy
72 rainfall in the area [8]. The soils of Obubra belong to the soil order, Ultisols which are
73 extensively weathered [2]. The soils are highly leached and therefore acidic in reaction probably
74 due to high amounts of rainfall in the area [6]. Their major constraints include the sandy nature
75 of the surface, prone to severe and internal erosion, low potassium reserve and high acidity thus

76 necessitating regular liming [20; 10]. The soils are generally suitable for most arable crops and
77 cash crops [10]. Therefore, the objective of this study was to investigate the comparison of the
78 effect of solid (granular) and liquid (foliar) fertilizers application on the growth and yield of
79 maize.

80

81 **MATERIALS AND METHODS**

82 **Study area**

83 This research was carried out in the month of April, 2018 in sandy loam texture soils
84 belonging to the soil order, Ultisols [2] at the Cross River University of Technology teaching and
85 research farm, Ovonum, Obubra Local Government Area of Cross River State, Nigeria. Obubra
86 lies between Latitude $06^{\circ} 5' 8.466''$ N and $08^{\circ} 32' 80''$ E. The rainfall distribution is between 2250
87 – 2500 mm, a mean annual rainfall of 2375 mm [2]. The average temperature and humidity are
88 25°C and 70 %. The study area has a number of economic activities such as lumbering, fishing
89 and craftsmanship. The area witnesses two major seasons usually characterized by heavy and
90 incessant showers.

91

92 **Experimental site, Procedures, Treatments and Experimental design**

93 The research site is about 0.5 km away from the University Lecture Halls. Before the
94 design of the experimental plot, composite soil samples were collected in a random manner using
95 soil auger and analyzed for physical and chemical properties of the soil. Thereafter, the
96 experimental plot was cleared during the month of March in the 2018 farming season. The plot
97 was designed and blocked into subplots, each measuring 6 m x 4 m (24 m^2). Fifteen subplots
98 containing eight ridges each were constructed making a total of one hundred and twenty ridges in
99 the experimental plot. The total area of the experimental plot was 25 m x 20 m which gives a

100 total of 500 m² (0.05 ha⁻¹). Each block was separated from the other with a distance of 1 m apart
 101 and between subplots 0.5 m apart. The experimental design used was Randomized Complete
 102 Block Design (RCBD) with five replications in three (3) treatments namely treatment one (T1) -
 103 Zero application at the rate of 0 kg ha⁻¹ treatment two (T2) – liquid fertilizer (NPK 20:20:20) at
 104 the rate of 100 mil of N, 50 mil of P₂O₅, 33.3 mil of Mp ha⁻¹ and treatment three (T3) – Solid
 105 fertilizer (NPK 20:20:20) at the rate of 44.4 kg of N, 40 kg of P₂O₅ and 33.3 kg of Mp ha⁻¹.

106

107 Table 1: Fertilizer type, rate applied on each plot and hectare

108

| TRTS. | Treatment and Replication | Fertilizer type | Rate applied/each plot | Rate applied/ha |
|-------|---|-----------------|---|--|
| T1 | T ₁ R ₁ , T ₁ R ₂ , T ₁ R ₃ , T ₁ R ₄ & T ₁ R ₅ | Nothingness | Nothingness | Nothingness |
| T2 | T ₂ R ₁ , T ₂ R ₂ , T ₂ R ₃ , T ₂ R ₄ & T ₂ R ₅ | NPK 20:20:20 | 0.24 mil of N, 0.12 mil of P ₂ O ₅ , 0.08 mil of Mp | 100 mil of N, 50 mil of P ₂ O ₅ , 33.3 mil of Mp |
| T3 | T ₃ R ₁ , T ₃ R ₂ , T ₃ R ₃ , T ₃ R ₄ & T ₃ R ₅ | NPK 20:20:20 | 0.11 kg of N, 0.10 kg of P ₂ O ₅ , 0.08 kg of Mp | 44.4 kg of N, 40 kg of P ₂ O ₅ , 33.3 kg of Mp |

109

110 **Sampling and data collection**

111 The planting material (maize seeds) was obtained from local market of Ikom, Cross River
 112 State, Nigeria. The central row plants were used for data collection where growth parameters
 113 namely plant height, number of leaf per plant with yield components such as number of cobs;

114 number of seeds per row and weight of grain after shelling were recorded. Plant height (cm) was
115 measured from the base of the plant to the upper of the top most leaves. The numbers of
116 functional leaves per plant was a visual count of the green leaves. The number of cobs was
117 through counting from randomly selected cobs and the grain after shelling was weighed.

118

119 **Laboratory analyses**

120 Five (5) composite soil samples were collected from the site and air-dried, gently crushed with
121 pestle and mortar and sieved through a 2.00 mm sieve to obtain the fine earth fraction for the
122 analysis. Particle size analysis was determined by Bouyoucos hydrometer methods using sodium
123 hexametaphosphate (VII) as dispersant [21]. Soil texture was determined using USDA soil
124 textural triangle [18]. Five (5) undisturbed soil samples were collected by 100 cm³ metallic cores
125 and oven-dried at 105⁰C to constant for determination of bulk density. The pH was determined
126 potentiometrically with a glass electrode pH meter in water at 1:2.5 soils: water ratio [3].
127 Organic Carbon was determined following the Walkley and Black wet oxidation method as
128 outlined by [13]. Total nitrogen was determined by the micro-kjeldhal method [21]. Available
129 phosphorus was determined by extraction with Bray P-I extractants described by [4].
130 Exchangeable acidity was determined by successive leaching of soil with neutral unbuffered 1N
131 KCl using 1:10 Soil: Liquid ratio. The amount of H⁺ and Al³⁺ in the leachate was determined by
132 the titration method [21]. Exchangeable cations were determined with 1N ammonium acetate
133 (pH 7.0) using 1:10 Soil: Water ratio. Ca⁺⁺ and Mg⁺⁺ in the filtrate were determined with an
134 atomic adsorption spectrophotometer (AAS) while Na⁺ and K⁺ were determined with a flame
135 photometeras described by [21]. Cation exchange capacity (CEC) was determined by the neutral
136 ammonium acetate (pH 7.0) method. While effective cation exchange capacity was calculated by

137 summing up exchangeable H^+ and Al^{3+} and exchangeable cations. Base saturation was
138 determined by dividing the summation of exchangeable bases (Ca^{2+} , Mg^{2+} , K^+ and Na^+) by the
139 effective cation exchange capacity and multiplies by 100. The formula is as follows:

140

$$\text{Base saturation} = \frac{\text{Total Exchangeable bases}}{\text{ECEC}} \times \frac{100}{1}$$

141

142 **Data analysis**

143 Data collected on various growths and yield parameters were subjected to analysis of
144 variance (ANOVA) in Randomized Complete Block Design (RCBD). The treatments mean were
145 separated by F-LSD test at 5% probability level using GenStat software version 8.10.

146

147 **RESULTS AND DISCUSSIONS**

Soil properties before trial of fertilizer

The result on the soil physical and chemical properties before trial of fertilizer is shown in Table 2. The results showed that the soil had a sandy loam texture with high proportion of sand content and deficient in nutrients. The soil pH (5.4) showed very strongly acidic milieu [11]. The organic carbon, total N, and available phosphorus were low. The low contents in organic carbon, total N and available P could be attributed to the effects of intensive cultivation of the soils in the area. This conforms to the work of [17]; [12] who stated that continuous cultivation of land results in the reduction of soil nutrients especially organic carbon. The low content of available P might be attributed to the pH. The exchangeable bases were also low. This might be attributed to high rainfall in the areas which leaches the basic cations down the profile. The exchangeable bases were generally low with no Mg^{2+} . This is an

indication of how the cations are leached by rain. The exchangeable acidity was high indicating the acidic condition of the soil. The soil requires fertility management practices.

148 Table 2: Results on soil properties before application of fertilizer

| Physico-chemical properties | Quantity |
|---------------------------------------|-------------------|
| Sand (%) | 74.0 |
| Silt (%) | 16.0 |
| Clay (%) | 10.0 |
| pH (H ₂ O) | 5.4 |
| Org. Carbon (%) | 1.13 |
| Total nitrogen (%) | 0.14 |
| Av. P (mg/kg) | 15.63 |
| Exchangeable cations (cmol/kg) | |
| Ca ²⁺ | 2.4 |
| Mg ²⁺ | 0 |
| K ⁺ | 0.09 |
| Na ⁺ | 0.07 |
| Exchangeable acidity (cmol/kg) | |
| Al ³⁺ | 1.32 |
| H ⁺ | 2.36 |
| ECEC | 6.64 |
| B.S (%) | 44.58 |
| Textural Class | Sandy loam |

149

150

151

152 **Plant height (cm)**

153 The plant heights were measured in centimeters (cm) in each subplot. The results are
 154 presented in Table 3. The results analyzed for the 6 and 8 weeks after planting were highly
 155 significantly ($P \leq 0.05$) difference. Treatment three (T₃) recorded the highest plant height, mean
 156 values of 57.94 and 64.02 followed by treatment two (T₂) which recorded mean values of 52.24
 157 and 58.24, followed by treatment one (T₁) which recorded the least number in plant height, mean
 158 values of 41 and 53.08. Treatment three (T₃) that recorded the highest followed by treatment two

159 (T₂) could be attributed to the effect of fertilizers applied which enhanced the growth and yield.
 160 This agrees with [1] who elucidated that there was high significant difference in maize plant
 161 height in plots treated with fertilizers compared to zero application.

162

163

Table 3: Results of plant heights at 6 weeks and 8 weeks after planting

| 6WAP | | | | | | | | 8WAP | | | | | | |
|---------------------|-----------|-------|-------|-------|-------|-------|-------|---------------------|----------|-------|-------|-------|-------|-------|
| TRTS. | R1 | R2 | R3 | R4 | R5 | TOTAL | MEAN | R1 | R2 | R3 | R4 | R5 | TOTAL | MEAN |
| 1 | 40.1 | 35.6 | 40 | 43.2 | 46.1 | 205 | 41 | 55 | 50 | 56.1 | 49.8 | 54.5 | 265.4 | 53.08 |
| 2 | 50.1 | 50.2 | 53.5 | 56.2 | 51.2 | 261.2 | 52.24 | 50 | 56 | 65.2 | 59.5 | 60.5 | 291.2 | 58.24 |
| 3 | 60 | 45 | 60.1 | 60.5 | 64.1 | 289.7 | 57.94 | 64 | 61.5 | 68.5 | 69.4 | 56.7 | 320.1 | 64.02 |
| BLK Total | 150.2 | 130.8 | 153.6 | 169.9 | 161.4 | 755.9 | | 169 | 167.5 | 189.5 | 178.7 | 171.7 | 876.7 | |
| F-LSD (0.05) | ** | | | | | | | F-LSD (0.05) | * | | | | | |

164

165

166 WAP = Week after Planting R = Replication, TRTS =Treatments, * = Significant; ** = not significant

167

168 **Number of leaves**

169 Numbers of leaves per plant on 10 plants in the middle row were counted and their mean
 170 obtained for each treatment at 6 and 8 WAP. The result is presented in Table 4. The result shows
 171 that number of leaves for 6 weeks was not significant ($P \geq 0.05$) while that of 8 weeks was
 172 significant ($P \geq 0.05$) with treatment three (T₃) recording the highest number of leaves, mean
 173 values of 10.62 and 13.6 respectively, followed by treatment two (T₂) which recorded mean
 174 values of 10.12 and 12.08 and treatment (T₁) recording the least mean values of 8.6 and 10.12 at

175 both 6weeks and 8 weeks after planting. The highest number of leaves recorded in treatment
 176 three (T₃) followed by treatment two (T₂) was due to the fertilizer application which boosted the
 177 growth of vegetative part of the plant.

178

Table 4: Results of number of leaves at 6 weeks and 8 weeks after planting

| TRTS. | 6WAP | | | | | | | 8WAP | | | | | | |
|---------------------|-----------|------|------|------|------|-------|-------|-----------------------|------|------|------|------|-------|-------|
| | R1 | R2 | R3 | R4 | R5 | TOTAL | MEAN | R1 | R2 | R3 | R4 | R5 | TOTAL | MEAN |
| 1 | 9.2 | 10 | 7.5 | 8.2 | 8.1 | 43 | 8.6 | 10.5 | 9.5 | 10.6 | 10 | 10 | 50.6 | 16.12 |
| 2 | 9.7 | 9.6 | 10.6 | 10.2 | 10.1 | 50.6 | 10.12 | 13 | 11 | 13 | 12 | 11.4 | 60.4 | 12.08 |
| 3 | 11.1 | 11 | 10.2 | 9.3 | 11.5 | 53.1 | 10.62 | 14 | 13.5 | 13 | 12.5 | 15 | 68 | 13.6 |
| BLK Total | 30 | 30.6 | 28.3 | 27.7 | 30.1 | 146.7 | | 37.5 | 33.8 | 36.6 | 34.5 | 36.4 | 179 | |
| F-LSD (0.05) | ** | | | | | | | F-LSD (0.05) * | | | | | | |

179

180

181 WAP = Week after Planting R = Replication, TRTS =Treatments, * = Significant, ** = not significant

182

183 **Number of cobs (kg)**

184 The result on the number of cobs in each subplot is presented in Table 5. The result
 185 shows that treatment three (T₃) recorded the highest number of cobs, a mean value of 7.4
 186 followed by treatment two (T₂) which recorded the mean value of 6.3 and treatment one (T₁)
 187 recorded the least mean value of 5.3. There was high significant (P≤0.05) difference in the
 188 number of cobs. This could be attributed to application of fertilizer resulting to taller plant which
 189 bears more cobs. This conforms to [5] who noticed that plant height is an important parameter of
 190 yield of maize as usually taller plant bears more cobs and offers more yield.

191

192

Table 5: Results of number of cobs in each subplot

| TRTS. | R1 | R2 | R3 | R4 | R5 | TOTAL | MEAN |
|-----------|------|-----|-----|----|------|-------|------|
| 1 | 5 | 5.5 | 5.5 | 5 | 5.5 | 26.5 | 5.3 |
| 2 | 6.5 | 6.5 | 6.5 | 6 | 6 | 31.5 | 6.3 |
| 3 | 7 | 8 | 7 | 7 | 7 | 37 | 7.4 |
| BLK Total | 18.5 | 20 | 19 | 19 | 18.5 | 95 | |

193 **F-LSD (0.05) ***

194 WAP = Week after Planting R = Replication, TRTS =Treatments, * = Significant

195

196 **1000 seeds (g)**

197 The result of 1000 seeds weight in each subplot is presented in Table 6. The result shows
 198 that treatment three (T₃) recorded the highest weight with mean value of 0.28 g followed by 0.18
 199 g recorded in treatment two (T₂) and treatment one (T₁) recorded the least mean value of 0.1. The
 200 result analyzed was significant (P \geq 0.05). The highest weight of seeds was recorded in treatment
 201 three (T₃) followed by treatment two (T₂) which might be attributed to the effect of fertilizers
 202 applied for better growth and grain filling of maize of crop.

203

Table 6: Result of 1000 seeds weight in each subplot

204

| TRTS | R1 | R2 | R3 | R4 | R5 | TOTAL | MEAN |
|-----------|-----|-----|-----|-----|-----|-------|------|
| 1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.5 | 0.1 |
| 2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.9 | 0.18 |
| 3 | 0.3 | 0.2 | 0.3 | 0.3 | 0.3 | 1.4 | 0.28 |
| BLK Total | 0.6 | 0.4 | 0.6 | 0.6 | 0.6 | 2.8 | |

205 **F-LSD (0.05) ***

206 WAP = Week after Planting R = Replication, TRTS =Treatments, * = Significant

207 **WEIGHT OF GRAIN AFTER SHELLING**

208 The result on the weight of grain after shelling is shown in Table 7. The result shows that
209 there was significant ($P \leq 0.05$) difference. Treatment three (T_3) recorded the highest number,
210 mean value of 1.52 followed by treatment two (T_2) which recorded the mean value of 1.34 and
211 treatment one (T_1) recorded the least, mean value of 1.02. The high values recorded in treatment
212 three (T_3) and treatment two (T_2) could be attributed to the fertilizers applied resulting in
213 maximum grain numbers. The result agrees with [7; 15] who reported that maize crop fertilized
214 with fertilizers produced maximum grain number per cob.

215

Table 7: Result of weight of grain after shelling in kg. 216

| TRTS | R1 | R2 | R3 | R4 | R5 | TOTAL | MEAN |
|---------------------|-----|-----|-----|-----|-----|-------|------|
| 1 | 0.9 | 0.8 | 1.2 | 1.0 | 1.2 | 5.1 | 1.02 |
| 2 | 1.3 | 1.4 | 1.4 | 1.3 | 1.3 | 6.7 | 1.34 |
| 3 | 1.3 | 1.6 | 1.5 | 1.7 | 1.5 | 7.6 | 1.52 |
| BLK Total | 3.5 | 3.8 | 4.1 | 4.0 | 4.0 | 19.4 | |
| F-LSD (0.05) | | * | | | | | |

217

218 WAP = Week after Planting R = Replication, TRTS = Treatments, * = Significant

219

220 **CONCLUSION AND RECOMMENDATION**

221 The study concludes that the soil was generally deficient in nutrients for growth of maize.
222 The application of treatments affected the physical and chemical properties of these soils in plots
223 applied with treatments as shown in the growth performance and yield of the crop (maize). This
224 shows that treatment three (T_3) performed the best, followed by treatment two (T_2) while
225 treatment one (T_1) came least in both growth and yield. The solid (granular) fertilizer was found
226 to be more effective and therefore should be recommended for maize production in the area and
227 nutrient management should also be adopted for the soil if it is to be put into agricultural use.

228 **References:**

- 229 1. Adekayode, FO and Ogunkoya, MO. Effect of quantity and placement distances of
230 inorganic 15-15-15 fertilizers in improving soil fertility status and the performance and
231 yield of maize in tropical rain forest zone of Nigeria, 2010. *Journal of Soil Science and*
232 *Environmental Management*.1: 155-163.
- 233 2. Agba, OB, Ubi, BE, Abam, P, Ogechi, J, Akeh, M, Odey, S and Ogar, N. Evaluation of
234 Agronomic performance of Maize (*Zea mays* L) under Different Rates of poultry manure
235 application in Ultisol of Obubra, Cross River State, Nigeria, 2012. *International Journal*
236 *of Agriculture and Forestry*. 2 (4): 138 – 144
- 237 3. Bates, RG. Determination of pH theory and practice, 1973. Wiley
- 238 4. Bray, RH and Kurtz, LT. Determination of total, organic and available forms of
239 phosphorus in soils.1945. *Soil Sci.*, 59: 39-46
- 240 5. Dilshad, MD, Lone, MD, Jilani, G., Malik, MA.,Yousaf, M, Khalid, A and Shamin, F
241 Integrated Plant Nutrient Management on maize under rainfed condition, 2010. *Pakistan*
242 *Journal of Nutrition*. 9: 896-901
- 243 6. Dowswell, CR, Paliwal, RL and Cantrell, RP. Maize in the Third World, 1996. West
244 View Press, Colorado, USA. ISBN- 13: 9780813389639. Page 268
- 245 7. Eleweanya, NP, Uguru, MI, Enebong, E E. and Okocha, PI. Correlation and path
246 coefficient analysis of grain yield related characters in maize (*Zea mays* L) under
247 Umudike conditions of Southeastern Nigeria, 2005. *Agro – Science Journal of Tropical*
248 *of Agriculture, Food, Environment and Extension*. 1, 24-28
- 249 8. Ezeaku, PL. Optimum NPK fertilizer rates based on soil data for grain maize (*Zea mays*
250 L) production in some soils of Southeastern Nigeria, 2008. *Agricultural Journal*. 3 (1):
251 36-41
- 252 9. Gualem, P, Gustafson, DM and Wicks, III.”Phosphorus Concentration uptake and Dry
253 Matter Yield of corn Hybrids”, 2011.*World Journal of Agricultural Sciences*, 7 (4): 418-
254 424, ISSN: 1817 – 3047
- 255 10. Ibia, TO, Uko-Itakha, IB, Edem, SO, Ogban, PI & Obi, JC. Evaluation of the Acid Soils
256 for sanitary Landfills in Akwa-Ibom State, Southern Nigeria, 2011. *Nigerian Journal of*
257 *Soil Science*. Vol 21(1):1-5
- 258 11. Myers, RJ. One – Hundred years of pH, 2010.*Journal of Chemical Education*. **87**: 30 –
259 33
- 260 12. Negassa, W and Gebrekidan, H. Forms of phosphorus and status of available
261 micronutrients under different land use systems of alfisols in Bako area of Ethiopia.
262 2003. *Journal of Natural Resources*. 5: 17-37
- 263 13. Nelson, OW and Sommers, LE. Total Carbon, Organic Carbon and Organic. In O. L.
264 Sparks (ed). *Methods of Soil Analysis Part 3, Chemical Methods*, 1996. Soil Science
265 Society of America Book Series Number 5. American Society of Agronomy, Madison
266 WIE, pp 961-1010.
- 267 14. Onasanya, RO, Aiyelari, OP, Onasanya, AW, Wilene, FE. and Onyelakin, OO. Effect of
268 different level of Nitrogen and phosphorus fertilizers on the growth and yield of maize

- 269 (Zea mays L) in Southwest Nigeria,2009. *Nigeria International Journal of Agricultural*
270 *Resources*. 4: 193 -203
- 271 15. Rasheed, M, Ali, H and Mahmood, T. Impact of Nitrogen and sulfur application on
272 growth and yield of maize (Zea mays L) crop, 2004. *J. Res. Sci.* 15: 153-157
- 273 16. Rhodes, LL and Eagles, HA. “Origins of maize in Zealand”, 1984. *Zealand Journal of*
274 *Agricultural Research*. 27 (2): 151-156
- 275 17. Salk, H, Varadachari, Cand Gosh, K.. Changes in carbon, nitrogen and phosphorus levels
276 due to deforestation and cultivation, 1998. A case study in Simpal National Park.
- 277 18. Soil Survey Staff .Soil Survey Manual. US Department of Agriculture, 1999. Hand Book
278 18. US Govt. Printing Office, Washington.
- 279 19. Tadesse, T, Dechassa, N, Bayu, W and Gebeyehu, S. Effects of farmyard manure and
280 inorganic fertilizer application on soil physicochemical properties and nutrient balance in
281 rain-fed lowland rice ecosystem 2013. *American Journal of Plant Science* 4: 309-316.
- 282 20. Thomas, EE, Kristin, MH and Mary, LS. Marginal Horticulturists or Maize
283 Agriculturists: Archaeobotanical, Paleopathological and Isotopic Evidence Relating to
284 Landford Tradition Maize Consumption, 2005. *Midcontinental Journal of Archaeology*.
285 30 (1):67-118
- 286 21. Udo, EJ, Ibia, TO, Ogunwale, JA, Ano, AO and Esu, IE. Manual of soil, plant and water
287 analysis, 2009. Sibon books limited Lagos Nigeria. pp.183.
- 288 22. Mallis, A. *Handbook of pest control*, 1982. Clevelad: Franzak Foster Co.
- 289 23. Hock J, Kranz, J. Studies on the epidemiology of the tar spot disease complex of maize in
290 Mexico, *Plant Pathology*, 1995. British Society for plant pathology. 44(3): 410 – 502:
291 doi: 10.1111/J.1365-3059.1995.tb01671.X

292
293
294
295
296
297