

## Effect of Nitrogen and Phosphorus on the Growth and Seed Yield of Spinach

### ABSTRACT

The experiment was conducted at the farm of Sher-e-Bangla Agricultural University, Dhaka, Bangladesh during Rabi (November 2017 to March 2018) season to find out the growth, yield and economic benefit of spinach seed as influenced by nitrogen and phosphorus. The research involved two factors. Factor A: Four levels of nitrogen;  $N_0 = 0$  kg/ha,  $N_1 = 27.6$  kg/ha,  $N_2 = 55.2$  kg/ha,  $N_3 = 82.8$  kg/ha, and factor B: Four levels of phosphorus;  $P_0 = 0$  kg/ha;  $P_1 = 15.84$  kg/ha,  $P_2 = 31.68$  kg/ha,  $P_3 = 47.52$  kg/ha. There were 16 treatment combinations in the experiment and laid out in Randomized Complete Block Design (RCBD) with three replications. Quality tests of seeds were done based on the germination test (%), seed vigor test (Electrical conductivity). In case of nitrogen, the highest seed yield ( $1.10 \text{ t ha}^{-1}$ ), germination percentage (87.33 %) and lowest value in EC test ( $11.87 \text{ dS/cm}$ ) were obtained from  $N_2$ , while the lowest seed yield ( $0.81 \text{ t ha}^{-1}$ ), germination percentage (79.33 %) and highest value in EC test ( $13.87 \text{ dS/cm}$ ) from  $N_0$ . For phosphorus levels, the highest seed yield ( $1.05 \text{ t ha}^{-1}$ ), germination percentage (86.58 %) and lowest value in EC test ( $11.79 \text{ dS/cm}$ ) were recorded from  $P_2$ , whereas the lowest seed yield ( $0.84 \text{ t ha}^{-1}$ ), germination percentage (79.91 %) and highest value in EC test ( $13.35 \text{ dS/cm}$ ) from  $P_0$ . Due to mutual effect, the highest seed yield ( $1.30 \text{ t ha}^{-1}$ ), germination percentage (91.33 %) and lowest value in EC test ( $10.2 \text{ dS/cm}$ ) were noted from  $N_2P_2$ , whereas the lowest seed yield ( $0.69 \text{ t ha}^{-1}$ ), germination percentage (72.66 %) and highest value in EC test ( $14.83 \text{ dS/cm}$ ) from  $N_0P_0$ . From the economic point of view, the highest Benefit-Cost Ratio (BCR) (1.59) was found in the treatment of  $N_2P_2$  and the lowest BCR (1.01) was found in the treatment of  $N_0P_0$ . It is apparent that the treatment combination  $N_2P_2$  gave the best performance for the seed yield and economic benefit of spinach.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

*Keywords: Growth, Nitrogen, Phosphorus, Seed Yield*

### 1. INTRODUCTION

Spinach (*Spinacia oleracea*) is a leafy green cool-season vegetable that is known for its nutritive value and is considered one of the most popular vegetables in Bangladesh. It is believed to have originated from Persia. Its leaves are a common edible vegetable. By weight, spinach consists of 91.4% water, 3.6% carbons, and 2.9% protein [1]. There are 23 calories in 100 grams of spinach and further benefits of spinach leaves consumption can be of more value e.g vital mineral elements in spinach compared to other vegetables e.g iron which is very good for women. The seed is produced for commercial consumption and for seed companies that supply seed throughout the country. In Bangladesh spinach occupies 22000 acres [2] with an annual production of 66000 tons. In comparison to other countries, this yields much lower. So, to use of quality seeds of high yield varieties are the foremost important technique for maximizing yield per unit area. Quality seed can increase vegetable production by up to 25-50% [1]. Farmers save seeds annually of about 50 tons and used

29 every year, which in most cases are of inferior in quality [3]. Appropriate fertilizer application  
30 to the plants greatly affects their growth and production. Nitrogen strongly stimulates growth,  
31 expansion of the crop canopy and interception of solar radiation [4]. Increasing the levels of  
32 nitrogen during the vegetative stage can strengthen and allow a plant to grow more rapidly  
33 and produce large amounts of succulent, green foliage, which in turn can generate higher  
34 yields [5]. Similarly, phosphorus (P) is an essential nutrient act as catalysts in the conversion  
35 of numerous key biochemical reactions in plants. Phosphorus stimulates root development,  
36 improves flower formation, seed production and improves crop quality and increases  
37 resistance to plant diseases [6]. Leafy vegetables, particularly, spinach is highly responsive  
38 to fertilization [7] and oxalates are the main indexes of the quality due to a very efficient  
39 uptake system and inefficient reductive systems [8]. The fertilizer requirements on sandy and  
40 sandy loams are 85 to 120 kg N, 75 to 85 kg P<sub>2</sub>O<sub>5</sub>, and 85 to 150 kg K<sub>2</sub>O. On heavier clay  
41 soils, 75 kg ha<sup>-1</sup> of each nutrient should be adequate. If the fertilizer is banded at seeding it  
42 should be placed along each side of the rows 2 to 3 inches below the level of the seed and 6  
43 inches to the side of the row; fertilizer should never come in contact with the seed and two or  
44 three splits of 85 to 120 kg ha<sup>-1</sup> N would be adequate as side-dressing [9]. An adequate  
45 supply of fertilizers can promote plant growth and increase crop production, but excessive  
46 and inappropriate use of chemical fertilizers causes accumulation of compounds in the  
47 edible products which can be detrimental to human health in addition cause environmental  
48 pollution and economic losses [10]. [11] found that N increased the spinach yield and  
49 enhanced the accumulation of N and P in leaves. [12] reported that application of 40 kg N +  
50 15.0 kg P<sub>2</sub>O<sub>5</sub> increased plant fresh yield by 27.2 and 42.3% and 16.3 and 10.4% in seed  
51 yield in relation to the control in the first and second seasons, respectively. [13] achieved the  
52 highest yield with 120 kg N ha<sup>-1</sup> Farmers in Bangladesh generally do not use any improved  
53 or special techniques for quality seed production. There for to achieve higher seeds yield  
54 with good quality fertilizer management is an important practice which cannot be  
55 emphasized. The present study was thus undertaken to evaluate the effects of nitrogen and  
56 phosphorus on the growth, yield and economic benefit of spinach (*Spinacia oleracea*) seed.

## 57 58 **2. MATERIAL AND METHODS**

### 59 **2.1. Experimental Site**

60 The experiment was conducted at the Agronomy Research Farm of Sher-e-Bangla  
61 Agricultural University (SAU), Dhaka, Bangladesh from November 2017 to March 2018.  
62 Experimental site situated an elevation of 8 meters above the sea level in the Agro-  
63 ecological zone of "Madhupur Tract" (AEZ-28). The soil was sandy loam and medium high  
64 land in texture having pH 5.46- 5.62.

### 65 66 **2.2 Experiment Frame Work**

67 The research was consisted of two factors: Factor A: Four levels of nitrogen; N<sub>0</sub>= 0 kg/ha  
68 N<sub>1</sub>= 27.6 kg/ha, N<sub>2</sub>=55.2 kg/ha, N<sub>3</sub>= 82.8 kg/ha, and factor B: Four levels of phosphorus; P<sub>0</sub>=  
69 0 kg/ha; P<sub>1</sub> = 15.84 kg/ha, P<sub>2</sub>= 31.68 kg/ha, P<sub>3</sub>= 47.52 kg/ha. The two factors experiment  
70 was laid out following the Randomized Complete Block Design (RCBD) with three  
71 replications. The experiment was divided into three equal blocks where each block was  
72 divided into 16 plots. Then 16 treatment combinations were allotted randomly in each block.  
73 The size of each unit plot was 1.5 m × 1 m. The distance maintained between two blocks  
74 and two plots were 0.75 m and 0.5 m, respectively. Row to row distance was 30 cm and  
75 plant to plant distance was 20 cm.

### 76 77 **2.3. Application of manure and fertilizers**

78 About 5 t ha<sup>-1</sup> well-decomposed cow dung was applied as the control (N<sub>0</sub>P<sub>0</sub>) treatment) plot  
79 and incorporated adequately to the soil during final land preparation whereas other plots  
80 were applied with inorganic fertilizer as per treatment. Doses of inorganic fertilizers (Urea  
81 and Triple superphosphate) were applied in the experimental plot according to the

82 treatments (Table 1). The whole amount of TSP and half the amount of urea and MoP (180  
 83 kg/ha) were also applied as basal dose before sowing of seed in the main field. 1<sup>st</sup> top  
 84 dressing of urea was applied when seedlings established in the main field about 10 days  
 85 after seed sowing. 2<sup>nd</sup> top dressing of urea and the rest amount of MoP was applied about  
 86 25days after 1st top dressing. Then the rest amount of urea was applied as 3<sup>rd</sup> installment  
 87 after flowering. Each top dressing was followed by manual irrigation.

88  
 89

**Table 1. Doses of nutrients application in the main field as per treatment**

Treatments	N rates (kg ha <sup>-1</sup> )	Fertilizers (kg ha <sup>-1</sup> )	Doses (g plot <sup>-1</sup> )	Treatments	P rates (kg ha <sup>-1</sup> )	Fertilizes (kg ha <sup>-1</sup> )	Doses (g plot <sup>-1</sup> )
	<b>N</b>	<b>Urea</b>	<b>Urea</b>		<b>P</b>	<b>TSP</b>	<b>TSP</b>
<b>N<sub>0</sub></b>	0	0	0	<b>P<sub>0</sub></b>	0	0	0
<b>N<sub>1</sub></b>	27.6	60	9	<b>P<sub>1</sub></b>	15.84	75	11.25
<b>N<sub>2</sub></b>	55.2	120	18	<b>P<sub>2</sub></b>	31.68	150	22.5
<b>N<sub>3</sub></b>	82.8	180	27	<b>P<sub>3</sub></b>	47.52	225	33.75

90  
 91

#### 2.4. Economic analysis

92 The cost of production was analyzed in order to find out the most economic treatment of  
 93 nitrogen and phosphorus for quality seed production of spinach. All the non-material and  
 94 material input costs and interests in running capital were considered for computing the cost  
 95 of production. The benefit-cost ratio (BCR) was calculated by the following formula:  
 96 **Benefit-cost ratio (BCR) = Gross return (tk/ha) ÷ Total cost of production (tk/ha)**

97  
 98

#### 2.5. Statistical analysis

99 The data obtained for different characters were statistically analyzed to observe the  
 100 significant difference among the treatment by using the STATISTIX-10 computer package  
 101 program. The mean values of all the characters were calculated and analysis of variance  
 102 was performed. The significance of the difference among the treatments means was  
 103 estimated by the Least Significant Different Test (LSD) at 5% level of probability.

104  
 105

### 3. RESULTS AND DISCUSSION

#### 3.1 Plant height (cm)

106 A significant variation was observed on the plant height of spinach due to the application of  
 107 different levels of nitrogen (Table 2.). The highest plant height 13.94cm and 47.69cm were  
 108 obtained before flowering and at the time of harvest respectively from N<sub>3</sub> while the lowest  
 109 plant height 11.01cm and 34.88cm were obtained before flowering and at the time of harvest  
 110 respectively from N<sub>0</sub>. P<sub>3</sub> showed the highest plant height 13.64cm and 46.68cm before  
 111 flowering and at the time of harvest respectively and the lowest plant height 11.60 cm and  
 112 35.18 cm at before flowering and at the time of harvest respectively in P<sub>0</sub> (Table 3).

114 The combined effect of different levels of nitrogen and phosphorus application showed a  
 115 significant effect on the plant height of spinach (Table 4). The highest plant height 15.21cm  
 116 and 55.25cm before flowering and at the time of harvest, respectively was observed in N<sub>3</sub>P<sub>3</sub>  
 117 while the lowest plant height 10.10cm and 27.7cm before flowering and at the time of  
 118 harvest, respectively in N<sub>0</sub>P<sub>0</sub>.

#### 3.2. Number of leaves per plant

120 The effect of nitrogen on the number of leaves per plant of spinach was significant (Table 2).  
 121 The highest number of leaves (8.46) was produced from N<sub>2</sub> and the lowest number of leaves

122 (6.42) was observed in  $N_0$ .  $P_2$  showed the maximum leaves per plant (8.30) and the  
 123 minimum leaves per plant (6.36) was observed in  $P_0$  (Table 3).

124 The combined effect of different levels of nitrogen and phosphorus showed a significant  
 125 effect on the number of leaves per plant of spinach (Table 4). The maximum leaves per plant  
 126 (10.35) were observed in  $N_2P_2$  and the lowest leaves per plant (5.01) were recorded with  
 127  $N_0P_0$ , [14] also found a similar result.  
 128

129 **Table 2.** Effect of nitrogen on plant height (before flowering and at the time of harvest)  
 130 leaves per plant of spinach

Treatments	Plant height before flowering (cm)	Plant height at the time of harvest (cm)	Leaves per plant
$N_0$	11.01	34.88	6.42
$N_1$	12.34	40.35	7.24
$N_2$	13.07	43.75	8.46
$N_3$	13.94	47.69	7.84
<b>LSD</b>	<b>0.87</b>	<b>3.12</b>	<b>0.54</b>
<b>CV %</b>	<b>8.34</b>	<b>9.00</b>	<b>8.70</b>

131 Here,  $N_0= 0$  kg/ha;  $N_1=27.6$  kg/ha;  $N_2= 55.2$  kg/ha;  $N_3= 82.8$  kg/ha

132 **Table 3.** Effect of phosphorus on plant height (before and at the time of harvest), Leaves per  
 133 plant of spinach

Treatments	Plant height before flowering	Plant height at the time of harvest	Leaves per plant
$P_0$	11.60	35.18	6.36
$P_1$	12.16	40.37	7.24
$P_2$	12.96	44.44	8.30
$P_3$	13.64	46.68	8.06
<b>LSD</b>	<b>0.87</b>	<b>3.11</b>	<b>0.543</b>
<b>CV %</b>	<b>8.34</b>	<b>9.00</b>	<b>8.70</b>

134 Here,  $P_0= 0$  kg/ha;  $P_1= 15.84$  kg/ha;  $P_2= 31.68$  kg/ha;  $P_3= 47.52$  kg/ha.

135 **Table 4.** Combined effect of nitrogen and phosphorus on the plant height (before and at the  
 136 time of harvesting), leaves per plant of spinach

Treatments	Plant height before flowering (cm)	Plant height at the time of harvest (cm)	Leaves per plant
$N_0P_0$	10.10	27.7	5.01
$N_0P_1$	10.63	34.34	6.25
$N_0P_2$	11.25	37.31	6.66
$N_0P_3$	12.07	40.18	7.76
$N_1P_0$	11.49	34.65	6.60
$N_1P_1$	12.15	41.03	7.33
$N_1P_2$	12.60	42.44	7.46
$N_1P_3$	13.12	43.29	7.58
$N_2P_0$	12.14	37.88	6.91
$N_2P_1$	12.54	21.80	7.81
$N_2P_2$	13.42	47.31	10.35
$N_2P_3$	14.17	48.01	8.76
$N_3P_0$	12.65	40.51	6.93
$N_3P_1$	13.35	44.30	7.56

<b>N<sub>3</sub> P<sub>2</sub></b>	14.56	50.71	8.73
<b>N<sub>3</sub> P<sub>3</sub></b>	15.21	55.25	8.13
<b>LSD</b>	<b>1.75</b>	<b>6.25</b>	<b>1.08</b>
<b>CV %</b>	<b>8.34</b>	<b>9.00</b>	<b>8.70</b>

137 Here, N<sub>0</sub>= 0 kg/ha; N<sub>1</sub>=27.6 kg/ha; N<sub>2</sub>= 55.2 kg/ha; N<sub>3</sub>= 82.8 kg/ha.  
 138 P<sub>0</sub>= 0 kg/ha; P<sub>1</sub>= 15.84 kg/ha; P<sub>2</sub>= 31.68 kg/ha; P<sub>3</sub>= 47.52 kg/ha.

139

140

### 3.3. Number of inflorescence per plant

141

Statistically significant differences were found on the number of inflorescence per plant of spinach due to the application of nitrogen (Table 5). The highest number of inflorescence per plant (8.03) was recorded from N<sub>2</sub> whereas, the lowest number (5.17) was observed from N<sub>0</sub>. The highest number of inflorescence per plant (7.80) was recorded from P<sub>2</sub> and the lowest (4.94) was found from P<sub>0</sub> (Table 6).

142

The combined effect of nitrogen and phosphorus showed significant variation in the number of inflorescences per plant (Table 7). The highest number of inflorescence per plant (10.53) was recorded from N<sub>2</sub>P<sub>2</sub> and the lowest number of inflorescence per plant (3.90) from N<sub>0</sub>P<sub>0</sub>.

143

144

145

### 3.4. Length of inflorescence

146

A significant variation was observed on the length of an inflorescence of spinach when different levels of nitrogen were applied (Table 5). The highest length of inflorescence (29.54cm) was recorded in N<sub>2</sub> and the lowest length of inflorescence (19.09 cm) from N<sub>0</sub>. The highest length of inflorescence (28.97 cm) was recorded in P<sub>3</sub> and the lowest length of inflorescence (20.58 cm) was recorded in P<sub>0</sub> (Table 6).

147

148

149

The combined effect of different levels of nitrogen and phosphorus showed a significant effect on the length of inflorescence (Table 7). The highest length of inflorescence (35.04 cm) was observed in N<sub>2</sub>P<sub>2</sub> and the lowest length of inflorescence (15.19 cm) was recorded in N<sub>0</sub>P<sub>0</sub>.

150

151

### 3.5. Number of seeds per inflorescence

152

Statistically significant differences were found on the number of seeds per inflorescence of spinach due to the application of different nitrogen levels (Table 5). The maximum seeds per inflorescence (58.68) was recorded from N<sub>2</sub> (55.2 kg/ha) and the minimum (38.15) was observed from N<sub>0</sub>. The highest seeds per inflorescence (57.19) were recorded from P<sub>2</sub> and the lowest number of seeds per inflorescence (39.13) in P<sub>0</sub> (Table 6).

153

154

155

The number of seeds per inflorescence was significantly influenced by the combined application of nitrogen and phosphorus (Table 7). The maximum number of seeds per inflorescence (70.41) was recorded from N<sub>2</sub>P<sub>2</sub> and the lowest number of seeds per inflorescence (35.33) in N<sub>0</sub>P<sub>0</sub>.

156

157

**Table 5. Effect of nitrogen on number of inflorescence per plant, inflorescence length and seeds per inflorescence of spinach**

158

159

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
N <sub>0</sub>	5.17	19.99	38.15
N <sub>1</sub>	6.30	23.56	47.37
N <sub>2</sub>	8.03	29.54	58.86
N <sub>3</sub>	7.13	29.01	56.28
<b>LSD</b>	<b>0.47</b>	<b>1.857</b>	<b>3.40</b>

160

161

162

163

164

165

166

167

168

169

170

171

172

173

	<b>CV %</b>	<b>8.48</b>	<b>8.73</b>	<b>8.12</b>
174	Here,	N <sub>0</sub> = 0 kg/ha;	N <sub>1</sub> =27.6 kg/ha;	N <sub>2</sub> = 55.2 kg/ha; N <sub>3</sub> = 82.8 kg/ha.
175				

176 **Table 6. Effect of phosphorus on number of inflorescence per plant, inflorescence length**  
177 **and seeds per inflorescence of spinach**

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
P <sub>0</sub>	4.94	20.58	39.13
P <sub>1</sub>	6.26	24.12	48.19
P <sub>2</sub>	7.80	28.43	57.19
P <sub>3</sub>	7.64	28.97	56.15
<b>LSD</b>	<b>0.48</b>	<b>1.85</b>	<b>3.39</b>
<b>CV %</b>	<b>8.48</b>	<b>8.73</b>	<b>8.12</b>

178 Here, P<sub>0</sub>= 0 kg/ha; P<sub>1</sub>= 15.84 kg/ha; P<sub>2</sub>= 31.68 kg/ha; P<sub>3</sub>= 47.52 kg/ha.

179 **Table 7. Combined effect of nitrogen and phosphorus on number of inflorescence per plant,**  
180 **inflorescence length and seeds per inflorescence of spinach**

Treatments	Number of inflorescence per plant	Inflorescence length (cm)	Seeds per inflorescence
N <sub>0</sub> P <sub>0</sub>	3.9	15.19	28.03
N <sub>0</sub> P <sub>1</sub>	5.03	18.95	35.33
N <sub>0</sub> P <sub>2</sub>	5.70	21.86	43.82
N <sub>0</sub> P <sub>3</sub>	6.06	23.96	45.43
N <sub>1</sub> P <sub>0</sub>	5.08	19.91	36.27
N <sub>1</sub> P <sub>1</sub>	6.23	22.91	47.79
N <sub>1</sub> P <sub>2</sub>	6.75	24.72	51.74
N <sub>1</sub> P <sub>3</sub>	7.15	26.69	53.70
N <sub>2</sub> P <sub>0</sub>	5.23	22.84	45.25
N <sub>2</sub> P <sub>1</sub>	7.05	26.15	54.29
N <sub>2</sub> P <sub>2</sub>	10.53	35.04	70.41
N <sub>2</sub> P <sub>3</sub>	9.31	34.15	65.48
N <sub>3</sub> P <sub>0</sub>	5.55	24.38	47.00
N <sub>3</sub> P <sub>1</sub>	6.73	28.48	55.34
N <sub>3</sub> P <sub>2</sub>	8.21	32.10	62.80
N <sub>3</sub> P <sub>3</sub>	8.03	31.10	60.00
<b>LSD</b>	<b>0.94</b>	<b>3.71</b>	<b>6.79</b>
<b>CV (%)</b>	<b>8.48</b>	<b>8.73</b>	<b>8.12</b>

181 Here, N<sub>0</sub>= 0 kg/ha; N<sub>1</sub>=27.6 kg/ha; N<sub>2</sub>= 55.2 kg/ha; N<sub>3</sub>= 82.8 kg/ha.  
182 P<sub>0</sub>= 0 kg/ha; P<sub>1</sub>= 15.84 kg/ha; P<sub>2</sub>= 31.68 kg/ha; P<sub>3</sub>= 47.52 kg/ha.

183

### 184 3.6. Seed yield per hectare (ton)

185 Statistically significant differences were found for seed yield per hectare of spinach due to  
186 different nitrogen levels (Table 8). The maximum seed yield (1.10 t ha<sup>-1</sup>) was recorded from  
187 N<sub>2</sub> and the lowest (0.81 t ha<sup>-1</sup>) was recorded from N<sub>0</sub>. The maximum seed yield (1.05 t ha<sup>-1</sup>)  
188 was recorded from P<sub>2</sub> and the lowest (0.84 t ha<sup>-1</sup>) was recorded from P<sub>0</sub> (Table 9.). The  
189 combined effect of nitrogen and phosphorus showed significant variation in the seed yield of  
190 spinach (Table 10). The highest seed yield (1.30 t/ha) was recorded from the combination of  
191 N<sub>2</sub>P<sub>2</sub> and the lowest (0.69 t ha<sup>-1</sup>) was recorded from N<sub>0</sub>P<sub>0</sub> treatment combination.

192

### 193 3.7. 1000 seed weight (g)

194 Statistically significant differences were found for 1000 seed weight of spinach due to the  
 195 nitrogen level (Table 8). The maximum 1000 seed weight (10.19 g) was recorded from N<sub>2</sub>  
 196 and the lowest (9.24 g) was recorded from N<sub>0</sub>. The maximum 1000 seed weight (10.20 g)  
 197 was recorded from P<sub>2</sub> and the lowest (9.23 g) was recorded from P<sub>0</sub> (Table 9). The combined  
 198 effect of nitrogen and phosphorus showed significant variation in 1000 seed weight of  
 199 spinach (Table 10). The highest 1000 seed weight (11.06 g) was recorded from the  
 200 combination of N<sub>2</sub>P<sub>2</sub> and the lowest (8.77 g) from N<sub>0</sub>P<sub>0</sub>.

### 201 3.8. Germination percentage

202 A significant difference was found on germination % due to fertilizer level (Table 8). The  
 203 maximum germination percentage (87.33 %) was recorded from N<sub>2</sub> and the minimum (79.33  
 204 %) was found in N<sub>0</sub>. [15] reported that seed yield and its quality of spinach obtained that  
 205 increasing the rate of nitrogen levels up to the levels at 60 kg N produced higher seed yield  
 206 with the best quality, germination percentage and germination rate. The maximum  
 207 germination percentage (87.08 %) was recorded from P<sub>3</sub> and the minimum (79.91 %) was  
 208 found from P<sub>0</sub> (Table 9).

209 The combined effect of nitrogen and phosphorus was significant on the germination  
 210 percentage of spinach (Table 10). The highest germination percentage (91.33 %) was  
 211 recorded from the combination of N<sub>2</sub>P<sub>2</sub> and the lowest germination percentage (72.66 %)  
 212 was recorded in N<sub>0</sub>P<sub>0</sub> (control). [14] experimented on the farm of the Department of  
 213 Horticulture, BSMRAU, Salna, Gazipur on 6 spinach genotypes to observe their seed  
 214 production potentiality and to evaluate the quality of produced seed. They reported that the  
 215 quality test of seed was done based on germination test (%), seed vigor test (Electrical  
 216 conductivity), moisture test (%) and thousand seed weight (g) of seeds.

### 217 3.9. Electrical conductivity test

218 A significant difference was found in electrical conductivity test value due to the application  
 219 of different levels of nitrogen (Table 8). The highest EC test value (13.87 dS/cm) was  
 220 recorded from treatment N<sub>0</sub> whereas, the minimum (11.16 dS/cm) was found in N<sub>2</sub>. The  
 221 maximum EC test value (13.35 dS/cm) was recorded from treatment P<sub>0</sub> and the lowest EC  
 222 test value (11.79 dS/cm) was recorded in P<sub>2</sub> (Table 9). **The combined effect of nitrogen and  
 223 phosphorus was significant on the EC test value of spinach (Table 10).** The highest EC test  
 224 value (14.83 dS/cm) was recorded from the combination (N<sub>0</sub>P<sub>0</sub>) and the lowest EC test value  
 225 (10.20 dS/cm) was recorded with N<sub>2</sub>P<sub>2</sub> treatment combination. Why highest in N<sub>2</sub>P<sub>2</sub> and not  
 226 N<sub>3</sub>P<sub>3</sub>  
 227

228 **Table 8. Effect of nitrogen on seed yield per hectare 1000 seed weight, germination**  
 229 **percentage and electrical conductivity test**

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
N <sub>0</sub>	0.81	9.24	79.33	13.87
N <sub>1</sub>	0.91	9.58	83.41	12.78
N <sub>2</sub>	1.10	10.19	87.33	11.16
N <sub>3</sub>	1.01	10.14	87.25	11.73
LSD	0.06	0.75	4.95	0.85
CV (%)	8.28	9.19	7.05	8.25

230

231 **Table 9.** Effect of phosphorus on seed yield per hectare, 1000 seed weight, germination  
 232 percentage and electrical conductivity test

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
P <sub>0</sub>	0.84 c	9.23 b	79.91	13.35
P <sub>1</sub>	0.93 b	9.56 ab	83.75	12.57
P <sub>2</sub>	1.05 a	10.20 a	86.58	11.79
P <sub>3</sub>	1.01 a	10.18 a	87.08	11.81
LSD	<b>0.067</b>	<b>0.75</b>	<b>4.95</b>	<b>0.85</b>
CV (%)	<b>8.28</b>	<b>9.19</b>	<b>7.05</b>	<b>8.25</b>

233 Here, P<sub>0</sub>= 0 kg/ha; P<sub>1</sub>= 15.84 kg/ha; P<sub>2</sub>= 31.68 kg/ha; P<sub>3</sub>= 47.52 kg/ha.

234 **Table 10.** The combined effect of nitrogen and phosphorus seed yield per hectare, 1000  
 235 seed weight, germination percentage and electrical conductivity test

Treatments	Seed yield per hectare (t)	1000 seed weight (g)	Germination (%)	Electrical conductivity test (ds/cm)
N <sub>0</sub> P <sub>0</sub>	0.69	8.77	72.66	14.83
N <sub>0</sub> P <sub>1</sub>	0.83	9.22	80.00	13.97
N <sub>0</sub> P <sub>2</sub>	0.85	9.43	81.00	13.37
N <sub>0</sub> P <sub>3</sub>	0.88	9.56	83.66	13.30
N <sub>1</sub> P <sub>0</sub>	0.83	9.15	79.00	13.55
N <sub>1</sub> P <sub>1</sub>	0.91	9.47	84.00	12.85
N <sub>1</sub> P <sub>2</sub>	0.95	9.76	84.66	12.58
N <sub>1</sub> P <sub>3</sub>	0.96	9.95	86.00	12.14
N <sub>2</sub> P <sub>0</sub>	0.93	9.26	83.33	12.59
N <sub>2</sub> P <sub>1</sub>	0.99	9.74	84.66	11.46
N <sub>2</sub> P <sub>2</sub>	1.30	11.06	91.33	10.20
N <sub>2</sub> P <sub>3</sub>	1.16	10.70	90.00	10.42
N <sub>3</sub> P <sub>0</sub>	0.92	9.73	84.66	12.45
N <sub>3</sub> P <sub>1</sub>	0.98	9.80	86.33	12.01
N <sub>3</sub> P <sub>2</sub>	1.09	10.53	89.33	11.03
N <sub>3</sub> P <sub>3</sub>	1.04	10.50	88.66	11.41
LSD	<b>0.13</b>	<b>1.50</b>	<b>9.91</b>	<b>1.70</b>
CV (%)	<b>8.28</b>	<b>9.19</b>	<b>7.05</b>	<b>8.25</b>

236 Here, N<sub>0</sub>= 0 kg/ha; N<sub>1</sub>=27.6 kg/ha; N<sub>2</sub>= 55.2 kg/ha; N<sub>3</sub>= 82.8 kg/ha.  
 237 P<sub>0</sub>= 0 kg/ha; P<sub>1</sub>= 15.84 kg/ha; P<sub>2</sub>= 31.68 kg/ha; P<sub>3</sub>= 47.52 kg/ha.

238

### 239 **3.10. Benefit cost ratio (BCR)**

240

241 The combination of nitrogen and phosphorus showed different benefit cost ratio. The highest  
 242 benefit cost ratio (1.59) which is economically more benefitted was obtained from N<sub>2</sub>P<sub>2</sub> and  
 243 the lowest benefit cost ratio (1.01) which is economically less benefitted was obtained from  
 244 the treatment combination of N<sub>0</sub>P<sub>0</sub>(Table.11).

245

246 **Table 11.** Cost and return analysis of spinach seed considering nitrogen and phosphorus

Treatments	Seed yield (t/ha)	Gross return (Tk/ha)	Total cost of production(Tk)	Net return (Tk/ha)	Benefit cost ratio (BCR)
N <sub>0</sub> P <sub>0</sub>	0.69	117000	116656.8	13156	1.01



N <sub>0</sub> P <sub>1</sub>	0.83	124500	118342	6158	1.05
N <sub>0</sub> P <sub>2</sub>	0.85	127500	120027.3	7472.7	1.06
N <sub>0</sub> P <sub>3</sub>	0.88	132000	121712.5	10287.5	1.08
N <sub>1</sub> P <sub>0</sub>	0.83	124500	117870.1	6629.9	1.05
N <sub>1</sub> P <sub>1</sub>	0.91	136500	119555.4	16944.6	1.14
N <sub>1</sub> P <sub>2</sub>	0.95	142500	121240.6	21259.4	1.17
N <sub>1</sub> P <sub>3</sub>	0.96	144000	122925.9	21074.1	1.17
N <sub>2</sub> P <sub>0</sub>	0.93	133500	119083.5	14416.5	1.12
N <sub>2</sub> P <sub>1</sub>	0.99	148500	120768.8	27731.2	1.22
N <sub>2</sub> P <sub>2</sub>	1.30	195000	122454	72546	1.59
N <sub>2</sub> P <sub>3</sub>	1.16	174000	124139.3	49860.7	1.40
N <sub>3</sub> P <sub>0</sub>	0.92	138000	120296.9	17703.1	1.14
N <sub>3</sub> P <sub>1</sub>	0.98	147000	121982.1	25017.9	1.20
N <sub>3</sub> P <sub>2</sub>	1.09	163500	123667.4	39832.6	1.32
N <sub>3</sub> P <sub>3</sub>	1.04	156000	125352.6	30647.4	1.24

247 Here,

- 248     > Sale of marketable seed @ Tk.150/kg  
249     > Gross return = Marketable yield x Tk./kg  
250     > Net income = Gross return-Total cost of production  
251     > BCR = Gross return ÷ cost of production

252

#### 253 4. CONCLUSION

254

255 Both seed yield and economic benefit of crop are important for the seed production.  
256 Application of nitrogen (55.2 kg/ha) represents higher seed yield in spinach than without no  
257 nitrogen and (82.8 kg/ha) nitrogen. According to the results of the present experiment it may  
258 be concluded that efficient production of spinach seed is increased by the application of  
259 phosphorus (31.68 kg/ha). The combined effect of nitrogen and phosphorus had positive  
260 effect on morphological characters, yield contributing characters but seed yield and seed  
261 quality of spinach increased by the combined application of (N 55.2 kg/ha+ P 31.68 kg/ha).  
262 On the basis of benefit cost ratio, it may be suggested that application of N 55.2 kg/ha with P  
263 31.68 kg/ha combination seemed to be more suitable for the highest values of gross return,  
264 net return as well as highest beneficial cost ratio of spinach..  
265

266

#### 267 COMPETING INTERESTS

268

269 The authors have declared that no competing interests exist.

270

#### 271 REFERENCES

272

- 273 1. George, R.A.T. (1985). Vegetable seed production. New York, America. p-318.  
274 2. BBS. (2016). Year book of Agricultural statistics of Bangladesh.2016 28<sup>th</sup> ed.  
275 Bangladesh Bureau of statistics. Statistics Division, Ministry of planning. Govt of the  
276 people Republic of Bangladesh.p-204.  
277 3. Anonymous. (1995). Requirement and supply of seeds. Report of a Task Force,  
278 Ministry of Agriculture, Government of the Peoples Republic of Bangladesh, Dhaka.  
279 p-284.  
280 4. Milford, G.F.J., Armstrong, M.J., Jarvis, P.J., Houghton, B.J., Bellett, D.M., Jones, J.  
281 and Leigh, R.A. (2000). Effects of potassium fertilizer on the yield, quality and

- 282 potassium offtake of sugar beet crops grown on soils of different potassium status.  
283 *J. Agric. Sci.* **135**: 1-10.
- 284 5. Eckert, D. (2010). Efficient Fertilizer Use of Nitrogen. John Willy and Sons, New  
285 York, America. pp. 1-19.
- 286 6. Griffith, B. (2010) Efficient Fertilizer Use - Phosphorus. McMillan Co. Amsterdam,  
287 Netherland. pp. 1-7.
- 288 7. Cantliffe, D. (1992). Nitrate accumulation in vegetable crops as affected by  
289 photoperiod and light duration (beets, radish, spinach, beans). *J. Am. Soc. Hort.*  
290 *Sci.*, **97**: 414-418.
- 291 8. Jaworska, G. (2005) Content of nitrates, nitrites, and oxalates in New Zealand  
292 spinach. *Food Chem.* **89**: 235-242.
- 293 9. Canali, S., Montemurro, F., Tittarelli, F., and Masetti, O. (2008).Effect of nitrogen  
294 fertilization reduction on yield, quality and N utilization of processing spinach. *J.*  
295 *Food. Agric. Environ.* **6**(3&4): 242-247.
- 296 10. Wang, Z. H., Zong, Z.Q., Li, S. X., and Chen, B. M. (2002). Nitrate accumulation in  
297 vegetables and its residual in vegetable fields. *Environ. Sci.* **23**(11): 79-83.
- 298 11. Fadaly, K.A., and Mishriky, J.F. (2000). Effect of nitrogen sources and levels on  
299 growth, yield and mineral composition of spinach. Bulletin of Faculty of Agriculture,  
300 University of Cairo. **41**(3): 973-988.
- 301 12. Assiouty, M.M., Sedera, S.A. and Giza, D. (2005). Effect of bio and chemical  
302 fertilizers on seed production and quality of spinach. *Int. J. Agril.Biol.* **7**(6): 947–952.
- 303 13. Boroujerdnia, M. and Ansari, N.A. (2007).Effect of Different Levels of Nitrogen  
304 Fertilizer and Cultivars on Growth, Yield and Yield Components of Romaine Lettuce  
305 (*Lactuca sativa* L.). *Middle Eastern and Russian J. Plant. Sci. Biotech.* **1**(2): 47-53.
- 306 14. Ambia, K., Hossain, M., Haque, M., Mosiur, A., Halder, C. and Krisna, S. (2016).  
307 Evaluation Seed Yield and Quality of Different Genotypes of Spinach (*Beta*  
308 *vulgaris*). *Sci. Agri.* **16** (2): 61-66.
- 309 15. Zaki, M.S. and Halawa, S.S. (2016). Effect of nitrogen fertilizer and growth  
310 stimulants on flowering behavior, seed yield and its quality of spinach. *Annal. Agric.*  
311 *Sci.* **54**(2): 345-354.