

1
2

Effects of a high dose of vitamin C along with thiamine in critically-ill patients with septic shock: a preliminary study

4
5
6
7

Running title: use of high dose vitamin c in septic shock

Abstract

9

Septic shock may occur in critically ill patients and despite antimicrobial treatment, it is associated with a high mortality rate. It is reasonable to look for new treatment modalities that might improve outcome. This is a randomized, double-blind control trial aiming at critically-ill septic patients in a tertiary hospital. Patients with quick sofa score of 2 and with organ dysfunction were included in this study. The intervention group received high doses of vitamin C at a dose of 50 mg/kg/four times daily along with thiamine at a dose of 200 mg/ twice daily) and the control group received normal saline for four days. The dose of vasopressors, procalcitonin and lactate clearance, and mean sequential organ failure assessment (SOFA) score were examined in the two groups. Patients were followed for 28 days. One hundred patients were allocated into two equal groups, and there was no difference between the two groups regarding baseline characteristics. Mean lactate concentration, SOFA score, days of antimicrobial therapy, and mortality at 28 days were similar between them. However, the mean procalcitonin concentration, and mean vasopressor treatment hours were significantly lower in the intervention group (p<0.05). Although Days of ICU stay were lower in the intervention group, It did not reach statistical significance. The results of this study showed that treatment with high dose vitamin C Reduces the vasopressor requirement without any effects on other parameters. Further studies with larger sample size are required for more generalizable results.

Keywords: Septic Shock, Vitamin C, Organ Failure

27

28

30

INTRODUCTION

32

Sepsis is a fatal complication of infections that causes physiologic, pathologic and biochemical dysfunction with a high mortality rate [1]. Different etiologic factors including bacterial, viral, parasites and fungal have been proposed for sepsis [2]. Host responses, innate immunity, coagulopathy, and inflammatory responses are involved in sepsis pathophysiology [3, 4]. The clinical features of sepsis are variable according to pathogens, infection sources, genetics and health status [3]. Respiratory and cardiovascular systems are affected more frequently [3]. Systemic inflammatory response syndrome (SIRS) and quick sequential organ failure assessment (qSOFA) were proposed for initial assessment of patients [2]. Previously Sepsis was defined as the occurrence of at least two of SIRS criteria along with confirmed or suspected infection [5]. The most recent revised definition has eliminated SIRS criteria and rely on sequential organ dysfunction as results of infections [1]. Recently, the third international consensus on sepsis definition recommend qSOFA for initial evaluation [1]. Quick SOFA defined as certain or suspected infection along with at least two of the following criteria: decline in level of consciousness, Systolic blood pressure < 100 mm Hg, respiratory rate > 22/min.

47

Septic shock is circulatory, cellular and metabolic dysfunctions in patients with sepsis. It is defined as mean arterial pressure \leq 65mmHg and serum lactate levels \geq 2 milimol despite adequate fluid resuscitation (1, 4). Different etiologic factors such as adrenal insufficiency, creation of reactive oxygen species, mitochondrial dysfunction, and vasopressin deficiency have been proposed for septic shock (6). Previous studies showed that antioxidant supplements may improve organ dysfunction, dose of vasopressors and mortality in septic shock [7-9].

54

Vitamin C is a water soluble acting as a free radical scavenger that may improve endothelial function, and micro- and macrovascular functions [10]. In addition, a decrease in serum ascorbate levels has been reported in previous studies after intensive care admission [8, 11, 12]. Previous studies showed the safety and efficacy of vitamin C doses at 25-50 mg/kg four times daily [7, 9]. Increased oxalate concentration seen at daily doses of more than 10 grams [13]. Thiamine deficiency is prevalent in sepsis [14, 15]. Thiamine is a key vitamin in cellular metabolism and improve Krebs cycle, aerobic respiration, branched-chain amino acids [16]. Therefore, the aim

of the present study is to evaluate the possible role of high-dose vitamin C along with thiamine as compared with placebo in critically-ill patients with septic shock according to the recommendations of Society of Critical Care Medicine guidelines [17].

65

METHOD

This study was a randomized, single blind controlled trial conducted in a tertiary hospital in Kermanshah. Patients were allocated in two groups according to the random number generation software. Quick sofa was used for initial examination of patients. Patients with qSOFA of 2 and more were examined for possible sepsis. Other examinations such as Para-clinic data (serum creatinine, liver function test, white blood cells), chest X-ray, and cultures (sputum, blood, spinal fluid and urine as indicated) were performed too.

73

Patients

One hundred Patients including 57 women who were diagnosed with septic shock and signed the informed consent (by patients or their next of kin) were included in the study. Pregnant patients, children and adolescents, and those with hypersensitivity to ascorbic acid, glucose-6 phosphate dehydrogenase deficiency, history of nephrolithiasis and immunosuppression were excluded from the study. Patients in the intervention group received high dose of vitamin C (50 mg/kg four times daily up to 6g/day) for 4 consecutive days along with thiamine 200 mg and patients in the placebo received normal saline. Other diagnostics and treatment modalities were similar for the two groups. All included patients with septic shock received norepinephrine at the initial dosage of 4-20 micrograms/minute, to sustain mean arterial pressure of more than 65 mm Hg. Furthermore, all patients received Hydrocortisone at the daily dose of 200 mg was started at time of norepinephrine initiation, continued for four days and then tapered over four days.

87

88

Measurements

During the study, arterial oxygen pressure (PaO₂) to the fraction of inspired oxygen (FiO₂), systolic and diastolic hypotension, and renal and hepatic functions were tested daily. Sequential Organ Failure Assessment (SOFA) [18] and Acute Physiology and Chronic Health Evaluation

75

(APACHE II) were recorded [19]. Acute Kidney Injury Network (AKIN) criteria were used for kidney function evaluation [20]. All patients, if survived, were followed for 28 days.

The mean concentration of lactate, procalcitonin at baseline and after 72 hours were recorded for all allocated patients. Additionally, Mean hours of vasopressor therapy, total days of mechanical ventilation, ICU stay, and mortality at 28 days were compared between the two groups.

98

99

Data Analysis

To perform the data analysis, we used the software SPSS-20. T-test was used for quantitative data and categorical data analysis was performed with Chi-squared test or Fisher as appropriate.

Wilcoxon and the Mann-Whitney-U tests were used.

104

105

106

RESULTS

108

During the study, 100 patients in two equal groups of fifty patients with mean age of 61.06 ± 16.02 years were included in the study. Demographic data are shown in table.1 where baseline characteristics were similar in the two groups. Chronic obstructive pulmonary disease (COPD) was the main reason for the admission of patients, and other baseline diseases are demonstrated in table 1. Pneumonia was the main source for sepsis and septic shock, which was seen in 65% of all participants followed by intra-abdominal (15 %), and meningitis sources (10%) respectively. None of patients received hormonal therapy. No definite source was detected in 10% of our patients. Mean SOFA score after 96 hours were similar between two groups (9.01 ± 3.92 vs. 10.67 ± 3.03 , for intervention and control groups respectively, $p=0.204$). There was no significant difference between intervention and control groups regarding mortality after 28 days (15 % vs. 17 % respectively, $p=0.36$). In our study, mean hours of vasopressor therapy was 77.52 ± 21.5 and 111.5 ± 15.75 hours in intervention and control groups respectively ($p=0.001$).

The results of the present study did not show significant differences between the two groups regarding days of ICU stay 9.87 ± 8.32 vs. 12.67 ± 6.99 ($p=0.32$), and days patients needed mechanical ventilation 6.67 ± 7.84 vs. 7.42 ± 6.23 ($p=0.87$). In addition, there was not any significant difference between the two groups regarding serum lactate after 72 hours 4.1 ± 2.6 vs. 5.1 ± 3.2 milimol ($p=0.65$). Although, a similar procalcitonin concentration was detected at baseline; however, mean serum procalcitonin concentration was significantly lower in the intervention group 2.21 ± 1.5 vs. 4.13 ± 2.31 ($p=0.001$) after 72 hours. No significant adverse event was seen in our patients.

132

133

DISCUSSION

135

In the present study, we evaluated the safety and efficacy of vitamin C in critically-ill patients with septic shock. High doses of vitamin C was well tolerated and associated with lower duration of vasopressor requirement. As known, critically-ill patients are at a high risk of antioxidants concentration [21]. *Fowler et al* studied the safety of high dose (200mg/kg/ day) for four days, in septic shock, where treatment with high dose of vitamin C significantly increased plasma vitamin C concentrations without significant adverse events [8].

In our study, mean hours of vasopressor therapy was 77.52 ± 21.5 and 111.5 ± 15.75 hours in intervention and control groups respectively ($p=0.001$). This data is concordant with *Marik et al* [7]. They evaluated the effect of early use of high dose vitamin C along with thiamine and hydrocortisone can decrease the duration of vasopressor therapy (18.3 ± 9.8 vs. 54.9 ± 28.4 hours respectively), organ dysfunction, and mortality. In addition, *zabet et al* reported lower mortality in patients who treated with high doses of vitamin C [9]. However, the mortality rate in our study didn't reduce in the intervention group. Results of our study is similar to *Litwak et al.* that showed treatment with marik's cocktail didn't significantly change outcomes (including mortality) in critically ill patients with septic shock compared with standard cares [22].

151

Mean duration of mechanical ventilation was not significantly lower in the intervention group. In the original study by *Marik* and associates vitamin C, thiamine, and hydrocortisone cocktail

was associated with lower duration of mechanical ventilation [9]. Our data are consistent with Shin *et al.* study that showed treatment with mentioned protocol versus control didn't affect duration of mechanical ventilation (5 (IQR, 3.0–15.0) vs. 5.0 (IQR, 3.0–10.0) days, $p= 0.63$) [23].

158

Duration of ICU stay was similar between the two groups. Shin *et al.* and Litwak *et al.* that separately showed treatment with this protocol didn't decrease duration of ICU stay [22, 23]. However, Previous studies showed that treatment with vitamins E and C may decrease organ failure as measured by the SOFA score, hospital but not ICU stay in critically-ill patients [8, 24].

163

Patients treated with Thiamine and vitamin C had similar serum lactate clearance after 72 hours. Serum lactate levels and clearance may both be useful predictors of mortality in critically-ill patients [25]. Recently, a single center retrospective cohort study showed that treatment with a high dose of intravenous thiamine (1500 mg/day) within 24 hours is associated with an increase in lactate clearance and decrease 28 days mortality [26]. As mentioned in our study, only low dose of thiamine along with high dose of vitamin C was used.

171

Although mean procalcitonin was similar in the two groups at baseline, the intervention group had significantly lower procalcitonin concentration. It confirms the previous study by Fowler *et al.* showing that treatment with a high dose of vitamin C decreases inflammatory biomarkers, including c-reactive protein and procalcitonin concentration [8]. They demonstrated that vitamin C infusion may decrease procalcitonin concentration as high as 50%. Therefore, clinicians should interpret procalcitonin concentration with caution.

179

180

Limitation

Our study had several limitations: Low sample size in both groups; Not being able to determine daily serum level of ascorbate. Although primary determined sample size size was 15 patients in each group, however, 100 patients were followed and included for the final analysis.

185

186

CONCLUSION

Results showed that treatment with a high dose vitamin C is well tolerated in critically-ill septic patients and may decrease total duration of vasopressor therapy and procalcitonin concentration without affecting the duration of ICU stay and one month mortality. Further studies with larger sample size is needed to confirm the possible role of high-dose vitamin C in similar populations.

192

Conflict of interest: The authors have declared that there is no conflict of interest.

194

ETHICAL APPROVAL

This study was approved by the Ethics Committee of Kermanshah University of Medical Sciences (IR.KUMS.REC.1396.341) and registered in IRCT (IRCT20130812014333N78).

195

198

199

200

References:

202

- 203 Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The third international consensus definitions for sepsis and septic shock (Sepsis-3). *JAMA*. 2016;315(8):801-10.
- 206 Angus DC, Van der Poll T. Severe sepsis and septic shock. *N Engl J Med*. 2013;369(9):840-51.
- 208 Angus DC, Van der Poll T. Severe sepsis and septic shock. *N Engl J Med*. 2013;369(9):840-51.
- 210 Gotts JE, Matthay MA. Sepsis: pathophysiology and clinical management. *BMJ*. 2016;353:i1585.
- 212 Levy MM, Fink MP, Marshall JC, Abraham E, Angus D, Cook D, et al. 2001 SCCM/ESICM/ACCP/ATS/SIS international sepsis definitions conference. *Intensive Care Med*. 2003;29(4):530-8.

6. 215 Landry DW, Oliver JA. The pathogenesis of vasodilatory shock. *NEJM* 2001;345(8):588-95. 216
7. 217 Marik PE, Khangoora V, Rivera R, Hooper MH, Catravas J. Hydrocortisone, vitamin C, and thiamine for the treatment of severe sepsis and septic shock: a retrospective before-after study. *Chest*. 2017;151(6):1229-1238. 218
8. 220 Syed AA, Knowlson S, Sculthorpe R, Farthing D, DeWilde C, Farthing CA, et al. Phase I safety trial of intravenous ascorbic acid in patients with severe sepsis. *J Transl Med*. 2014 31;12:32. 219
9. 223 Zabet MH, Mohammadi M, Ramezani M, Khalili HJ. Effect of high-dose Ascorbic acid on vasopressor's requirement in septic shock. *J Res Pharm Pract*. 2016; 5(2): 94–100. 220
10. 225 Moskowitz A, Andersen LW, Huang DT, Berg KM, Grossestreuer AV, Marik PE, et al. Ascorbic acid, corticosteroids, and thiamine in sepsis: a review of the biologic rationale and the present state of clinical evaluation. *Crit Care*. 2018;22(1):283. 221
11. 228 Oudemans-van Straaten HM, Spoelstra-de Man AM, de Waard MC. Vitamin C revisited. *Crit Care*. 2014;18(4):460 222
12. 230 Carr AC, Shaw GM, Natarajan R. Ascorbate-dependent vasopressor synthesis: a rationale for vitamin C administration in severe sepsis and septic shock? *Crit Care*. 2015; 19: 418. 223
13. 232 de Grooth H-J, Manubulu-Choo W-P, Zandvliet AS, Spoelstra-de Man AM, Girbes AR, Swank EL, et al. Vitamin C pharmacokinetics in critically ill patients: a randomized trial of four IV regimens. *Chest*. 2018;153(6):1368-1377. 224
14. 235 Costa NA, Gut AL, de Souza Dorna M, Pimentel JAC, Cozzolino SMF, Azevedo PS, et al. Serum thiamine concentration and oxidative stress as predictors of mortality in patients with septic shock. *Journal of critical care. J Crit Care*. 2014;29(2):249-52. 225
15. 238 Donnino MW, Carney E, Cocchi MN, Barbash I, Chase M, Joyce N, et al. Thiamine deficiency in critically ill patients with sepsis. *J Crit Care*. 2010;25(4):576-81. 226
16. 240 Mallat J, Lemyze M, Thevenin D. Do not forget to give thiamine to your septic shock patient! *Journal of thoracic disease. J Thorac Dis*. 2016; 8(6): 1062–1066. 227
17. 242 Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving sepsis campaign: international guidelines for management of sepsis and septic shock: 2016. *Intensive Care Med*. 2017;43(3):304-377 228

18. 245 Vincent J-L, Moreno R, Takala J, Willatts S, De Mendonça A, Bruining H, et al. The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. Intensive care medicine. Intensive Care Med. 1996;22(7):707-10..
19. 248 Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. Crit Care Med. 1985 Oct;13(10):818-29.
20. 250 Thomas ME, Blaine C, Dawnay A, Devonald MA, Ftouh S, Laing C, et al. The definition of acute kidney injury and its use in practice. Kidney Int. 2015;87(1):62-73.
21. 252 Carr AC, Rosengrave PC, Bayer S, Chambers S, Mehrrens J, Shaw GMJCC. Hypovitaminosis C and vitamin C deficiency in critically ill patients despite recommended enteral and parenteral intakes. Crit Care. 2017;21(1):300.
22. 254 Litwak JJ, Cho N, Nguyen HB, Moussavi K, Bushell T. Vitamin C, Hydrocortisone, and Thiamine for the Treatment of Severe Sepsis and Septic Shock: A Retrospective Analysis of Real-World Application. Journal of clinical medicine. 2019;8(4):478.
23. 258 Shin TG, Kim YJ, Ryoo SM, Hwang SY, Jo IJ, Chung SP, Choi SH, et al. Early vitamin C and thiamine administration to patients with septic shock in emergency departments: Propensity score-based analysis of a before-and-after cohort study. Journal of clinical medicine. 2019;8(1):102.
24. 262 Nathens AB, Neff MJ, Jurkovich GJ, Klotz P, Farver K, Ruzinski JT, et al. Randomized, prospective trial of antioxidant supplementation in critically ill surgical patients. Ann Surg. 2002;136(6):814-22.
25. 265 Ryoo SM, Lee J, Lee Y-S, Lee JH, Lim KS, Huh JW, et al. Lactate level versus lactate clearance for predicting mortality in patients with septic shock defined by Sepsis-3. Crit Care Med. 2018;46(6):e489-e495.
26. 268 Woolum JA, Abner EL, Kelly A, Bastin MLT, Morris PE, Flannery AH. Effect of thiamine administration on lactate clearance and mortality in patients with septic shock. Crit Care Med. 2018;46(11):1747-1752

271

272

273

274

275

276

277

278

279

Table 1 Baseline characteristics of study population


	Intervention group	Placebo	P
Age (years)	56.21 ±13.62	61.07± 16.85	0.8
Female	27 (54%)	30 (60%)	0.7
Body mass index (kg/m2)	25.75±2.29	25.43±2.43	0.43
Mean GFR (milliliter/min)	55±10	60±13	0.6
Mean SOFA score	10.2±2.5	9.8±3.8	0.32
Mean procalcitonin level (ng/ml)	1.25±1.6	1.99±1.56	0.146
Underling diseases / Reason for admission			0.75
COPD exacerbation	30	35	
Diabetes	20	15	
Guillan barre	2	0	
Chronic renal failure	17	20	
Hypertension	10	8	
Secondary Peritonintis	15	13	
Systemic lupus erythematosus	3	0	
Cerebrovascular accident	15	10	
Heart failure	10	7	
Epilepsy	3	1	

CO~~PD~~: choronic obstructive pulmonary disease,

282

283

284



287

Figure 288 Mean procacitonin concentration before and end of intervention (p<0.0001)

286

UNDER PBL